

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
30 October 2003 (30.10.2003)

PCT

(10) International Publication Number  
**WO 03/088908 A2**

- (51) International Patent Classification<sup>7</sup>: **A61K** Circle, Morrisville, NC 27560 (US). **GROSS, Michael, F.** [US/US]; 6200 Chesden Dr., Durham, NC 27713 (US).
- (21) International Application Number: PCT/US03/11807
- (22) International Filing Date: 16 April 2003 (16.04.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/374,279 19 April 2002 (19.04.2002) US
- (71) Applicants (for all designated States except US): **BRISTOL-MYERS SQUIBB COMPANY** [US/US]; P. O. Box 4000, Route 206 and Provinceline Road, Princeton, NJ 08543-4000 (US). **ICAGEN, INC.** [US/US]; 4222 Emperor Boulevard, Suite 350, Durham, NC 27703 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **LLOYD, John** [US/US]; 1702 Yardley Rd., Yardley, PA 19067 (US). **JEON, Yoon, T.** [US/US]; 10 Saddlewood Ct., Belle Mead, NJ 08502 (US). **FINLAY, Heather** [GB/US]; 363 Province Line Rd., Skillman, NJ 08558 (US). **YAN, Lin** [US/US]; 4 Schindler Ct., East Brunswick, NJ 08816 (US). **BEAUDOIN, Serge** [CA/US]; 2215 Apt. Duck Pond
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: HETEROCYCLO INHIBITORS OF POTASSIUM CHANNEL FUNCTION

(57) Abstract: Novel heterocyclo compounds useful as inhibitors of potassium channel function (especially inhibitors of the Kv1 subfamily of voltage gated K<sup>+</sup> channels, especially inhibitors Kv1.5 which has been linked to the ultra-rapidly activating delayed rectifier K<sup>+</sup> current IK<sub>ur</sub>), methods of using such compounds in the prevention and treatment of arrhythmia and IK<sub>ur</sub>-associated conditions, and pharmaceutical compositions containing such compounds.

WO 03/088908 A2

## HETEROCYCLO INHIBITORS OF POTASSIUM CHANNEL FUNCTION

### FIELD OF THE INVENTION

The present invention provides for heterocyclyl compounds useful as  
5 inhibitors of potassium channel function (especially inhibitors of the  $K_v1$  subfamily of voltage gated  $K^+$  channels, more especially inhibitors  $K_v1.5$  which has been linked to the ultra-rapidly activating delayed rectifier  $K^+$  current  $I_{Kur}$ ) and to pharmaceutical compositions containing such compounds. The present invention further provides for methods of using such compounds in the treatment of arrhythmia,  $I_{Kur}$ -associated  
10 disorders, and other disorders mediated by ion channel function.

### BACKGROUND OF THE INVENTION

The importance of potassium channels was first recognized approximately fifty years ago when Hodgkin and Huxley discovered that potassium ions contributed  
15 to the current that excited the squid giant axon. Research in the area, however, was hampered by the lack of selective, high affinity ligands for potassium channels. But the advent of recombinant DNA techniques and single cell and whole cell voltage clamp techniques has changed the slow pace of the field. Indeed, potassium channels that exhibit functional, pharmacological and tissue distribution characteristics have  
20 been cloned. These cloned potassium channels are useful targets in assays for identifying candidate compounds for the treatment of various disease states. Potassium channels have turned out to be the most diverse family of ion channels discovered to date. They modulate a number of cellular events such as muscle contraction, neuro-endocrine secretion, frequency and duration of action potentials,  
25 electrolyte homeostasis, and resting membrane potential.

Potassium channels are expressed in eukaryotic and procaryotic cells and are elements in the control of electrical and non-electrical cellular functions. Potassium channels have been classified according to their biophysical and pharmacological characteristics. Subclasses of these channels have been named based on amino acid  
30 sequence and functional properties. Salient among these are the voltage dependent potassium channels, for example voltage gated potassium channels (e.g.,  $K_v1$ ,  $K_v2$ ,  $K_v3$ ,  $K_v4$ ). Subtypes within these subclasses have been characterized as to their

putative function, pharmacology and distribution in cells and tissues (Chandy and Gutman, "Voltage-gated potassium channel genes" in Handbook of Receptors and Channels – Ligand and Voltage-gated Ion Channels, ed. R.A. North, 1995; Doupnik et al., Curr. Opin. Neurobiol. 5:268, 1995). For example, the K<sub>v</sub>1 class of potassium channels is further subdivided depending on the molecular sequence of the channel, for example K<sub>v</sub>1.1, K<sub>v</sub>1.2, K<sub>v</sub>1.3, K<sub>v</sub>1.4, K<sub>v</sub>1.5, K<sub>v</sub>1.6, and K<sub>v</sub>1.7. Functional voltage-gated K<sup>+</sup> channels can exist as multimeric structures formed by the association of either identical or dissimilar subunits. This phenomena is thought to account for the wide diversity of K<sup>+</sup> channels. However, subunit compositions of native K<sup>+</sup> channels and the physiologic role that particular channels play are, in most cases, still unclear.

Membrane depolarization by K<sub>v</sub>1.3 inhibition has been shown to be an effective method to prevent T-cell proliferation and therefore has applications in many autoimmune conditions. Inhibition of K<sup>+</sup> channels in the plasma membrane of human T-lymphocytes has been postulated to play a role in eliciting immunosuppressive responses by regulating intracellular Ca<sup>++</sup> homeostasis, which has been found to be important in T-cell activation.

The K<sub>v</sub>1.3 voltage-gated potassium channel is found in neurons, blood cells, osteoclasts and T-lymphocytes. The Chandy and Cahalan laboratories proposed a hypothesis that blocking the K<sub>v</sub>1.3 channel would elicit an immunosuppressant response. (Chandy et al., J. Exp. Med. 160, 369, 1984; Decoursey et al., Nature, 307, 465, 1984). However, the K<sup>+</sup> channel blockers employed in their studies were non-selective. Until research with the peptide margatoxin, a peptide found in scorpion venom, no specific inhibitor of the K<sub>v</sub>1.3 channel existed to test this hypothesis. Although a laboratory (Price et al., Proc. Natl. Acad. Sci. USA, 86, 10171, 1989) showed that charybdotoxin would block K<sub>v</sub>1.3 in human T-cells, charybdotoxin was subsequently shown to inhibit four different K<sup>+</sup> channels (K<sub>v</sub>1.3 and three distinct small conductance Ca<sup>++</sup> activated K<sup>+</sup> channels) in human T-lymphocytes, limiting the use of this toxin as a probe for the physiological role of K<sub>v</sub>1.3 (Leonard et al., Proc. Natl. Acad. Sci. USA, 89, 10094, 1992). Margatoxin, on the other hand, blocks only K<sub>v</sub>1.3 in T-cells, and has immunosuppressant activity on both in in vitro and in vivo models. (Lin et al., J. exp. Med, 177, 637, 1993). The therapeutic utility of this compound, however, is limited by its potent toxicity. Recently, a class of compounds

has been reported that may be an attractive alternative to the above mentioned drugs, see for example U.S. Patent Nos. 5,670,504; 5,631,282; 5,696,156; 5,679,705; and 5,696,156. While addressing some of the activity/toxicity problems of previous drugs, these compounds tend to be of large molecular weight and are generally produced by synthetic manipulation of a natural product, isolation of which is cumbersome and labor intensive.

Immunoregulatory abnormalities have been shown to exist in a wide variety of autoimmune and chronic inflammatory diseases, including systemic lupus erythematosus, chronic rheumatoid arthritis, type I and II diabetes mellitus, inflammatory bowel disease, biliary cirrhosis, uveitis, multiple sclerosis and other disorders such as Crohn's disease, ulcerative colitis, bullous pemphigoid, sarcoidosis, psoriasis, ichthyosis, Graves ophthalmopathy and asthma.

Although the underlying pathogenesis of each of these conditions may be quite different, they have in common the appearance of a variety of auto-antibodies and self-reactive lymphocytes. Such self-reactivity may be due, in part, to a loss of the homeostatic controls under which the normal immune system operates. Similarly, following a bone-marrow or an organ transplantation, the host lymphocytes recognize the foreign tissue antigens and begin to produce antibodies which lead to graft rejection.

One end result of an autoimmune or a rejection process is tissue destruction caused by inflammatory cells and the mediators they release. Anti-inflammatory agents such as NSAID's act principally by blocking the effect or secretion of these mediators but do nothing to modify the immunologic basis of the disease. On the other hand, cytotoxic agents, such as cyclophosphamide, act in such a nonspecific fashion that both the normal and autoimmune responses are shut off. Indeed, patients treated with such nonspecific immunosuppressive agents are as likely to succumb from infection as they are from their autoimmune disease.

Cyclosporin A (CsA), which was approved by the US FDA in 1983 is currently the leading drug used to prevent rejection of transplanted organs. In 1993, FK-506 (Prograf) was approved by the US FDA for the prevention of rejection in liver transplantation. CsA and FK-506 act by inhibiting the body's immune system from mobilizing its vast arsenal of natural protecting agents to reject the transplant's

foreign protein. In 1994, CsA was approved by the US FDA for the treatment of severe psoriasis and has been approved by European regulatory agencies for the treatment of atopic dermatitis. Though they are effective in fighting transplant rejection, CsA and FK-506 are known to cause several undesirable side effects including nephrotoxicity, neurotoxicity, and gastrointestinal discomfort. Therefore, a selective immunosuppressant without these side effects still remains to be developed. Potassium channel inhibitors promise to be the solution to this problem.

Atrial fibrillation (AF) and atrial flutter are the most common cardiac arrhythmias in clinical practice and are likely to increase in prevalence with the aging of the population. Currently, AF affects more than 1 million Americans annually, represents over 5% of all admissions for cardiovascular diseases and causes more than 80,000 strokes each year in the United States. While AF is rarely a lethal arrhythmia, it is responsible for substantial morbidity and can lead to complications such as the development of congestive heart failure or thromboembolism. Currently available Class I and Class III antiarrhythmic drugs reduce the rate of recurrence of AF, but are of limited use because of a variety of potentially adverse effects including ventricular proarrhythmia. Because current therapy is inadequate and fraught with side effects, there is a clear need to develop new therapeutic approaches.

Antiarrhythmic agents of Class III are drugs that cause a selective prolongation of the duration of the action potential without significant cardiac depression. Available drugs in this class are limited in number. Examples such as sotalol and amiodarone have been shown to possess interesting Class III properties (Singh B.N., Vaughan Williams E.M. "A Third Class of Anti-Arrhythmic Action: Effects On Atrial And Ventricular Intracellular Potentials And Other Pharmacological Actions On Cardiac Muscle, of MJ 1999 and AH 3747" Br. J. Pharmacol 1970; 39:675-689. and Singh B.N., Vaughan Williams E.M., "The Effect of Amiodarone, A New Anti-Anginal Drug, On Cardiac Muscle", Br J. Pharmacol 1970; 39:657-667), but these are not selective Class III agents. Sotalol also possesses Class II effects which may cause cardiac depression and is contraindicated in certain susceptible patients. Amiodarone, also is not a selective Class III antiarrhythmic agent because it possesses multiple electrophysiological actions and is severely limited by side effects (Nademanee, K. "The Amiodarone Odessey". J. Am. Coll. Cardiol. 1992;20:1063-1065.) Drugs of

this class are expected to be effective in preventing ventricular fibrillation. Selective class III agents, by definition, are not considered to cause myocardial depression or an induction of arrhythmias due to inhibition of conduction of the action potential as seen with Class I antiarrhythmic agents.

- 5           Class III agents increase myocardial refractoriness via a prolongation of cardiac action potential duration. Theoretically, prolongation of the cardiac action potential can be achieved by enhancing inward currents (i.e.  $\text{Na}^+$  or  $\text{Ca}^{2+}$  currents; hereinafter  $I_{\text{Na}}$  and  $I_{\text{Ca}}$ , respectively) or by reducing outward repolarizing potassium ( $\text{K}^+$ ) currents. The delayed rectifier ( $I_{\text{K}}$ )  $\text{K}^+$  current is the main outward current
- 10 involved in the overall repolarization process during the action potential plateau, whereas the transient outward ( $I_{\text{to}}$ ) and inward rectifier ( $I_{\text{K1}}$ )  $\text{K}^+$  currents are responsible for the rapid initial and terminal phases of repolarization, respectively. Cellular electrophysiologic studies have demonstrated that  $I_{\text{K}}$  consists of two pharmacologically and kinetically distinct  $\text{K}^+$  current subtypes,  $I_{\text{Kr}}$  (rapidly activating
- 15 and deactivating) and  $I_{\text{Ks}}$  (slowly activating and deactivating) (Sanguinetti and Jurkiewicz, Two Components Of Cardiac Delayed Rectifier  $\text{K}^+$  Current: Differential Sensitivity To Block By Class III Antiarrhythmic Agents, J Gen Physiol 1990, 96:195-215). Class III antiarrhythmic agents currently in development, including d-sotalol, dofetilide (UK-68,798), almokalant (H234/09), E-4031 and
- 20 methanesulfonamide-N-[1'-6-cyano-1,2,3,4-tetrahydro-2-naphthalenyl]-3,4-dihydro-4-hydroxyspiro[2H-1-benzopyran-2,4'-piperidin]-6yl]monochloride, predominantly, if not exclusively, block  $I_{\text{Kr}}$ . Although, amiodarone is a blocker of  $I_{\text{Ks}}$  (Balser J.R. Bennett, P.B., Hondeghem, L.M. and Roden, D.M. "Suppression Of Time-Dependent Outward Current In Guinea Pig Ventricular Myocytes: Actions Of Quinidine And
- 25 Amiodarone. Circ. Res. 1991, 69:519-529), it also blocks  $I_{\text{Na}}$  and  $I_{\text{Ca}}$ , effects thyroid function, is as a nonspecific adrenergic blocker, and acts as an inhibitor of the enzyme phospholipase (Nademanee, K. "The Amiodarone Odessey" J.Am. Coll. Cardiol.1992;20:1063-1065). Therefore its method of treating arrhythmia is uncertain. Most Class III agents that are known to be in development predominantly
- 30 block  $I_{\text{Kr}}$ .

Reentrant excitation (reentry) has been shown to be a prominent mechanism underlying supraventricular arrhythmias in man. Reentrant excitation requires a

critical balance between slow conduction velocity and sufficiently brief refractory periods to allow for the initiation and maintenance of multiple reentry circuits to coexist simultaneously and sustain AF. Increasing myocardial refractoriness by prolonging action potential duration (APD), prevents and/or terminates reentrant  
5 arrhythmias. Most selective Class III antiarrhythmic agents currently in development, such as d-sotalol and dofetilide predominantly, if not exclusively, block  $I_{kr}$ , the rapidly activating component of  $I_K$  found both in the human atrium and ventricle.

Since these  $I_{kr}$  blockers increase APD and refractoriness both in atria and ventricle without affecting conduction per se, theoretically they represent potential  
10 useful agents for the treatment of arrhythmias like AF. These agents have a liability in that they have an enhanced risk of proarrhythmia at slow heart rates. For example, torsades de points has been observed when these compounds are utilized (Roden, D.M. "Current Status of Class III Antiarrhythmic Drug Therapy", Am J. Cardiol, 1993; 72:44B-49B). This exaggerated effect at slow heart rates has been termed  
15 "reverse frequency-dependence", and is in contrast to frequency-independent or frequency-dependent actions (Hondeghe, L.M. "Development of Class III Antiarrhythmic Agents". J.Cadiovasc.Cardiol. 20 (Suppl.2):S17-S22).

The slowly activating component of the delayed rectifier ( $I_{ks}$ ) potentially overcomes some of the limitations of  $I_{kr}$  blockers associated with ventricular  
20 arrhythmias. Because of its slow activation kinetics however, the role of  $I_{ks}$  in atrial repolarization may be limited due to the relatively short APD of the atrium. Consequently, although  $I_{ks}$  blockers may provide distinct advantage in the case of ventricular arrhythmias, their ability to affect SVT is considered to be minimal.

The ultra-rapidly activating delayed rectifier  $K^+$  current ( $I_{kur}$ ) is believed to  
25 represent the native counterpart to a cloned potassium channel designated Kv1.5 and, while present in human atrium, it appears to be absent in human ventricle. Furthermore, because of its rapidity of activation and limited slow inactivation,  $I_{kur}$  is believed to contribute significantly to repolarization in human atrium. Consequently, a specific blocker of  $I_{kur}$ , that is a compound which blocks Kv1.5, would overcome  
30 the short coming of other compounds by prolonging refractoriness by retarding repolarization in the human atrium without causing the delays in ventricular

reporlarization that underlie arrhythmogenic after depolarizations and acquired long QT syndrome observed during treatment with current Class III drugs.


In intact human atrial myocytes an ultra-rapidly activating delayed rectifier  $K^+$  current  $I_{Kur}$  which is also known as the sustained outward current,  $I_{sus}$  or  $I_{so}$ , has been identified and this current has properties and kinetics identical to those expressed by the human  $K^+$  channel clone (hKv1.5, HK2) when isolated from human heart and stably expressed in human (HEK-293) cell lines (Wang et al., 1993, Circ Res 73:1061-1076; Fedida et al., 1993, Circ Res 73:210-216; Snyders et al., 1993, J Gen Physiol 101:513-543) and originally cloned from rat brain (Swanson et al., 1990, Neuron 4:929-939). Although various antiarrhythmic agents are now available on the market, those having both satisfactory efficacy and a high margin of safety have not been obtained. For example, antiarrhythmic agents of Class I according to the classification scheme of Vaughan-Williams ("Classification Of Antiarrhythmic Drugs: In: Cardiac Arrhythmias, edited by: E. Sandoe, E. Flensted-Jensen, K. Olesen; Sweden, Astra, Sodertalje, pp449-472, 1981) which cause a selective inhibition of the maximum velocity of the upstroke of the action potential ( $V_{max}$ ) are inadequate for preventing ventricular fibrillation. In addition, they have problems regarding safety, namely, they cause a depression of myocardial contractility and have a tendency to induce arrhythmias due to an inhibition of impulse conduction. Beta-adrenoceptor blockers and calcium antagonists which belong to Class II and IV, respectively, have a defect in that their effects are either limited to a certain type of arrhythmia or are contraindicated because of their cardiac depressant properties in certain patients with cardiovascular disease. Their safety, however, is higher than that of the antiarrhythmic agents of Class I.

25

### SUMMARY OF THE INVENTION

The present invention provides heterocycl compounds of the following formula I, including enantiomers, diastereomers, and salts thereof, useful as inhibitors of potassium channel function (especially inhibitors of the  $K_v1$  subfamily of voltage gated  $K^+$  channels, more especially inhibitors of  $K_v1.5$  which has been linked to the ultra-rapidly activating delayed rectifier  $K^+$  current,  $I_{Kur}$ ) for the treatment of disorders such as arrhythmia and  $I_{Kur}$ -associated disorders:

30




including enantiomers, diastereomers and salts thereof wherein



5 m and p are independently 0, 1, 2 or 3 provided that the sum of m and p is at least 2;

Q is  $\text{NR}^1$ , O, S, S(O) or S(O)<sub>2</sub>;

R<sup>1</sup> is



10



15

$\text{---C(=NR}^{8b})\text{R}^{8c}$ ,  $\text{---SO}_2\text{R}^{8c}$ ,  $\text{---OC(O)CCl}_3$ ,  $\text{---C(=S)R}^{8c}$ , optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclo, perfluoroalkyl, cyano, hydroxy, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted alkenyl, or optionally substituted alkynyl;

20

R<sup>2</sup> is heteroaryl, (heteroaryl)alkyl, aryl, (aryl)alkyl, heterocyclo, (heterocyclo)alkyl, alkyl or cycloalkyl, any of which may be optionally independently substituted with one or more groups T<sup>1</sup>, T<sup>2</sup> or T<sup>3</sup>;


J is a bond, C<sub>1-4</sub> alkylene optionally independently substituted with one or more

groups T<sup>1a</sup>, T<sup>2a</sup> or T<sup>3a</sup>, or C<sub>1-4</sub> alkenylene optionally independently substituted

25

with one or more groups T<sup>1a</sup>, T<sup>2a</sup> or T<sup>3a</sup>;

R<sup>3</sup> is



$R^4$  is H, alkyl, haloalkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, (aryl)alkyl or heteroaryl any of which may be optionally independently substituted with one or more groups  $T^{1b}$ ,  $T^{2b}$  or  $T^{3b}$ ;

$R^5$  is

- (a)  $\text{---NR}^{6a}\text{R}^{7a}$ , cyano or  
 (b) heteroaryl, (heteroaryl)alkyl, aryl, (aryl)alkyl, alkyl, cycloalkyl, (cycloalkyl)alkyl, heterocyclo, (heterocyclo)alkyl, or alkyl any of which may be optionally independently substituted with one or more groups  $T^{1c}$ ,  $T^{2c}$  or  $T^{3c}$ ;

$R^6$ ,  $R^{6a}$ ,  $R^7$ ,  $R^{7a}$ ,  $R^8$ ,  $R^{8a}$ ,  $R^{8a1}$ ,  $R^{8a2}$ , and  $R^{8a3}$  are independently H, alkyl, hydroxy, alkoxy, aryloxy, heterocycloxy, heteroaryloxy, (hydroxy)alkyl, (alkoxy)alkyl, (aryloxy)alkyl, (heterocycloxy)alkyl, (heteroaryloxy)alkyl, (cyano)alkyl, (alkenyl)alkyl, (alkynyl)alkyl, cycloalkyl, (cycloalkyl)alkyl, aryl, (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, heterocyclo, (heterocyclo)alkyl,  $\text{---C(O)R}^9$ ,

- CO<sub>2</sub>R<sup>9</sup>, -C(O)-NR<sup>9</sup>R<sup>10</sup>, or -NR<sup>9</sup>R<sup>10</sup> any of which may be optionally independently substituted with one or more groups T<sup>1d</sup>, T<sup>2d</sup> or T<sup>3d</sup>;  
 or R<sup>6</sup> and R<sup>7</sup>, or R<sup>6a</sup> and R<sup>7a</sup> together with the nitrogen atom to which they are attached may combine to form a 4 to 8 membered heterocyclo ring optionally  
 5 independently substituted with one or more groups T<sup>1d</sup>, T<sup>2d</sup> or T<sup>3d</sup>;  
 or one of R<sup>6</sup> or R<sup>7</sup>, may combine with one of R<sup>8</sup>, R<sup>8a</sup> or R<sup>9</sup> to form a saturated or unsaturated 5 to 8 membered ring optionally independently substituted with one or more groups T<sup>1d</sup>, T<sup>2d</sup> or T<sup>3d</sup>.  
 or one of R<sup>6a</sup> or R<sup>7a</sup>, may combine with R<sup>8a1</sup> to form a saturated or unsaturated 5 to 8  
 10 membered ring optionally independently substituted with one or more groups T<sup>1d</sup>, T<sup>2d</sup> or T<sup>3d</sup>  
 R<sup>8b</sup> is H, alkyl, aryl, cyano, nitro, acyl or -SO<sub>2</sub>(alkyl) where the alkyl and aryl groups may be optionally independently substituted with one or more groups T<sup>1d</sup>, T<sup>2d</sup> or T<sup>3d</sup>;  
 15 R<sup>8c</sup> is H, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, arylalkyl, heterocyclo, heteroaryl, alkoxy or aryloxy any of which may be optionally independently substituted with one or more groups T<sup>1d</sup>, T<sup>2d</sup> or T<sup>3d</sup>;  
 R<sup>8d</sup> is R<sup>4</sup>, COR<sup>4</sup>, CO<sub>2</sub>R<sup>4</sup>, SO<sub>2</sub>R<sup>4</sup>, CONR<sup>6</sup>R<sup>7</sup>, or SO<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>;  
 R<sup>9</sup> and R<sup>10</sup> are independently H, alkyl, hydroxy, alkoxy, aryloxy, heterocycloxy,  
 20 heteroaryloxy, (hydroxy)alkyl, (alkoxy)alkyl, (aryloxy)alkyl, (heterocycloxy)alkyl, (heteroaryloxy)alkyl, cycloalkyl, (cycloalkyl)alkyl, aryl, (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, heterocyclo, or (heterocyclo)alkyl any of which may be optionally independently substituted with one or more groups T<sup>1f</sup>, T<sup>2f</sup> or T<sup>3f</sup>  
 25 or R<sup>9</sup> and R<sup>10</sup> together with the nitrogen atom to which they are attached may combine to form a saturated or unsaturated ring which may be optionally independently substituted with one or more groups T<sup>1f</sup>, T<sup>2f</sup> or T<sup>3f</sup>;  
 W is =NR<sup>8a1</sup>, =N-CO<sub>2</sub>R<sup>8a1</sup>, =N-COR<sup>8a1</sup>, =N-CN, =N-SO<sub>2</sub>R<sup>8a1</sup>, or  

$$\begin{array}{c} \text{=C-NO}_2 \\ | \\ \text{H} \end{array};$$
  
 30 X<sup>1</sup> is O, S, NR<sup>8a2</sup> or CH<sub>2</sub>;  
 Z, Z<sup>1</sup> and Z<sup>2</sup> are independently =O, =S, =NR<sup>8a3</sup> or =N-CN;

$R^X$  is one or more optional substituents, attached to any available ring carbon atom,  
independently selected from  $T^{1g}$ ,  $T^{2g}$  or  $T^{3g}$ ;

$T^{1-1g}$ ,  $T^{2-2g}$ , and  $T^{3-3g}$  are each independently

- (1) hydrogen or  $T^6$ , where  $T^6$  is
    - (i) alkyl, (hydroxy)alkyl, (alkoxy)alkyl, alkenyl, alkynyl, cycloalkyl, (cycloalkyl)alkyl, cycloalkenyl, (cycloalkenyl)alkyl, aryl, (aryl)alkyl, heterocyclo, (heterocylco)alkyl, heteroaryl, or (heteroaryl)alkyl;
    - (ii) a group (i) which is itself substituted by one or more of the same or different groups (i); or
    - (iii) a group (i) or (ii) which is independently substituted by one or more (preferably 1 to 3) of the following groups (2) to (13) of the definition of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ ,
  - (2)  $-OH$  or  $-OT^6$ ,
  - (3)  $-SH$  or  $-ST^6$ ,
  - (4)  $-C(O)_tH$ ,  $-C(O)_tT^6$ , or  $-O-C(O)T^6$ , where  $t$  is 1 or 2;
  - (5)  $-SO_3H$ ,  $-S(O)_tT^6$ , or  $S(O)_tN(T^9)T^6$ ,
  - (6) halo,
  - (7) cyano,
  - (8) nitro,
  - (9)  $-T^4-NT^7T^8$ ,
  - (10)  $-T^4-N(T^9)-T^5-NT^7T^8$ ,
  - (11)  $-T^4-N(T^{10})-T^5-T^6$ ,
  - (12)  $-T^4-N(T^{10})-T^5-H$ ,
  - (13) oxo,
- $T^4$  and  $T^5$  are each independently
- (1) a single bond,
  - (2)  $-T^{11}-S(O)_t-T^{12}-$ ,
  - (3)  $-T^{11}-C(O)-T^{12}-$ ,
  - (4)  $-T^{11}-C(S)-T^{12}-$ ,
  - (5)  $-T^{11}-O-T^{12}-$ ,
  - (6)  $-T^{11}-S-T^{12}-$ ,

- (7)  $-T^{11}-O-C(O)-T^{12}-$ ,
- (8)  $-T^{11}-C(O)-O-T^{12}-$ ,
- (9)  $-T^{11}-C(=NT^{9a})-T^{12}-$ , or
- (10)  $-T^{11}-C(O)-C(O)-T^{12}-$

5  $T^7, T^8, T^9, T^{9a}$  and  $T^{10}$

- (1) are each independently hydrogen or a group provided in the definition of  $T^6$ , or
- (2)  $T^7$  and  $T^8$  may together be alkylene or alkenylene, completing a 3- to 8-membered saturated or unsaturated ring together with the atoms to which they are attached, which ring is unsubstituted or substituted with one or more groups listed in the description of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ , or
- (3)  $T^7$  or  $T^8$ , together with  $T^9$ , may be alkylene or alkenylene completing a 3- to 8-membered saturated or unsaturated ring together with the nitrogen atoms to which they are attached, which ring is unsubstituted or substituted with one or more groups listed in the description of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ , or
- (4)  $T^7$  and  $T^8$  or  $T^9$  and  $T^{10}$  together with the nitrogen atom to which they are attached may combine to form a group  $-N=CT^{13}T^{14}$  where  $T^{13}$  and  $T^{14}$  are each independently H or a group provided in the definition of  $T^6$ ; and

20  $T^{11}$  and  $T^{12}$  are each independently

- (1) a single bond,
- (2) alkylene,
- (3) alkenylene, or
- (4) alkynylene.


25

The present invention provides novel methods for the prevention and treatment of arrhythmia and  $I_{Kur}$ -associated disorders employing one or more compounds of the formula I, enantiomers, diastereomers or pharmaceutically acceptable salts thereof. In particular the present invention provides a novel method for the selective prevention and treatment of supraventricular arrhythmias.

30


Preferred compounds within the scope of formula I include compounds and salts thereof wherein one or more, and especially all of Q, R<sup>2</sup>, J and R<sup>3</sup> are selected from the following definitions:

Q is NR<sup>1</sup> or O;




R<sup>2</sup> is aryl, (aryl)alkyl or heteroaryl any of which may be optionally independently substituted with one or more T<sup>1</sup>, T<sup>2</sup> T<sup>3</sup>;

J is a bond or methylene; and




More preferred compounds within the scope of formula I include compounds and salts thereof wherein one or more, and especially all of Q, R<sup>2</sup>, J and R<sup>3</sup> are selected from the following definitions:

15 Q is NR<sup>1</sup>;



20 R<sup>2</sup> is aryl, (aryl)alkyl or heteroaryl (especially where aryl is phenyl and heteroaryl is thiophenyl) any of which may be optionally independently substituted with one or more T<sup>1</sup>, T<sup>2</sup> T<sup>3</sup>;

J is a bond or methylene;



$R^4$  is alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, (aryl)alkyl, heteroaryl or (heteroaryl)alkyl any of which may be optionally independently substituted with one or more  $T^{1b}$ ,  $T^{2b}$   $T^{3b}$ ;





$R^5$  is



5 (a)  $-NR^{6a}R^{7a}$  or

(b) aryl, (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, heterocyclo or (heterocyclo)alkyl any of which may be optionally independently substituted with one or more  $T^{1c}$ ,  $T^{2c}$   $T^{3c}$ ;

10  $R^6$ ,  $R^{6a}$ ,  $R^7$  and  $R^{7a}$  are independently H, alkyl, alkenyl, alkynyl, aryl, (aryl)alkyl, (alkoxy)alkyl, cycloalkyl, (cycloalkyl)alkyl, (hydroxy)alkyl, heteroaryl, (heteroaryl)alkyl, heterocyclo, (heterocyclo)alkyl, (aryloxy)alkyl,  $-C(O)R^9$ ,  $-CO_2R^9$ , or  $-C(O)-NR^9R^{10}$  any of which may be optionally independently substituted with one or more  $T^{1d}$ ,  $T^{2d}$   $T^{3d}$ ;

15 or  $R^6$  and  $R^7$ , or  $R^{6a}$  and  $R^{7a}$  together with the nitrogen atom to which they are attached combine to form an optionally substituted 4 to 8 membered

heterocyclo ring (e.g., , , , ,



20

$R^{8a}$  is H, alkyl, or (aryl)alkyl;

$R^{8c}$  is

(a) alkyl, aryl, heteroaryl any of which may be optionally independently substituted with one or more  $T^{1d}$ ,  $T^{2d}$   $T^{3d}$ ; or


25 (b)  $-NR^9R^{10}$ ;

W is =N-CN;

Z<sup>1</sup> is =O or =N-CN; and


T<sup>1</sup>, T<sup>1b</sup>, T<sup>1c</sup>, T<sup>1d</sup>, T<sup>2</sup>, T<sup>2b</sup>, T<sup>2c</sup>, T<sup>2d</sup>, T<sup>3</sup>, T<sup>3b</sup>, T<sup>3c</sup> and T<sup>3d</sup> are independently halo, cyano, alkyl, aryl, (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, haloalkyl, -OH, -OT<sup>6</sup>,  
 5 -C(O)<sub>t</sub>T<sup>6</sup>, -SO<sub>2</sub>T<sup>6</sup>, -T<sup>4</sup>NT<sup>7</sup>T<sup>8</sup>, or -T<sup>4</sup>N(T<sup>10</sup>)T<sup>5</sup>-T<sup>6</sup>.

Preferred compounds generally have the structure



10

Preferred -JR<sup>3</sup> moieties include:



15

### DETAILED DESCRIPTION OF THE INVENTION

The following are definitions of terms used in this specification. The initial definition provided for a group or term herein applies to that group or term throughout

the present specification, individually or as part of another group, unless otherwise indicated.

The terms "alk" or "alkyl" refer to straight or branched chain hydrocarbon groups having 1 to 12 carbon atoms, preferably 1 to 8 carbon atoms, such as methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, t-butyl, pentyl, hexyl, heptyl, octyl, etc. Lower alkyl groups, that is, alkyl groups of 1 to 6 carbon atoms, are generally most preferred. The term "substituted alkyl" refers to alkyl groups substituted with one or more groups listed in the definition of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ , preferably selected from cyano, halo, oxo, hydroxy,  $-OT^6$ ,  $-C(O)_tT^6$ ,  $-OC(O)T^6$ ,  $-T^4-NT^7T^8$ ,  $-T^4-N(T^9)-T^5-T^6$ ,  $-S(O)_tT^6$  or  $-S(O)_tN(T^9)T^6$ .

The term "alkenyl" refers to straight or branched chain hydrocarbon groups having 2 to 12 carbon atoms, preferably 2 to 4 carbon atoms, and at least one double carbon to carbon bond (either cis or trans), such as ethenyl. The term "substituted alkenyl" refers to alkenyl groups substituted with one or more groups listed in the definition of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ , preferably selected from cyano, halo, oxo, hydroxy,  $-OT^6$ ,  $-C(O)_tT^6$ ,  $-OC(O)T^6$ ,  $-T^4-NT^7T^8$ ,  $-T^4-N(T^9)-T^5-T^6$ ,  $-S(O)_tT^6$  or  $-S(O)_tN(T^9)T^6$ .

The term "alkynyl" refers to straight or branched chain hydrocarbon groups having 2 to 12 carbon atoms, preferably 2 to 4 carbon atoms, and at least one triple carbon to carbon bond, such as ethynyl. The term "substituted alkynyl" refers to alkynyl groups substituted with one or more groups listed in the definition of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ , preferably selected from cyano, halo, oxo, hydroxy,  $-OT^6$ ,  $-C(O)_tT^6$ ,  $-OC(O)T^6$ ,  $-T^4-NT^7T^8$ ,  $-T^4-N(T^9)-T^5-T^6$ ,  $-S(O)_tT^6$  or  $-S(O)_tN(T^9)T^6$ .


The term "alkylene" refers to a straight chain bridge of 1 to 4 carbon atoms connected by single bonds (e.g.,  $-(CH_2)_x-$  wherein x is 1 to 5), which may be substituted with one or more groups listed in the definition of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ , preferably selected from cyano, halo, oxo, hydroxy,  $-OT^6$ ,  $-C(O)_tT^6$ ,  $-OC(O)T^6$ ,  $-T^4-NT^7T^8$ ,  $-T^4-N(T^9)-T^5-T^6$ ,  $-S(O)_tT^6$  or  $-S(O)_tN(T^9)T^6$ .

The term "alkenylene" refers to a straight chain bridge of 2 to 5 carbon atoms having one or two double bonds that is connected by single bonds and may be substituted with one or more groups listed in the definition of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ , preferably selected from cyano, halo, oxo, hydroxy,  $-OT^6$ ,  $-C(O)_tT^6$ ,  $-OC(O)T^6$ ,

$-T^4-NT^7T^8$ ,  $-T^4-N(T^9)-T^5-T^6$ ,  $-S(O)_tT^6$  or  $-S(O)_tN(T^9)T^6$ . Exemplary alkenylene groups are  $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$ ,  $-\text{CH}_2-\text{CH}=\text{CH}-$ ,  $-\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-$ ,  $-\text{C}(\text{CH}_3)_2\text{CH}=\text{CH}-$  and  $-\text{CH}(\text{C}_2\text{H}_5)-\text{CH}=\text{CH}-$ .

The term "alkynylene" refers to a straight chain bridge of 2 to 5 carbon atoms that has a triple bond therein, is connected by single bonds, and may be substituted with one or more groups listed in the definition of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ , preferably selected from cyano, halo, oxo, hydroxy,  $-\text{OT}^6$ ,  $-\text{C}(\text{O})_tT^6$ ,  $-\text{OC}(\text{O})T^6$ ,  $-T^4-NT^7T^8$ ,  $-T^4-N(T^9)-T^5-T^6$ ,  $-S(O)_tT^6$  or  $-S(O)_tN(T^9)T^6$ . Exemplary alkynylene groups are  $-\text{C}\equiv\text{C}-$ ,  $-\text{CH}_2-\text{C}\equiv\text{C}-$ ,  $-\text{CH}(\text{CH}_3)-\text{C}\equiv\text{C}-$  and  $-\text{C}\equiv\text{C}-\text{CH}(\text{C}_2\text{H}_5)\text{CH}_2-$ .


The terms "ar" or "aryl" refer to aromatic homocyclic (i.e., hydrocarbon) mono-, bi- or tricyclic ring-containing groups preferably having 6 to 14 members such as phenyl, naphthyl and biphenyl, as well as such rings fused to a cycloalkyl, cycloalkenyl, heterocyclo, or heteroaryl ring. Examples include:





The term "substituted aryl" refers to aryl groups substituted with one or more groups listed in the definition of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ , preferably selected cyano, halo, oxo, hydroxy,  $-\text{OT}^6$ ,  $-\text{C}(\text{O})_tT^6$ ,  $-\text{OC}(\text{O})T^6$ ,  $-T^4-NT^7T^8$ ,  $-T^4-N(T^9)-T^5-T^6$ ,  $-S(O)_tT^6$  or  $-S(O)_tN(T^9)T^6$ .

The term "cycloalkyl" refers to saturated and partially unsaturated (containing 1 or 2 double bonds) cyclic hydrocarbon groups containing 1 to 3 rings, including monocycloalkyl, bicycloalkyl and tricycloalkyl, containing a total of 3 to 20 carbons forming the rings, preferably 3 to 7 carbons forming the ring, and which may be fused to 1 or 2 aromatic or heterocyclo rings, which include cyclopropyl, cyclobutyl,

cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclodecyl, cyclododecyl,  
cyclohexenyl,





5





terms "substituted cycloalkyl" refers to cycloalkyl groups substituted with one or  
10 more groups listed in the definition of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ , preferably selected from  
cyano, halo, oxo, hydroxy,  $-OT^6$ ,  $-C(O)_tT^6$ ,  $-OC(O)T^6$ ,  $-T^4-NT^7T^8$ ,  
 $-T^4-N(T^9)-T^5-T^6$ ,  $-S(O)_tT^6$  or  $-S(O)_tN(T^9)T^6$ .

The terms "halogen" and "halo" refer to fluorine, chlorine, bromine and iodine.




The terms "heterocycle", "heterocyclic", "heterocyclyl", "heterocyclic group"  
15 or "heterocyclo" refer to fully saturated or partially or unsaturated cyclic groups (for  
example, 3 to 13 member monocyclic, 7 to 17 member bicyclic, or 10 to 20 member  
tricyclic ring systems, preferably containing a total of 3 to 10 ring atoms) which have  
at least one heteroatom in at least one carbon atom-containing ring. Each ring of the  
heterocyclic group containing a heteroatom may have 1, 2, 3 or 4 heteroatoms selected  
20 from nitrogen atoms, oxygen atoms and/or sulfur atoms, where the nitrogen and sulfur  
heteroatoms may optionally be oxidized and the nitrogen heteroatoms may optionally  
be substituted or quaternized. The heterocyclic group may be attached at any  
heteroatom or carbon atom of the ring or ring system. The rings of multi-ring  
heterocycles may be either fused, bridged and/or joined through one or more spiro  
25 unions. Exemplary heterocyclic groups include



5



10









and the like.

15




The terms "substituted heterocycle", "substituted heterocyclic", "substituted heterocyclic group" and "substituted heterocyclo" refer to heterocycle, heterocyclic and heterocyclo groups substituted with one or more groups listed in the definition of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ , preferably selected from cyano, halo, oxo, hydroxy,  $-OT^6$ ,  
 5  $-C(O)_tT^6$ ,  $-OC(O)T^6$ ,  $-T^4-NT^7T^8$ ,  $-T^4-N(T^9)-T^5-T^6$ ,  $-S(O)_tT^6$  or  $-S(O)_tN(T^9)T^6$ .

The term "heteroaryl" as used herein alone or as part of another group refers to a 5- 6- or 7- membered aromatic rings containing from 1 to 4 nitrogen atoms and/or 1 or 2 oxygen or sulfur atoms provided that the ring contains at least 1 carbon atom and  
 10 no more than 4 heteroatoms. The heteroaryl ring is linked through an available carbon or nitrogen atom. Also included within the definition of heteroaryl are such rings fused to a cycloalkyl, aryl, cycloheteroalkyl, or another heteroaryl ring. One, two, or three available carbon or nitrogen atoms in the heteroaryl ring can be optionally substituted with substituents listed in the description of  $T^1$ ,  $T^2$  and  $T^3$ . Also  
 15 an available nitrogen or sulfur atom in the heteroaryl ring can be oxidized. Examples of heteroaryl rings include






5



10



Throughout the specification, groups and substituents thereof may be chosen  
10 to provide stable moieties and compounds.

The compounds of the present invention form salts which are also within the scope of this invention. Reference to a compound of the present invention herein is understood to include reference to salts thereof, unless otherwise indicated. The term "salt(s)", as employed herein, denotes acidic and/or basic salts formed with inorganic and/or organic acids and bases. In addition, when a compound of formula I contains both a basic moiety and an acidic moiety, zwitterions ("inner salts") may be formed and are included within the term "salt(s)" as used herein. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful, e.g., in isolation or purification steps which may be employed during preparation. Salts of the compounds of the formula I may be formed, for example, by reacting a compound I with an amount of acid or base, such as an equivalent amount,

in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

The compounds of the present invention which contain a basic moiety may form salts with a variety of organic and inorganic acids. Exemplary acid addition salts include acetates (such as those formed with acetic acid or trihaloacetic acid, for example, trifluoroacetic acid), adipates, alginates, ascorbates, aspartates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, cyclopentanepropionates, digluconates, dodecylsulfates, ethanesulfonates, fumarates, glucoheptanoates, glycerophosphates, hemisulfates, heptanoates, hexanoates, hydrochlorides (formed with hydrochloric acid), hydrobromides (formed with hydrogen bromide), hydroiodides, 2-hydroxyethanesulfonates, lactates, maleates (formed with maleic acid), methanesulfonates (formed with methanesulfonic acid), 2-naphthalenesulfonates, nicotines, nitrates, oxalates, pectinates, persulfates, 3-phenylpropionates, phosphates, picrates, pivalates, propionates, salicylates, succinates, sulfates (such as those formed with sulfuric acid), sulfonates (such as those mentioned herein), tartrates, thiocyanates, toluenesulfonates such as tosylates, undecanoates, and the like.

The compounds of the present invention which contain an acidic moiety may form salts with a variety of organic and inorganic bases. Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases (for example, organic amines) such as benzathines, dicyclohexylamines, hydrabamines (formed with N,N-bis(dehydroabietyl)ethylenediamine), N-methyl-D-glucamines, N-methyl-D-glucamides, t-butyl amines, and salts with amino acids such as arginine, lysine and the like.

Basic nitrogen-containing groups may be quaternized with agents such as lower alkyl halides (e.g. methyl, ethyl, propyl, and butyl chlorides, bromides and iodides), dialkyl sulfates (e.g. dimethyl, diethyl, dibutyl, and diamyl sulfates), long chain halides (e.g. decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides), aralkyl halides (e.g. benzyl and phenethyl bromides), and others.

Prodrugs and solvates of the compounds of the invention are also contemplated herein. The term "prodrug", as employed herein, denotes a compound

which, upon administration to a subject, undergoes chemical conversion by metabolic or chemical processes to yield a compound of the formula I, or a salt and/or solvate thereof. Solvates of the compounds of formula I are preferably hydrates.

To the extent that compounds of the the present invention, and salts thereof,  
5 may exist in their tautomeric form, all such tautomeric forms are contemplated herein as part of the present invention.


All stereoisomers of the present compounds, such as those which may exist due to asymmetric carbons on the various R and Z substituents, including enantiomeric forms (which may exist even in the absence of asymmetric carbons) and  
10 diastereomeric forms, are contemplated within the scope of this invention. Individual stereoisomers of the compounds of the invention may, for example, be substantially free of other isomers, or may be admixed, for example, as racemates or with all other, or other selected, stereoisomers. The chiral centers of the present invention can have the S or R configuration as defined by the IUPAC 1974 Recommendations.

15 The terms "including", "such as", "for example" and the like are intended to refer to exemplary embodiments and not to limit the scope of the present invention.

SCHEMES

Compounds of the formula I may be prepared using the sequence of steps outlined below.


5

SCHEME 1

Compounds of the formula I may be prepared using the sequence of steps outlined in scheme 1. Specifically, compounds of the formula I where R<sup>2</sup> is heteroaryl or substituted heteroaryl and R<sup>5</sup> is aryl, substituted aryl, heteroaryl or substituted heteroaryl may be prepared using Scheme 1. Heteroaryl acetonitrile **1** is deprotonated and alkylated to form the N-protected piperidine ring **2**. Reduction of nitrile **2** generates primary amine **3**. The amine is subsequently acylated with Heterocycle2 acyl chloride, deprotected and the resulting amine is taken onto final product carbamates, sulfonamides, sulfenyl ureas and cyanoguanidines.

Compounds of formula I where R<sup>3</sup> is  $\text{---}\overset{\text{Z}^1}{\parallel}\text{---R}^5$ , Z<sup>1</sup> is O and R<sup>5</sup> is -NR<sup>6a</sup>R<sup>7a</sup> may be prepared as described in Scheme 2.

## SCHEME 2



- 5 Protection of the nitrogen atom of compound 1 gives compound 2. One skilled in the art will recognize a variety of nitrogen protecting groups that are known in the literature. In this example, suitable nitrogen protecting groups include the benzyl (Bn), *N*-*tert*-butoxycarbonyl (Boc) and carbobenzyloxy (CBz) groups. The carboxylic acid moiety of compound 2 may be coupled with an amine  $\text{HNR}^{6a}\text{R}^{7a}$
- 10 using a variety of coupling procedures known in the literature to provide carboxamide compound 3. The nitrogen atom of compound 3 may be deprotected and made to react with a carboxylic acid (e.g.;  $\text{R}^4\text{CO}_2\text{H}$ ; shown in Scheme 1) in the presence of a coupling agent or an acid chloride (e.g.;  $\text{R}^4\text{COCl}$ ) in the presence of an acid scavenger such as triethylamine or polystyrene-diisopropylethylamine resin to give compound 5
- 15 where Q is  $\text{NR}^1$  and  $\text{R}^1$  is  $\text{—}\overset{\text{O}}{\parallel}\text{—}\text{R}^4$ . In addition to carboxylic acids or acid chlorides, one skilled in the art will recognize that the piperidine nitrogen atom of compound 4 may be made to react with a number of other readily available raw materials to provide compounds of formula I. For example, compound 4 may be made to react with sulfonyl chlorides (e.g.;  $\text{R}^{8c}\text{SO}_2\text{Cl}$ ) in the presence of an acid scavenger to
- 20 provide compounds of formula I where Q is  $\text{NR}^1$  and  $\text{R}^1$  is  $\text{—}\overset{\text{O}}{\parallel}\text{S}\text{—}\text{R}^{8c}$ . Compound 4 may be made to react with isocyanates (e.g.;  $\text{R}^7\text{R}^6\text{NCO}$ ) to provide compounds of formula I where Q is  $\text{NR}^1$  and  $\text{R}^1$  is  $\text{—}\overset{\text{O}}{\parallel}\text{NR}^6\text{R}^7$ . Compound 4 may be made to react


with chloroformates (e.g.;  $R^4\text{OCOCI}$ ) in the presence of an acid scavenger to provide compounds of formula I where Q is  $\text{NR}^1$  and  $R^1$  is  $\text{—}\overset{\text{O}}{\parallel}\text{—O—R}^4$ . Compound 4 may be made to react with certain heteroaryl groups (e.g; heteroaryl-X where X is a leaving group such as a halogen atom) in the presence of an acid scavenger to provide

5 compound 5 where Q is  $\text{NR}^1$  and  $R^1$  is heteroaryl. This reaction is may be performed in an organic solvent such a tetrahydrofuran or acetonitrile at elevated temperatures. Alternatively, this reaction may be performed in the presence of a palladium catalyst to provide compound 5 where Q is  $\text{NR}^1$  and  $R^1$  is heteroaryl.

Additional compounds of formula I where  $R^3$  is  $\text{—}\overset{\text{Z}^1}{\parallel}\text{—R}^5$ ,  $\text{Z}^1$  is O and  $R^5$  is  $\text{—NR}^{6a}\text{R}^{7a}$  may be prepared using as described in Scheme 3 and Scheme 4.

10

### SCHEME 3




15

Referring to Scheme 2, compound 1 may be made to react with diphenyl *N*-cyanocarbonimidate in a solvent such as tetrahydrofuran, acetonitrile or isopropanol to provide compound 2. Typically this reaction is conducted at elevated temperatures. Compound 2 may be made to react with an amine  $\text{HNR}^6\text{R}^7$  to provide cyano

20 guanidine compound 3 where Q is  $\text{NR}^1$  and  $R^1$  is  $\text{—}\overset{\text{W}}{\parallel}\text{—NR}^6\text{R}^7$  and W is N-CN.


## SCHEME 4



- 5 Referring to Scheme 3, compound 1 may be made to react with compound 2 to provide sulfenyl urea compound 3 where Q is  $\text{NR}^1$  and  $\text{R}^1$  is  $\text{—}\overset{\text{O}}{\overset{\text{O}}{\text{S}}}\text{—NR}^6\text{R}^7$ . Compound 2 may be prepared by reacting 2-chloroethanol with chlorosulfonyl isocyanate followed by an amine  $\text{HNR}^6\text{R}^7$  in the presence of an acid scavenger such as triethylamine in an organic solvent such as dichloromethane. One skilled in the art
- 10 will recognize certain sulfenyl ureas may be prepared from commercially available raw materials. For example, compound 1 may be made to react with dimethylsulfamoyl chloride in a solvent such as tetrahydrofuran or dichloromethane in the presence of an acid scavenger such as triethylamine or polystyrene-diisopropylethylamine resin to provide compound 3 where Q is  $\text{NR}^1$  and  $\text{R}^1$  is
- 15  $\text{—}\overset{\text{O}}{\overset{\text{O}}{\text{S}}}\text{—NR}^6\text{R}^7$  and  $\text{R}^6$  and  $\text{R}^7$  are each methyl. Compound 1 may be made to react with sulfamide in a solvent such as 1,4-dioxane at elevated temperature to provide compound 3 where Q is  $\text{NR}^1$  and  $\text{R}^1$  is  $\text{—}\overset{\text{O}}{\overset{\text{O}}{\text{S}}}\text{—NR}^6\text{R}^7$  and  $\text{R}^6$  and  $\text{R}^7$  are each hydrogen.


Compounds of formula I where J is  $\text{C}_1$ -alkylene (e.g.;  $\text{CH}_2$ ),  $\text{R}^3$  is  $\text{—}\overset{\text{Z}^1}{\text{C}}\text{—R}^5$ ,  $\text{Z}^1$  is O and  $\text{R}^5$  is  $\text{—NR}^6\text{R}^7$  may be prepared using as described in Scheme 5.

## SCHEME 5



- 5 The carboxylic acid compound 1 may be treated with thionyl chloride to convert the carboxylic acid to the acid chloride that is made to react with diazomethane in a solvent such as ethyl ether to produce a diazo intermediate compound 2. Compound 2 may be irradiated under ultra-violet light ( $\lambda = 365 \text{ nm}$ ) in a solvent such as methanol to provide an ester that may be hydrolyzed to the
- 10 carboxylic acid compound 3 by treatment with aqueous lithium hydroxide. The carboxylic acid moiety of compound 3 may be coupled with an amine  $\text{HNR}^{6a}\text{R}^{7a}$  using a variety of coupling procedures known in the literature to provide carboxamide compound 4. The nitrogen atom of compound 3 may be deprotected and the nitrogen atom further derivatized (see Schemes 1-3) to provide compounds of formula I.


- 15 Compounds of formula I where J is a bond,  $\text{R}^3$  is  $\text{—N—}\overset{\text{Z}^1}{\underset{\text{R}^{8a}}{\parallel}}\text{—R}^5$ ,  $\text{Z}^1$  is O,  $\text{R}^5$  is  $\text{—NR}^{6a}\text{R}^{7a}$  and  $\text{R}^{8a}$  is hydrogen may be prepared using as described in Scheme 6.

SCHEME 6

- 5 Compound 1 may be made to react with diphenylphosphoryl azide in a solvent such as chlorobenzene at an elevated temperature to produce isocyanate compound 2. Compound 2 may be made to react with an amine HNR<sup>6a</sup>R<sup>7a</sup> to produce the urea

compound 3 where J is a bond, R<sup>3</sup> is  $\text{---}\text{N}^{\text{Z}^1}\text{---}\text{R}^5$ , Z<sup>1</sup> is O, R<sup>5</sup> is  $\text{---}\text{NR}^{6a}\text{R}^{7a}$  and R<sup>8a</sup> is hydrogen.

- 10 Compounds of formula I where J is a bond, R<sup>3</sup> is  $\text{---}\text{N}^{\text{Z}^1}\text{---}\text{OR}^5$ , Z<sup>1</sup> is O and R<sup>8a</sup> is hydrogen may be prepared using as described in Scheme 7.

SCHEME 7

15


Isocyanate compound 1 may be made to react with an alcohol R<sup>5</sup>OH in a solvent such as chlorobenzene or tetrahydrofuran to provide compound 2 where J is a

bond, R<sup>3</sup> is  $\text{---}\text{N}^{\text{Z}^1}\text{---}\text{OR}^5$ , Z<sup>1</sup> is O and R<sup>8a</sup> is hydrogen.

Compounds of formula I where J is a bond,  $R^3$  is  $-\text{N}(\text{R}^{8a})\text{C}(\text{Z}^1)\text{R}^5$ ,  $Z^1$  is O,  $R^5$  is not  $\text{NR}^{6a}\text{R}^{7a}$  and  $R^{8a}$  is hydrogen may be prepared using as described in Scheme 8.

# SCHEME 8


5



Compound 1 (see Scheme 6;  $R^5$  is *tert*-butyl) may be deprotected by treatment with trifluoroacetic acid in a solvent such as dichloromethane to provide amine compound 2. The amino group of compound 2 may be made to react with a carboxylic acid (e.g.;  $R^5\text{CO}_2\text{H}$ ; shown in Scheme 7) in the presence of a coupling agent or an acid chloride (e.g.;  $R^5\text{COCl}$ ) in the presence of an acid scavenger such as triethylamine or polystyrene-diisopropylethylamine resin to give compound 3 where J is a bond,  $R^3$  is  $-\text{N}(\text{R}^{8a})\text{C}(\text{Z}^1)\text{R}^5$ ,  $Z^1$  is O,  $R^5$  is not  $\text{NR}^{6a}\text{R}^{7a}$  and  $R^{8a}$  is hydrogen. In

addition to carboxylic acids or acid chlorides, one skilled in the art will recognize that compound 2 may be made to react with a number of other readily available raw materials to provide compounds of formula I. For example, compound 2 may be made to react with sulfonyl chlorides (e.g.;  $R^5\text{SO}_2\text{Cl}$ ) in the presence of an acid scavenger to provide compounds of formula I where J is a bond and  $R^3$  is  $-\text{N}(\text{R}^{8a})\text{S}(\text{O})_2\text{R}^5$ .

Compounds of formula I where  $R^3$  is  $R^5$ ,  $R^5$  is  $\text{NR}^{6a}\text{R}^{7a}$  and  $R^{6a}$  is heteroaryl may be prepared as described in Scheme 9.


SCHEME 9

- 5 Amine compound 1 may be made to react with a substituted aryl or heteroaryl compound where X is a halogen atom, triflate or similar leaving group to provide compound 2. This reaction may be conducted in an organic solvent such as tetrahydrofuran or acetonitrile at an elevated temperature. Alternatively, this reaction may be performed in the presence of a palladium catalyst to provide compound 2
- 10 where R<sup>3</sup> is R<sup>5</sup>, R<sup>5</sup> is NR<sup>6a</sup>R<sup>7a</sup> and R<sup>6a</sup> is heteroaryl.

Compounds of formula I where R<sup>3</sup> is R<sup>5</sup> and R<sup>5</sup> is heteroaryl may be prepared as described in Scheme 10 and Scheme 11.

SCHEME 10


15



- Referring to Scheme 9, acid chloride compound 1 may be made to react compound 2 in an organic solvent such as dichloromethane to provide carboxamide compound 3. Cyclization of compound 2 in the presence of an acid such as acetic acid at elevated temperature provides compound 3 where R<sup>3</sup> is R<sup>5</sup> and R<sup>5</sup> is heteroaryl (e.g.; benzimidazole). One skilled in the art will recognize that carboxylic acids or acid chlorides may be converted to a wide variety of heteroaryl groups. For example, compound 1 may be made to react with a 2-amino phenol to provide compounds of
- 20

formula I where  $R^3$  is  $R^5$  and  $R^5$  is heteroaryl (e.g.; benzoxazole). Compound 1 may be made to react with a 2-amino benzenethiol to provide compounds of formula I where  $R^3$  is  $R^5$  and  $R^5$  is heteroaryl (e.g.; benzthiazole). Compound 1 may be made to react with an *N*-hydroxyamidine to provide compounds of formula 3 where  $R^3$  is  $R^5$  and  $R^5$  is heteroaryl (e.g.; oxadiazole).

### SCHEME 11




10

Referring to Scheme 10, cyano compound 1 may be made to react with sodium azide in a solvent such as methyl sulfoxide at elevated temperatures to provide tetrazole compound 2. Compound 2 may be made to react with  $T^{1c}-X$  where  $X$  is a leaving group such as a halogen atom or triflate to provide compound 3 where  $R^3$  is  $R^5$  and  $R^5$  is heteroaryl (e.g.; tetrazole).


15

Compounds of formula I where  $R^2$  is alkyl, cycloalkyl or (aryl)alkyl may be prepared as described in Scheme 12.


## SCHEME 12



- 5            Protection of the nitrogen atom of compound 1 provides compound 2.
- Deprotonation of compound 2 with a base such as lithium diisopropylamide in an organic solvent such as tetrahydrofuran at a low temperature followed by reaction with a benzyl halide, for example, provides compound 3 where  $R^2$  is (aryl)alkyl. One skilled in the art will recognize that aldehydes and ketones may also be made to react with
- 10   compound 2 after deprotonation with lithium diisopropylamide. The cyano group of compound 3 may be reduced with lithium aluminum hydride in a solvent such as tetrahydrofuran to provide amino compound 4. Alternatively, the cyano group of compound 3 may be hydrolyzed with aqueous sodium hydroxide to provide carboxylic acid compound 5. Compounds of formula I may be prepared from
- 15   compounds 4 and 5 as described previously.

SCHEME 13

- 5 Compounds of the formula 1 where  $p = 3$ ,  $m = 2$ ,  $Q = NR_1$ ,  $R_2 = \text{aryl}$  may be prepared as described in Scheme 13. The lactam was prepared from intermediate cyclohexanone which was synthesized as described in Journal of Medicinal Chemistry, **1998**, 821. The lactam nitrogen was protected using standard protecting group methodology and the lactam carbonyl group and the nitrile group were reduced either simultaneously or subsequently. The primary amine was acylated, the protecting group was removed and the azepane was further functionalized to the sulfinyl urea, the carbamate, the amide or alkylated directly.

SCHEME 14

Compounds of formula I where Q is O, R<sup>3</sup> is  $\text{---N} \begin{array}{c} \text{Z}^1 \\ \text{||} \\ \text{R}^{8a} \end{array} \text{---R}^5$ , Z<sup>1</sup> is O, and R<sup>8a</sup> is

hydrogen may be prepared as described in Scheme 14. Compound 1 may be made to react with bis(2-chloroethyl)ether under phase-transfer catalysis conditions to provide the cyclized product compound 2. The nitrile group of compound 2 may be reduced using various methods, including treatment with lithium aluminum hydride or hydrogenation in the presence of platinum (IV) oxide, to provide the amine compound 3. The amino group of compound 3 may be made to react with a carboxylic acid (e.g.; R<sup>5</sup>CO<sub>2</sub>H) in the presence of a coupling agent in an organic solvent such as tetrahydrofuran to provide the acylated product compound 4.

Additional compounds within the scope of the present invention can be prepared from the compounds obtained by the above described methods through conversion of the substituent groups to other functionality by the usual methods of chemical synthesis, as illustrated in the following examples.

Compounds of formula I that contain chiral centers may be obtained in non-racemic form by non-racemic synthesis or resolution by methods well known to those skilled in the art. Compounds that are non-racemic are designated as "chiral" in the examples.

In the examples described below it may be necessary to protect reactive functionality such as hydroxy, amino, thio or carboxy groups, where these are desired in the final product, to avoid their unwanted participation in reactions. The introduction and removal of protecting groups are well known to those skilled in the art, for example see (Green, T. W. in "Protective Groups in Organic Synthesis", John Wiley and Sons, 1991).

### Utility

Compounds within the scope of the present invention inhibit the K<sub>v</sub>1 subfamily of voltage-gated K<sup>+</sup> channels, and as such are useful in the treatment and/or prevention of various disorders: cardiac arrhythmias, including supraventricular arrhythmias, atrial arrhythmias, atrial flutter, atrial fibrillation, complications of cardiac ischemia, and use as heart rate control agents; angina pectoris including relief

of Prinzmetal's symptoms, vasospastic symptoms and variant symptoms;  
 gastrointestinal disorders including reflux esophagitis, functional dyspepsia, motility  
 disorders (including constipation and diarrhea), and irritable bowel syndrome;  
 disorders of vascular and visceral smooth muscle including asthma, chronic  
 5 obstructive pulmonary disease, adult respiratory distress syndrome, peripheral  
 vascular disease (including intermittent claudication), venous insufficiency,  
 impotence, cerebral and coronary spasm and Raynaud's disease; inflammatory and  
 immunological disease including inflammatory bowel disease, rheumatoid arthritis,  
 graft rejection, asthma, chronic obstructive pulmonary disease, cystic fibrosis and  
 10 atherosclerosis; cell proliferative disorders including restenosis and cancer (including  
 leukemia); disorders of the auditory system; disorders of the visual system including  
 macular degeneration and cataracts; diabetes including diabetic retinopathy, diabetic  
 nephropathy and diabetic neuropathy; muscle disease including myotonia and  
 wasting; peripheral neuropathy; cognitive disorders; migraine; memory loss including  
 15 Alzheimer's and dementia; CNS mediated motor dysfunction including Parkinson's  
 disease, and ataxia; epilepsy; and other ion channel mediated disorders.

As inhibitors of the  $K_v1$  subfamily of voltage-gated  $K^+$  channels compounds of  
 the present invention are useful to treat a variety of disorders including resistance by  
 transplantation of organs or tissue, graft-versus-host diseases brought about by  
 20 medulla ossium transplantation, rheumatoid arthritis, systemic lupus erythematosus,  
 hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type I diabetes uveitis,  
 juvenile-onset or recent-onset diabetes mellitus, posterior uveitis, allergic  
 encephalomyelitis, glomerulonephritis, infectious diseases caused by  
 pathogenic microorganisms, inflammatory and hyperproliferative skin diseases,  
 25 psoriasis, atopic dermatitis, contact dermatitis, eczematous dermatitis, seborrheic  
 dermatitis, lichen planus, pemphigus, bullous pemphigoid, epidermolysis bullosa,  
 urticaria, angioedemas, vasculitides, erythemas, cutaneous eosinophilias, Lupus  
 erythematosus, acne, alopecia areata, keratoconjunctivitis, vernal conjunctivitis,  
 uveitis associated with Behcet's disease, keratitis, herpetic keratitis, conical cornea,  
 30 dystrophia epithelialis corneae, corneal leukoma, ocular pemphigus, Mooren's ulcer  
 scleritis, Graves' ophthalmopathy, Vogt-Koyanagi-Harada syndrome, sarcoidosis,  
 pollen allergies, reversible obstructive airway disease, bronchial asthma, allergic

asthma, intrinsic asthma, extrinsic asthma, dust asthma, chronic or inveterate asthma,  
 late asthma and airway hyper-responsiveness, bronchitis, gastric ulcers, vascular  
 damage caused by ischemic diseases and thrombosis, ischemic bowel diseases,  
 inflammatory bowel diseases, necrotizing enterocolitis, intestinal lesions associated  
 5 with thermal burns and leukotriene B<sub>4</sub>-mediated diseases, Coeliac diseases, proctitis,  
 eosinophilic gastroenteritis, mastocytosis, Crohn's disease, ulcerative colitis,  
 migraine, rhinitis, eczema, interstitial nephritis, Good-pasture's syndrome, hemolytic-  
 uremic syndrome, diabetic nephropathy, multiple myositis, Guillain-Barre syndrome,  
 Meniere's disease, polyneuritis, multiple neuritis, mononeuritis, radiculopathy,  
 10 hyperthyroidism, Basedow's disease, pure red cell aplasia, aplastic anemia, hypoplastic  
 anemia, idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia,  
 agranulocytosis, pernicious anemia, megaloblastic anemia, anerythroplasia,  
 osteoporosis, sarcoidosis, fibroid lung, idiopathic interstitial pneumonia,  
 dermatomyositis, leukoderma vulgaris, ichthyosis vulgaris, photoallergic sensitivity,  
 15 cutaneous T cell lymphoma, arteriosclerosis, atherosclerosis, aortitis syndrome,  
 polyarteritis nodosa, myocarditis, scleroderma, Wegener's granuloma, Sjogren's  
 syndrome, adiposis, eosinophilic fascitis, lesions of gingiva, periodontium, alveolar  
 bone, substantia osses dentis, glomerulonephritis, male pattern alopecia or alopecia  
 senilis by preventing epilation or providing hair germination and/or promoting hair  
 20 generation and hair growth, muscular dystrophy; Pyoderma and Sezary's syndrome,  
 Addison's disease, ischemia-reperfusion injury of organs which occurs upon  
 preservation, transplantation or ischemic disease, endotoxin-shock,  
 pseudomembranous colitis, colitis caused by drug or radiation, ischemic acute renal  
 insufficiency, chronic renal insufficiency, toxinoses caused by lung-oxygen or drugs,  
 25 lung cancer, pulmonary emphysema, cataracts, siderosis, retinitis, pigmentosa, senile  
 macular degeneration, vitreal scarring, corneal alkali burn, dermatitis erythema  
 multiforme, linear IgA bullous dermatitis and cement dermatitis, gingivitis,  
 periodontitis, sepsis, pancreatitis, diseases caused by environmental pollution, aging,  
 carcinogenesis, metastasis of carcinoma and hypobaropathy, disease caused by  
 30 histamine or leukotriene-C<sub>4</sub> release, Behcet's disease, autoimmune hepatitis, primary  
 biliary cirrhosis sclerosing cholangitis, partial liver resection, acute liver necrosis,  
 necrosis caused by toxin, viral hepatitis, shock, or anoxia, B-virus hepatitis, non-

A/non-B hepatitis, cirrhosis, alcoholic cirrhosis, hepatic failure, fulminant hepatic failure, late-onset hepatic failure, "acute-on-chronic" liver failure, augmentation of chemotherapeutic effect, cytomegalovirus infection, HCMV infection, AIDS, cancer, senile dementia, trauma, and chronic bacterial infection.

5           The compounds of the present invention are antiarrhythmic agents which are useful in the prevention and treatment (including partial alleviation or cure) of arrhythmias. As inhibitors of  $K_v1.5$  compounds within the scope of the present invention are particularly useful in the selective prevention and treatment of supraventricular arrhythmias such as atrial fibrillation, and atrial flutter. By "selective  
10 prevention and treatment of supraventricular arrhythmias" is meant the prevention or treatment of supraventricular arrhythmias wherein the ratio of the prolongation of the atrial effective refractory period to the prolongation of the ventricular effective refractory period is greater than 1:1. This ratio is preferably greater than 4:1, more preferably greater than 10:1, and most preferably such that prolongation of the atrial  
15 effective refractory response period is achieved without significantly detectable prolongation of the ventricular effective refractory period.

          In addition, the compounds within the scope of the present invention block  $I_{Kur}$ , and thus may be useful in the prevention and treatment of all  $I_{Kur}$ -associated conditions. An " $I_{Kur}$ -associated condition" is a disorder which may be prevented,  
20 partially alleviated or cured by the administration of an  $I_{Kur}$  blocker. The  $K_v1.5$  gene is known to be expressed in stomach tissue, intestinal/colon tissue, the pulmonary artery, and pancreatic beta cells. Thus, administration of an  $I_{Kur}$  blocker could provide useful treatment for disorders such as: reflux esophagitis, functional dyspepsia, constipation, asthma, and diabetes. Additionally,  $K_v1.5$  is known to be expressed in  
25 the anterior pituitary. Thus, administration of an  $I_{Kur}$  blocker could stimulate growth hormone secretion.  $I_{Kur}$  inhibitors can additionally be useful in cell proliferative disorders such as leukemia, and autoimmune diseases such as rheumatoid arthritis and transplant rejection.

          The present invention thus provides methods for the prevention or treatment of  
30 one or more of the aforementioned disorders, comprising the step of administering to a subject in need thereof an effective amount of at least one compound of the present invention. Other therapeutic agents such as those described below may be employed

with the inventive compounds in the present methods. In the methods of the present invention, such other therapeutic agent(s) may be administered prior to, simultaneously with or following the administration of the compound(s) of the present invention.

5           The present invention also provides pharmaceutical compositions comprising at least one of the compounds of the present invention or salts thereof capable of preventing or treating one or more of the aforementioned disorders in an amount effective therefor, and a pharmaceutically acceptable vehicle or diluent. The compositions of the present invention may contain other therapeutic agents as  
10 described below, and may be formulated, for example, by employing conventional solid or liquid vehicles or diluents, as well as pharmaceutical additives of a type appropriate to the mode of desired administration (for example, excipients, binders, preservatives, stabilizers, flavors, etc.) according to techniques such as those well known in the art of pharmaceutical formulation.

15           The compounds of the present invention may be administered by any suitable means, for example, orally, such as in the form of tablets, capsules, granules or powders; sublingually; buccally; parenterally, such as by subcutaneous, intravenous, intramuscular, or intrasternal injection or infusion techniques (e.g., as sterile  
20 injectable aqueous or non-aqueous solutions or suspensions); nasally such as by inhalation spray; topically, such as in the form of a cream or ointment; or rectally such as in the form of suppositories; in dosage unit formulations containing non-toxic, pharmaceutically acceptable vehicles or diluents. The present compounds may, for example, be administered in a form suitable for immediate release or extended release. Immediate release or extended release may be achieved by the use of suitable  
25 pharmaceutical compositions comprising the present compounds, or, particularly in the case of extended release, by the use of devices such as subcutaneous implants or osmotic pumps. In the case where the compounds of formula I are being administered to prevent or treat arrhythmias, the compounds may be administered to achieve chemical conversion to normal sinus rhythm, or may optionally be used in  
30 conjunction with electrical cardioconversion.

Exemplary compositions for oral administration include suspensions which may contain, for example, microcrystalline cellulose for imparting bulk, alginic acid

or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners or flavoring agents such as those known in the art; and immediate release tablets which may contain, for example, microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and/or lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants such as those known in the art. The compounds of formula I may also be delivered through the oral cavity by sublingual and/or buccal administration. Molded tablets, compressed tablets or freeze-dried tablets are exemplary forms which may be used. Exemplary compositions include those formulating the present compound(s) with fast dissolving diluents such as mannitol, lactose, sucrose and/or cyclodextrins. Also included in such formulations may be high molecular weight excipients such as celluloses (avicel) or polyethylene glycols (PEG). Such formulations may also include an excipient to aid mucosal adhesion such as hydroxy propyl cellulose (HPC), hydroxy propyl methyl cellulose (HPMC), sodium carboxy methyl cellulose (SCMC), maleic anhydride copolymer (e.g., Gantrez), and agents to control release such as polyacrylic copolymer (e.g., Carbopol 934). Lubricants, glidants, flavors, coloring agents and stabilizers may also be added for ease of fabrication and use.

Exemplary compositions for nasal aerosol or inhalation administration include solutions in saline which may contain, for example, benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, and/or other solubilizing or dispersing agents such as those known in the art.

Exemplary compositions for parenteral administration include injectable solutions or suspensions which may contain, for example, suitable non-toxic, parenterally acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution, an isotonic sodium chloride solution, or other suitable dispersing or wetting and suspending agents, including synthetic mono- or diglycerides, and fatty acids, including oleic acid.

Exemplary compositions for rectal administration include suppositories which may contain, for example, a suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters or polyethylene glycols, which are solid at ordinary temperatures, but liquify and/or dissolve in the rectal cavity to release the drug.

Exemplary compositions for topical administration include a topical carrier such as Plastibase (mineral oil gelled with polyethylene).

The effective amount of a compound of the present invention may be determined by one of ordinary skill in the art, and includes exemplary dosage amounts for an adult human of from about 0.001 to 100 mg/kg of body weight of active compound per day, which may be administered in a single dose or in the form of individual divided doses, such as from 1 to 4 times per day. It will be understood that the specific dose level and frequency of dosage for any particular subject may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the species, age, body weight, general health, sex and diet of the subject, the mode and time of administration, rate of excretion, drug combination, and severity of the particular condition. Preferred subjects for treatment include animals, most preferably mammalian species such as humans, and domestic animals such as dogs, cats and the like, subject to the aforementioned disorders.

The compounds of the present invention may be employed alone or in combination with each other and/or other suitable therapeutic agents useful in the treatment of the aforementioned disorders or other disorders, including: other antiarrhythmic agents such as Class I agents (e.g., propafenone), Class II agents (e.g., carvediol and propranolol), Class III agents (e.g., sotalol, dofetilide, amiodarone, azimilide and ibutilide), Class IV agents (e.g., diltiazem and verapamil), 5HT antagonists (e.g., sulamserod, serraline and tropsetron), and dronedarone; calcium channel blockers (both L-type and T-type) such as diltiazem, verapamil, nifedipine, amlodipine and mybefradil; Cyclooxygenase inhibitors (i.e., COX-1 and/or COX-2 inhibitors) such as aspirin, indomethacin, ibuprofen, piroxicam, naproxen, celebrex, vioxx and NSAIDs; anti-platelet agents such as GPIIb/IIIa blockers (e.g., abciximab, eptifibatide and tirofiban), P2Y<sub>12</sub> antagonists (e.g., clopidogrel, ticlopidine and CS-747), thromboxane receptor antagonists (e.g., ifetroban), aspirin, and PDE-III inhibitors (e.g., dipyridamole) with or without aspirin; diuretics such as chlorothiazide, hydrochlorothiazide, flumethiazide, hydroflumethiazide, bendroflumethiazide, methylchlorothiazide, trichloromethiazide, polythiazide, benzthiazide, ethacrynic acid, tricyclic nafen, chlorthalidone, furosemide, musolimine,

bumetanide, triamterene, amiloride, and spironolactone; anti-hypertensive agents such as alpha adrenergic blockers, beta adrenergic blockers, calcium channel blockers, diuretics, renin inhibitors, ACE inhibitors, (e.g., captopril, zofenopril, fosinopril, enalapril, ceranopril, cilazopril, delapril, pentopril, quinapril, ramipril, lisinopril), A II antagonists (e.g., losartan, irbesartan, valsartan), ET antagonists (e.g. sitaxsentan, atrsentan and compounds disclosed in U.S. Patent Nos. 5,612,359 and 6,043,265), Dual ET/AII antagonist (e.g., compounds disclosed in WO 00/01389), neutral endopeptidase (NEP) inhibitors, vasopepsidase inhibitors (dual NEP-ACE inhibitors) (e.g., omapatrilat and gemopatrilat), nitrates, and combinations of such anti-

hypertensive agents; antithrombotic/thrombolytic agents such as tissue plasminogen activator (tPA), recombinant tPA, tenecteplase (TNK), lanoteplase (nPA), factor VIIa inhibitors, factor Xa inhibitors, thrombin inhibitors (e.g., hirudin and argatroban), PAI-1 inhibitors (i.e., inactivators of tissue plasminogen activator inhibitors),  $\alpha$ 2-antiplasmin inhibitors, streptokinase, urokinase, prourokinase, anisoylated plasminogen

streptokinase activator complex, and animal or salivary gland plasminogen activators; anticoagulants such as warfarin and heparins (including unfractionated and low molecular weight heparins such as enoxaparin and dalteparin); HMG-CoA reductase inhibitors such as pravastatin lovastatin, atorvastatin, simvastatin, NK-104 (a.k.a. itavastatin, or nisvastatin or nisbastatin) and ZD-4522 (a.k.a. rosuvastatin, or

atavastatin or visastatin); other cholesterol/lipid lowering agents such as squalene synthetase inhibitors, fibrates, and bile acid sequestrants (e.g., questran); antiproliferative agents such as cyclosporin A, taxol, FK 506, and adriamycin; antitumor agents such as taxol, adriamycin, epothilones, cisplatin and carboplatin; anti-diabetic agents such as biguanides (e.g. metformin), glucosidase inhibitors (e.g. acarbose), insulins, meglitinides (e.g. repaglinide), sulfonylureas (e.g. glimepiride, glyburide and glipizide), biguanide/glyburide combinations (i.e., glucovance), thiozolidinediones (e.g. troglitazone, rosiglitazone and pioglitazone), PPAR-gamma agonists,  $\alpha$ P2 inhibitors, and DP4 inhibitors; thyroid mimetics (including thyroid receptor antagonists) (e.g., thyrotropin, polythyroid, KB-130015, and dronedarone);

Mineralocorticoid receptor antagonists such as spironolactone and eplerinone; growth hormone secretagogues; anti-osteoporosis agents (e.g., alendronate and raloxifene); hormone replacement therapy agents such as estrogen (including conjugated estrogens

in premarin), and estradiol; antidepressants such as nefazodone and sertraline; antianxiety agents such as diazepam, lorazepam, buspirone, and hydroxyzine pamoate; oral contraceptives; anti-ulcer and gastroesophageal reflux disease agents such as famotidine, ranitidine, and omeprazole; anti-obesity agents such as orlistat; 5 cardiac glycosides including digitalis and ouabain; phosphodiesterase inhibitors including PDE III inhibitors (e.g. cilostazol), and PDE V inhibitors (e.g., sildenafil); protein tyrosine kinase inhibitors; steroidal anti-inflammatory agents such as prednisone, and dexamethasone; and other anti-inflammatory agents such as enbrel.

The above other therapeutic agents, when employed in combination with the 10 compounds of the present invention, may be used, for example, in those amounts indicated in the Physicians' Desk Reference (PDR) or as otherwise determined by one of ordinary skill in the art.


Assays to determine the degree of activity of a compound as an  $I_{Kur}$  inhibitor are well known in the art and are described in references such as *J. Gen. Physiol.* 15 Apr;101(4):513-43, and *Br. J. Pharmacol.* 1995 May;115(2):267-74.

Assays to determine the degree of activity of a compound as an inhibitor of other members of the  $K_v1$  subfamily are also well known in the art. For example, inhibition of  $K_v1.1$ ,  $K_v1.2$  and  $K_v1.3$  can be measured using procedures described by Grissmer S, et al., *Mol Pharmacol* 1994 Jun;45(6):1227-34. Inhibition of  $K_v1.4$  can 20 be measured using procedures described by Petersen KR, and Nerbonne JM, *Pflugers Arch* 1999 Feb;437(3):381-92. Inhibition of  $K_v1.6$  can be measured using procedures described by Bowlby MR, and Levitan IB, *J Neurophysiol* 1995 Jun;73(6):2221-9. And inhibition of  $K_v1.7$  can be measured using procedures described by Kalman K, et al., *J Biol Chem* 1998 Mar 6;273(10):5851-7.

25 Compounds within the scope of the present invention demonstrate activity in  $K_v1$  assays such as the ones described above.


All documents cited in the present specification are incorporated herein by reference in their entirety.

The following examples and preparations describe the manner and process of 30 making and using the invention and are illustrative rather than limiting. It is to be understood that there may be other embodiments which fall within the spirit and scope of the invention as defined by the claims appended hereto.

EXAMPLESEXAMPLE 1

5

2-Methoxy-N-(4-thiophen-2-yl-piperidin-4-ylmethyl)-benzamide

Synthesis

10

**Compound 1:** Compound 1 is commercially available.

**Compound 2:** Compound 1 (20 g, 0.11 mol) was suspended in 200 mL dichloromethane. Benzoylchloride (17 mL, 0.14 mol) was added. At 0 °C TEA (42 mL, 0.30mol) in dichloromethane (10 mL) was added slowly to the reaction mixture. The reaction mixture was stirred at room temperature for 14 h, diluted with Ethyl acetate (500 mL), washed with saturated NaHCO<sub>3</sub> (2 × 250 mL), 1N HCl (2 × 250 mL), dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was purified by silica gel chromatography using Hexanes/Ethyl acetate (4/1, 1/1) as eluent to give an orange oil compound 2, 25.85 g (94% yield).

**Compound 3:** NaH (6.0 g, 0.25 mol) was suspended in DMF (100 mL). At 0 °C 2-thiopheneacetonitrile (7.5 g, 0.061 mol) was added followed by the addition of compound **2** (12 g, 0.048 mol) in 100 mL DMF. The reaction was stirred at 0 °C for 0.5 h and then at ambient temperature for 14 h. The reaction mixture was poured into ice and extracted with Ethyl acetate (250 mL). The aqueous layer was extracted with Ethyl acetate (2 × 250 mL). The organic layers were combined and washed with 10% LiCl (2 × 200 mL), dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was purified by silica gel flash chromatography eluted with 2/1 and 1/1 Hexane:Ethyl acetate yielding 9.14 g (64% isolated yield) of compound **3** as a dark brown solid.

HPLC Rt 2.87 min, Purity 100%, YMC S5 column 4.6 × 50 mm, 4 min gradient 0 to 100% MeOH (90% in water, 0.2% PPA) UV detection at 220nm. LCMS Rt 1.47 min, [M+1] 297.22 Phenomenex S5 column 4.6 × 30 mm, 2 min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.74 – 2.32 ppm, 4H, m; 3.22– 3.43 ppm, 2H, m; 3.78 – 3.84 ppm, 1H, broadpeak; 4.79 ppm, 1H, broad peak; 6.94 ppm, 1H, dd, J=5.0 Hz and 3.6 Hz; 7.08 ppm, 1H, dd, J=3.3 Hz and 1.1 Hz; 7.24 ppm, 1H, dd, J=5.0 Hz and 1.1 Hz; 7.34 – 7.42 ppm, 5H, m.

**Compound 4:** To a solution of compound **3** (2.1 g, 7.2 mmol) in THF (40 mL) was added LAH (20 mL, 20 mmol, 1.0M solution in THF). The reaction mixture was heated to reflux for 1 h then allowed to cool to ambient temperature. The solution was cooled to 0 °C and quenched with water (5.3 mL), 1N NaOH (3.4 mL) and water (5.3 mL). The quenched reaction mixture was stirred at ambient temperature for 0.5 h. The slurry was filtered through a celite pad and the filtrate was concentrated. The residue was diluted with EtOAc (200 mL), washed with saturated NaHCO<sub>3</sub> (2 × 100 mL), dried over MgSO<sub>4</sub>, filtered and concentrated to give a brown oil compound **4** sufficiently pure to be taken on to the next step. HPLC Rt 0.20 min and 0.59 min, Purity 100%, YMC S5 column 4.6 × 50 mm, 4 min gradient 0 to 100% MeOH (90% in water, 0.2% PPA) UV detection at 220nm. LCMS Rt 0.73 min, [M+1] 287.39 Phenomenex S5 column 4.6 × 30 mm, 2min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.85 ppm, 2H, m; 2.05 – 2.08 ppm, 2H, m; 2.22 – 2.27 ppm, 2H, m; 2.67 ppm, 1H, s; 2.69 ppm, 2H,


m; 3.44 ppm, 2H, s; 6.77 – 6.78 ppm, 1H, m; 6.94 ppm, 1H, dd, J=5.0 Hz and 3.3 Hz; 7.17 – 7.30 ppm, 6H, m.

**Compound 5:** Compound **4** was dissolved in dichloromethane (30 mL) and ortho-anisoyl chloride (1.1 mL, 7.4 mmol) was added followed by the addition of  
5 TEA (2.3 mL, 16 mmol). The reaction was stirred at ambient temperature for 14 h, diluted with dichloromethane (200 mL), washed with saturated NaHCO<sub>3</sub>, dried over MgSO<sub>4</sub>, filtered and concentrated. The product amide compound **5** was eluted with 2:1 and 1:1 hexane:ethylacetate as a white foam (2.19g, 77% yield for two steps). HPLC Rt 2.50 min, Purity 100%, YMC S5 column 4.6 × 50 mm, 4 min gradient 0 to  
10 100% MeOH (90% in water, 0.2% PPA) UV detection at 220nm. LCMS Rt 1.36 min, [M+1] 421.27 Phenomenex S5 column 4.6 × 30 mm, 2 min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CD<sub>3</sub>OD) 1.98 – 2.00 ppm, 2H, m; 2.15 – 2.18 ppm, 2H, m; 2.44 ppm, 2H, m; 2.74 – 2.76 ppm, 2H, m; 3.53 ppm, 2H, s; 3.65 ppm, 2H, s; 3.77 ppm, 3H, s; 7.02 – 7.09 ppm, 4H, m; 7.25  
15 – 7.31 ppm, 5H, m; 7.39 – 7.40 ppm, 1H, m; 7.45 – 7.48 ppm, 1H, m; 7.93 – 7.95 ppm, 1H, m.

**Compound 6:** Compound **5** (2.2 g, 5.2 mmol) was dissolved in dichloroethane (40 mL) and TEA (3.6 mL, 26 mmol) was added. At 0 °C chloroethyl chloroformate (1.1 mL, 10 mmol) in dichloroethane (20 mL) was added. The reaction  
20 mixture was stirred at ambient temperature for 1 h, then another 1.1 mL chloroethyl chloroformate in dichloroethane (20 mL) was added. The reaction was stopped till no starting material left according to LC-MS. The reaction mixture was concentrated and dried on oil pump for 0.5 h. MeOH (40 mL) was added to the residue and was heated to reflux for 4 h. The reaction mixture was concentrated and the residue was purified  
25 by a silica gel pad eluted with 1:1 EtOAc:Hexane and then 1:1 MeOH:DCM yielding a pale yellow solid 2-methoxy-N-(4-thiophen-2-yl-piperidin-4-ylmethyl)-benzamide **6** (1.63 g, 95% yield). HPLC Rt 2.09 min, Purity 100%, YMC S5 column 4.6 × 50 mm, 4 min gradient 0 to 100% MeOH (90% in water, 0.2% PPA) UV detection at 220nm. LCMS Rt 1.20 min, [M+1] 331.41 Phenomenex S5 column 4.6 × 30 mm, 2 min  
30 gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 2.01 – 2.04 ppm, 2H, m; 2.14 – 2.18 ppm, 2H, m; 2.92 – 2.94 ppm, 2H,


m; 3.16 ppm, 2H, m; 3.64 ppm, 2H, s; 3.67 ppm, 3H, s; 6.82 – 6.86 ppm, 2H, m;  
6.96– 7.00 ppm, 2H, m; 7.23 – 7.25 ppm, 1H, m; 7.32 – 7.37 ppm, 1H, m; 8.09 –  
8.11 ppm, 1H, m.

5

EXAMPLE 2

4-[(2-Methoxy-benzoylamino)-methyl]-4-thiophen-2-yl-piperidine-  
1-carboxylic acid ethyl ester

10

Synthesis

15


**Compound 1:** Compound 1 was prepared using methodology described in Example 1.

**Compound 2:** Compound 1 (14 mg, 0.043 mmol) was dissolved in dichloromethane (0.20 mL). At ambient temperature TEA (12 uL, 0.086 mmol) was added followed by the addition of ethyl chloroformate (4.90 mg, 0.045 mmol) in dichloromethane (0.10 mL). The reaction mixture was stirred for 2.5 h and then concentrated. 4-[(2-Methoxy-benzoylamino)-methyl]-4-thiophen-2-yl-piperidine-1-carboxylic acid ethyl ester 2 (13.7 mg, 79% yield) was isolated by Prep-HPLC as a colorless oil. HPLC Rt 3.43 min, Purity 100%, YMC S5 column 4.6 × 50 mm, 4 min gradient 0 to 100% MeOH (90% in water, 0.2% PPA) UV detection at 220nm.


LCMS Rt 1.78 min, [M+1] 403.17 YMC S5 column 4.6 × 30 mm, 2 min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.18 ppm, 3H, t, J=7.1 Hz; 1.81 – 1.84 ppm, 2H, m; 2.00 – 2.04 ppm, 2H, m; 3.18 – 3.25 ppm, 2H, m; 3.67 ppm, 3H, s; 3.73 – 3.77 ppm, 4H, m; 4.05 ppm, 2H, q, J=7.1 Hz; 6.83 – 6.88 ppm, 2H, m; 6.98 – 7.02 ppm, 2H, m; 7.25 ppm, 1H, dd, J=5.0 Hz and 0.76 Hz; 7.34 – 7.39 ppm, 1H, m; 7.83 ppm, 1H, m; 8.10 ppm, 1H, dd, J=7.8 Hz and 1.7 Hz.

### EXAMPLE 3

Example 3 was prepared using methodology described in Example 2.


Example	Structure	Name	[M+1]
3		4-[(2-Methoxy-benzoylamino)-methyl]-4-thiophen-2-yl-piperidine-1-carboxylic acid tert-butyl ester	430

### EXAMPLE 4



2-Methoxy-N-[1-(propane-1-sulfonyl)-4-thiophen-2-yl-piperidin-4-ylmethyl]-benzamide

# Synthesis





5            **Compound 1:** Compound **1** was prepared using methodology described in Example 1.

**Compound 3:** Compound **1** (14 mg, 0.043 mmol) was dissolved in dichloromethane (0.20 mL). At ambient temperature TEA (12 uL, 0.086 mmol) was added followed by the addition of propyl sulfonylchloride (6.4 mg, 0.045 mmol) in  
 10 dichloromethane (0.10 mL). The reaction mixture was stirred for 1 h and then concentrated. 2-Methoxy-N-[1-(propane-1-sulfonyl)-4-thiophen-2-yl-piperidin-4-ylmethyl]-benzamide (14.5 mg, 77% yield) was isolated by Prep-HPLC as a colorless oil. HPLC Rt 3.21 min, Purity 100%, YMC S5 column 4.6 × 50 mm, 4 min gradient 0 to 100% MeOH (90% in water, 0.2% PPA) UV detection at 220nm. LCMS Rt 1.67  
 15 min, [M+1] 437.15 YMC S5 column 4.6 × 30 mm, 2 min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 0.97 ppm, 3H, t, J=7.5 Hz; 1.73 – 1.79 ppm, 2H, m; 1.92 – 1.98 ppm, 2H, m; 2.09 – 2.14 ppm, 2H, m; 2.78 – 2.82 ppm, 2H, m; 3.19 – 3.25 ppm, 2H, m; 3.40 – 3.45 ppm, 2H, m; 3.66 ppm, 3H, s; 3.71 ppm, 2H, d, J=6.2 Hz; 6.84 – 6.88 ppm, 2H, m; 6.99 – 7.03 ppm,  
 20 2H, m; 7.27 ppm, 1H, dd, J=5.0 Hz and 0.74 Hz; 7.37 – 7.41 ppm, 1H, m; 7.98 ppm, 1H, m; 8.07 ppm, 1H, dd, J=7.8 Hz and 1.8 Hz.


EXAMPLES 5 AND 6

Examples 5 and 6 were synthesized using methodology described in Example

4.


Example	Structure	Name	[M+1]
5		N-(1-Benzenesulfonyl-4-thiophen-2-yl-piperidin-4-ylmethyl)-2-methoxybenzamide	470
6		N-[1-(4-Fluoro-benzenesulfonyl)-4-thiophen-2-yl-piperidin-4-ylmethyl]-2-methoxybenzamide	488

5

EXAMPLE 7

2-Methoxy-N-[4-thiophen-2-yl-1-(2,2,2-trifluoro-ethylsulfamoyl)-piperidin-4-ylmethyl]-benzamide

10

Synthesis

5        **Compound 1:** Compound 1 is commercially available.

**Compound 2:** Compound 2 is commercially available.

**Compound 3:** Compound 1 (26 mg, 0.18 mmol) was dissolved in dichloromethane (0.50 mL). At 0 °C compound 2 (15 mg, 0.18 mmol) in dichloromethane (0.50 mL) was added and the reaction mixture was stirred for 1 h.

10    Trifluoroethylamine (18 mg, 0.18 mmol) in dichloromethane (0.50 mL) was added followed by the addition of TEA (75 uL, 0.54 mmol). The reaction mixture was stirred at ambient temperature for 2 h and then at 35 °C for 14 h. The reaction was diluted with dichloromethane (20 mL), washed with 1N HCl (10 mL), dried over  $\text{MgSO}_4$ , filtered and concentrated to yield compound 3 sufficient pure to be taken to

15    the next step.

**Compound 4:** Compound 4 was prepared using methodology described in Example 1.






**Title Compound:** Compound 4 (20 mg, 0.060 mmol) and TEA (130 uL, 0.94 mmol) were dissolved into acetonitrile (1.0 mL). To this mixture compound 3 was

20    added and the reaction mixture was heated at 95 °C for 14 h. The reaction was concentrated and the residue was purified by PrepHPLC yielding 2-methoxy-N-[4-thiophen-2-yl-1-(2,2,2-trifluoro-ethylsulfamoyl)-piperidin-4-ylmethyl]-benzamide (17.2 mg, 58% yield) as a clear oil. HPLC Rt 3.25 min, Purity 100%, YMC S5 column 4.6 × 50 mm, 4 min gradient 0 to 100% MeOH (90% in water, 0.2% PPA)


UV detection at 220nm. LCMS Rt 1.66 min, [M+1] 492.14 YMC S5 column 4.6 × 30 mm, 2 min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.94 – 1.96 ppm, 2H, m; 2.01 – 2.12 ppm, 2H, m; 3.11 – 3.13 ppm, 2H, m; 3.43 ppm, 2H, m; 3.56 – 3.59 ppm, 2H, m; 3.64 ppm, 2H, s; 3.66 ppm, 3H, s; 4.95 ppm, 1H, t, J=5.7 Hz; 6.85 – 6.86 ppm, 2H, m; 6.99 – 7.00 ppm, 2H, m; 7.25 – 7.26 ppm, 1H, m; 7.35 – 7.36 ppm, 1H, m; 7.75 – 7.78 ppm, 1H, m; 8.12 ppm, 1H, dd, J=6.2 Hz and 1.3 Hz.

### EXAMPLES 8 TO 14

10 Examples 8 to 14 were prepared using methodology described in Example 7.


Example	Structure	Name	[M+1]
8		N-(1-Dimethylsulfamoyl-4-thiophen-2-yl-piperidin-4-ylmethyl)-2-methoxy-benzamide	437
9		N-[1-(4-Fluoro-benzylsulfamoyl)-4-thiophen-2-yl-piperidin-4-ylmethyl]-2-methoxy-benzamide	517
10		(4-Fluoro-benzyl)-carbamic acid 2-{4-[(2-methoxy-benzoylamino)-methyl]-4-thiophen-2-yl-piperidine-1-sulfonylamino}-ethyl ester	604
11		2-Methoxy-N-(1-phenylsulfamoyl-4-thiophen-2-yl-piperidin-4-ylmethyl)-benzamide	485
12		2-Methoxy-N-(1-methylsulfamoyl-4-thiophen-2-yl-piperidin-4-ylmethyl)-benzamide	423

Example	Structure	Name	[M+1]
13		N-{1-[1-(4-Fluoro-phenyl)-ethylsulfamoyl]-4-thiophen-2-yl-piperidin-4-ylmethyl}-2-methoxy-benzamide	531
14		2-Methoxy-N-(1-propylsulfamoyl-4-thiophen-2-yl-piperidin-4-ylmethyl)-benzamide	451

EXAMPLE 15

5                    2-Methoxy-N-(4-phenyl-1-sulfamoyl-piperidin-4-ylmethyl)-benzamide

# Synthesis



5        **Compound 1:** Compound 1 is commercially available.

**Compound 2:** To 1-N-benzyl-4-phenyl-4-cyanopiperidine.HCl (10.0 g, 31.97 mmol) was added 220 mL of THF and the reaction flask was cooled to 0°C. Lithium aluminum hydride (4.85 g, 127.86 mmol) was added slowly and the reaction mixed 12 h at room temperature. The reaction was quenched by the addition of 5 mL of water,  
 10    15 mL of 15% aqueous solution of sodium hydroxide followed by 5 mL of water. The organic fraction was dried over anhydrous sodium sulfate, filtered and the filtrate was concentrated to give compound 2 that was used without further purification. LCMS Rt 1.70 min, [M+1] 281.0.


**Compound 3:** To a solution of 1-N-benzyl-4-phenyl-4-  
 15    aminomethylpiperidine (4.0 g, 14.26 mmol) and triethylamine (3.0 g, 21.39 mmol) in 18 mL of dichloromethane was added  $o$ -anisoyl chloride (425  $\mu\text{L}$ , 2.85 mmol) at 0°C. The reaction was stirred for 12 h and quenched with (100 mL) of 1 M hydrochloric acid. Dichloromethane (100 mL) was also added and the aqueous layer was washed with dichloromethane (50 mL portions, 2x). The organic fractions were combined  
 20    and washed with 1 N sodium hydroxide (50 mL portions, 2x) followed by brine (100

mL). The organics were dried over anhydrous sodium sulfate and concentrated. 4.54 g (77% yield) of compound **3** was obtained. LCMS Rt 1.41 min, [M+1] 415.1.

**Compound 4:** To a solution of N-(1-benzyl-4-phenyl-piperidin-4-ylmethyl)-benzamide (4.54 g, 10.95 mmol) in ethanol (100 mL) was added 10% palladium/carbon (1.40 g). The reaction mixture was stirred in a hydrogen atmosphere (50 psi) for 78 h. After filtration, the filtrate was concentrated and purified using column chromatography on silica gel using 9:1:0.1 chloroform:methanol:ammonium hydroxide as the eluent to give a yellow oil. After lyophilization 2.7 g (76 % yield) of compound **4** as a white/yellow powdery solid was obtained. LCMS Rt 1.43 min, [M+1] 325.3.


**Title Compound:** To compound **4** (50 mg, 0.154 mmol) in 1,4-dioxane (1.7 mL) was added sulfamide (148 mg, 1.54 mmol) and then stirred at 100°C overnight. The solution was cooled to room temperature and the solvent was concentrated under reduced pressure. The crude material was diluted with dichloromethane (10 mL) and washed with water (10 mL) (2x), brine (10 mL) (1x) and dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under reduced pressure to give a white solid. The crude was purified using preparative HPLC and lyophilized to give 54 mg (87% yield) of 2-methoxy-N-(4-phenyl-1-sulfamoyl-piperidin-4-ylmethyl)-benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, rt): δ ppm) 2.11-2.19 (2 H, m), 2.22-2.25 (2 H, m), 3.24-3.26 (2 H, m), 3.35-3.38 (2 H, m), 3.57 (3 H, s), 3.67 (2 H, d, J = 6 Hz), 4.43 (2 H, s), 6.86 (1 H, d, J = 8.3 Hz), 7.05 (1 H, t, J = 7.5 Hz), 7.30-7.45 (6 H, m), 7.63 (1H, s), 8.16 (1 H, d, J = 7.5). LCMS Rt 1.50 min, [M+1] 404.2.

#### EXAMPLE 16




N-(1-Dimethylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-methoxy-benzamide

## Synthesis




5      **Compound 1:** Compound **1** was prepared as described in Example 15.

**Title Compound:** To compound **1** (35 mg, 0.108 mmol) in tetrahydrofuran (1.5 mL) was added polystyrene-diisopropylethylamine resin (394 mg, 1.4 mmol) and dimethylsulfamoyl chloride (35  $\mu$ L, 0.324 mmol). The reaction was mixed overnight at room temperature. Excess dimethylsulfamoyl chloride (35  $\mu$ L, 0.324 mmol) was added to the reaction to drive it to completion. To the reaction was added AP-Trisamine (232 mg, 0.972 mmol) and the reaction mixed for 6 hr at room temperature. After filtration, the solvent was concentrated under reduced pressure. The crude material was purified using the preparative HPLC and lyophilized to give N-(1-dimethylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-methoxy-benzamide 34.5 mg (73% yield).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , rt):  $\delta$  ppm) 1.94-1.99 (2 H, m), 2.01-2.67 (2 H, m), 2.79 (6 H, s), 3.16-3.24 (2 H, m), 3.42-3.5 (2 H, m), 3.58 (3 H, s), 3.74 (2 H, d,  $J = 6.3$  Hz), 6.86 (1 H, d,  $J = 8.1$  Hz), 7.05 (1 H, t,  $J = 7.5$  Hz), 7.29-7.46 (6 H, m), 7.59 (1 H, s), 8.18 (1 H, dd,  $J = 1.7, 7.8$  Hz). LCMS Rt 1.36 min,  $[\text{M}+1]$  432.3.

EXAMPLE 17

2-Methoxy-N-[1-(2-methoxy-ethylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-benzamide

5

Synthesis

10

**Compound 1:** Compound 1 is commercially available.

**Compound 2:** Compound 2 is commercially available.

**Compound 3:** Compound 3 was prepared using methodology described in Example 7 using 2-methoxy-ethylamine instead of 2,2,2-trifluoro-ethylamine.

15

**Compound 4:** Compound 4 was as described in Example 15.

**Title Compound:** 2-Methoxy-N-[1-(2-methoxy-ethylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-benzamide was prepared using methodology described in

Example 7.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , rt):  $\delta$  ppm) 1.95-2.03 (2 H, m), 2.2-2.29 (2 H, m), 3.12-3.22 (4 H, m), 3.30 (3 H, s), 3.41-3.48 (4 H, m), 3.58 (3 H, s), 3.73 (2 H, d,  $J = 6.1$  Hz), 4.55 (1 H, t,  $J = 5.8, 11.7$  Hz), 6.86 (1 H, d,  $J = 8.2$  Hz), 7.04 (1 H, dt,  $J = 0.9, 7.5$  Hz), 7.28-7.45 (6 H, m), 7.59 (1 H, t,  $J = 5.7, 8.65$  Hz), 8.17 (1 H, dd,  $J = 2.0, 7.85$  Hz). LCMS Rt 1.36 min,  $[\text{M}+1]$  462.


### EXAMPLES 18 TO 22


Examples 18 to 22 were prepared using methodology described in Example

17.


10

Example	Structure	Name	M+H
18		N-(1-Benzylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-methoxy-benzamide	495
19		2-Methoxy-N-(4-phenyl-1-propylsulfamoyl-piperidin-4-ylmethyl)-benzamide	447
20		N-[1-(4-Fluoro-benzylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	513
21		N-(1-Allylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-methoxy-benzamide	445

Example	Structure	Name	M+H
22		N-[1-(2-Hydroxy-ethylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	449

EXAMPLE 23

- 5                    2-Methoxy-N-[1-(2-oxo-oxazolidine-3-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-benzamide

Synthesis

10


**Compound 1:** Compound **1** was prepared as described in Example 15.

**Title Compound:** A solution of chlorosulfonyl isocyanate (0.12 mL; 1.4 mmol) in dichloromethane (20 mL) was cooled to 0°C and treated with 2-chloroethanol (0.094 mL; 1.3 mmol). After 2 h a solution of compound **1** (0.42g; 1.3 mmol) and triethylamine (0.72 mL; 5.2 mmol) in dichloromethane (15 mL) was added dropwise. When the addition was complete the cooling bath was removed and the reaction mixture was allowed to stir at room temperature for 24 h. 20% aqueous hydrochloric acid was added and the organic layer was separated, washed with

saturated aqueous sodium chloride, dried (anhydrous sodium sulfate), filtered and concentrated. The residue was purified by column chromatography on silica gel using 9:1 ethyl acetate:hexane as the eluent to give 0.3 g of 2-methoxy-N-[1-(2-oxo-oxazolidine-3-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-benzamide as a white solid.


5 LCMS  $m/z = 475$  (M+H)<sup>+</sup>

#### EXAMPLE 24



10

#### Synthesis



15


**Compound 1:** Compound 1 is commercially available.

**Compound 2:** Trifluoroethylamine (6.6 mg, 0.067 mmol) and compound 1 (16 mg, 0.067 mmol) were added into iPrOH (1.0 mL). The reaction mixture was heated at 95 °C for 6 h.


**Compound 3:** Compound 3 was prepared as described in Example 1.

**Title Compound:** Compound **3** (20 mg, 0.060 mmol) in acetonitrile (0.50 mL) was added to compound **2** in iPrOH.. The reaction was heated at 95 °C for 14 h and concentrated. The residue was purified through Prep HPLC to yield the title compound (4.4 mg, 15% yield) as a clear oil. HPLC Rt 3.13 min, Purity 91%, YMC S5 column 4.6 × 50 mm, 4 min gradient 0 to 100% MeOH (90% in water, 0.2% PPA) UV detection at 220nm. LCMS Rt 1.62 min, [M+1] 480.18 YMC S5 column 4.6 × 30 mm, 2 min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.91 – 1.99 ppm, 2H, m; 2.12 – 2.16 ppm, 2H, m; 3.40 – 3.47 ppm, 2H, m; 3.65 – 3.68 ppm, 5H, m; 3.77 – 3.83 ppm, 2H, m; 3.94 – 3.97 ppm, 2H, m; 5.32 ppm, 1H, m; 6.85 – 6.87 ppm, 2H, m; 6.98 – 7.02 ppm, 2H, m; 7.27 ppm, 1H, dd, J=5.0 Hz and 0.8 Hz; 7.35 – 7.40 ppm, 1H, m; 7.82 ppm, 1H, m; 8.06 – 8.09 ppm, 1H, m.

### EXAMPLE 25



### Scheme



**Compound 1:** Compound **1** was prepared as described in Example 15.

**Compound 2:** A solution of compound **1** (0.63 g; 1.9 mmol) in anhydrous acetonitrile (20 mL) was treated with diphenyl *N*-cyanocarbonimidate (0.95 g; 4.0




mmol) and the reaction mixture was heated to 85°C. After 19 h the acetonitrile was removed by evaporation and the residue was portioned between ethyl acetate and saturated aqueous sodium bicarbonate. The organic layer was separated, washed with saturated aqueous sodium chloride, dried (anhydrous sodium sulfate), filtered and concentrated. Column chromatography on silica gel using 7:3 ethyl acetate:hexane as the eluent gave 0.89 g of compound 2 as a white solid. LCMS  $m/z = 470$  (M+H)<sup>+</sup>








**Title Compound:** Compound 3 (0.1 g; 0.2 mmol) was treated with 7 N ammonia in methanol (1.5 mL) and heated to 45°C in a sealed tube for 1 h. The methanol and ammonia was removed by evaporation and the residue was portioned between ethyl acetate and 1 N sodium hydroxide. The organic layer was separated, washed with saturated aqueous sodium chloride, dried (anhydrous sodium sulfate), filtered and concentrated. The crude product was purified by recrystallization from ethyl acetate to give 0.064 g of the title compound as white crystals. LCMS  $m/z = 392$  (M+H)<sup>+</sup>








15

### EXAMPLES 26 TO 58

Examples 26 to 58 were prepared using methodology described in Example 24 and Example 25.


Example	Structure	Name	[M+1]
26			519
27			473
28			439

Example	Structure	Name	[M+1]
29			411
30			392
31			421
32			501
33			407
34			435
35			449


Example	Structure	Name	[M+1]
36			461
37			433
38			437
39			447
40			431
41			515
42			447

Example	Structure	Name	[M+1]
43			463
44			435
45			439
46			465
47			426
48			426
49			421
50			408

Example	Structure	Name	[M+1]
51			391
52			484
53			460
54			495
55			511
56			445
57			463
58			487

EXAMPLE 59

5

Synthesis

10 **Compound 1:** Compound 1 was prepared as described in Example 1.

**Compound 2:** To a solution of compound 1 (3.5 g, 12 mmol) in dry THF (70 mL) was added LAH (35 mL, 35 mmol, 1.0M solution in tetrahydrofuran). The reaction mixture was heated to reflux for 1.5 h then allowed to cool to ambient temperature. The solution was cooled to 0 °C and quenched with water (9.2 mL), 1N  
 15 NaOH (5.8 mL) and water (9.2 mL). The quenched reaction mixture was stirred at

ambient temperature for 0.5 h. The slurry was filtered through a celite pad and the filtrate was concentrated. The residue was diluted with EtOAc (200 mL), washed with saturated NaHCO<sub>3</sub> (2 × 100 mL), dried over MgSO<sub>4</sub>, filtered and concentrated to give a brown oil. This residue was dissolved in dichloromethane (75 mL) and

5 carbonic acid di-tert-butyl ester (2.9 g, 13 mmol) and TEA (1.8 mL, 13 mmol) were added subsequently. The reaction mixture was stirred at ambient temperature for 18 h and then concentrated. The product compound **2** was purified by flash silica gel chromatography elution with 2:1 hexane:ethylacetate (3.5g, 75% yield for two steps). HPLC Rt 2.44 min, Purity 100%, YMC S5 column 4.6 × 50 mm, 4 min gradient 0 to

10 100% MeOH (90% in water, 0.2% PPA) UV detection at 220nm. LCMS Rt 1.39 min, [M+1] 387.24 Phenomenex S5 column 4.6 × 30 mm, 2 min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.32 ppm, 9H, s; 1.80 ppm, 2H, m; 1.96 ppm, 2H, m; 2.22 ppm, 2H, m; 2.55 – 2.58 ppm, 2H, m; 3.21 ppm, 2H, d, J = 6.3 Hz; 3.35 ppm, 2H, s; 4.30 ppm, 1H, m; 6.76 ppm,

15 1H, dd, J = 0.8 Hz and 3.4 Hz; 6.90 – 6.91 ppm, 1H, m; 7.13 – 7.21 ppm, 6H, m.

**Compound 3:** Compound **2** (1.7 g, 4.5 mmol) was dissolved in dichloroethane (35 mL) and TEA (3.1 mL, 22.4 mmol) was added. At 0 °C chloroethyl chloroformate (0.97 mL, 8.9 mmol) in dichloroethane (17 mL) was added. The reaction mixture was stirred at ambient temperature for 1 h, then another 1.0 mL

20 chloroethyl chloroformate and 3.0 mL TEA were added. The reaction mixture was stirred at ambient temperature for a further 2h then concentrated and dried under high vacuum for 0.5 h. MeOH (20 mL) was added to the residue and was heated to reflux for 4 h. The reaction mixture was concentrated and the residue was taken directly to the next step.

25 **Compound 4:** Compound **3** was dissolved in dichloromethane (60 mL). At ambient temperature TEA (1.9 mL, 13.4 mmol) was added followed by the addition of 4-Fluoro-benzenesulfonyl chloride (1.0 g, 5.4 mmol). The reaction mixture was stirred for 18 h and then concentrated. Compound **4** (1.44 g, 71% yield) was isolated as a white solid by column chromatography using 2:1 Hexane:EtOAc. HPLC Rt 3.56

30 min, Purity 100%, YMC S5 column 4.6 × 50 mm, 4 min gradient 0 to 100% MeOH (90% in water, 0.2% PPA) UV detection at 220nm. LCMS Rt 1.82 min, [M+23]





477.13 YMC S5 column 4.6 × 30 mm, 2 min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.32 ppm, 9H, s; 1.86 – 1.88 ppm, 2H, m; 2.03 – 2.06 ppm, 2H, m; 2.70 – 2.74 ppm, 2H, m; 3.13 ppm, 2H, d, J = 6.6 Hz; 3.40 – 3.42 ppm, 2H, m; 4.30 ppm, 1H, m; 6.70 ppm, 1H, d, J = 3.3 Hz; 5 6.87 – 6.89 ppm, 1H, m; 7.07 – 7.10 ppm, 1H, m; 7.13 – 7.14 ppm, 1H, m; 7.65 – 7.68 ppm, 2H, m.


**Compound 5:** Compound 4 (55 mg, 0.12 mmol) was dissolved in dichloromethane (1.3 mL). To this solution TFA (180 μL, 2.3 mmol) in dichloromethane (0.50 mL) was added and the reaction was stirred at ambient 10 temperature till no starting material was detected according to LC-MS. The reaction mixture was concentrated and neutralized with TEA. To the concentrated residue was added diphenyl cyanocarbonimide (32 mg, 0.13 mmol) and isopropyl alcohol (1.0 mL). The reaction mixture was heated to reflux for 18 h.

**Title Compound:** Cyclohexyl amine (19 mg, 0.19 mmol) was added to 15 compound 5 and the reaction was heated at 95 °C for 24h. The reaction mixture was concentrated and purified by Prep-HPLC YMC ODS S5 20x100mm, 16 min gradient 40 to 100% MeOH (90% in water 0.1%TFA) at 20mL/min UV detection at 220nm to give the title compound as a yellow oil (23.8 mg, 39% yield). HPLC Rt 3.49 min, Purity 100%, YMC S5 column 4.6 × 50 mm, 4 min gradient 0 to 100% MeOH (90% 20 in water, 0.2% PPA) UV detection at 220nm. LCMS Rt 1.81 min, [M+1] 504.24 YMC S5 column 4.6 × 30 mm, 2 min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.01 – 1.24 ppm, 5H, m; 1.56 – 1.58 ppm, 1H, m; 1.66 – 1.77 ppm, 4H, m; 1.91 – 1.94 ppm, 2H, m; 2.15 – 2.18 ppm, 2H, m; 2.75 – 2.79 ppm, 2H, m; 2.98 – 3.01 ppm, 1H, m; 3.27 ppm, 2H, d, J = 6.1 Hz; 25 3.45 – 3.47 ppm, 2H, m; 6.81 ppm, 1H, dd, J = 3.9 Hz and 1.1 Hz; 6.98 ppm, 1H, dd, J = 3.3 Hz and 4.9 Hz; 7.13 – 7.19 ppm, 2H, m; 7.25 – 7.29 ppm, 1H, m; 7.71 – 7.76 ppm, 2H, m.


EXAMPLES 60 TO 63

Examples 60 to 63 were prepared as described in Example 59.

Example	Structure	Name	[M+1]
60			435
61			477
62			541
63			505

EXAMPLE 64

5

Synthesis

**Compound 1:** Compound 1 was prepared using methodology described in  
 10 Example 59.







**Compound 2:** To the solution of compound 1 (500 mg, 1.1 mmol) in dichloromethane (5.0 mL) was added TFA (3.0 mL) in dichloromethane (12 mL) and the reaction was stirred for 1.5 h. Then the mixture was concentrated and neutralized with TEA. At 0 °C 2-chloroethanol (74  $\mu$ L, 1.1 mmol) in dichloromethane (7.5 mL)

was added to the solutions of chlorosulfonyl isocyanate (96  $\mu$ L, 1.1 mmol) in dichloromethane (7.5  $\mu$ L). The reaction mixture was stirred at 0 °C for 1 h. The crude amine in dichloromethane (5.0 mL) and TEA (460  $\mu$ L, 3.3 mmol) were added. The reaction was stirred for 18 h and concentrated. Compound **2** (260 mg, 47% yield) was purified as a white solid by column chromatography using 1:1 Hexane:EtOAc. HPLC Rt 2.86 min, Purity 100%, YMC S5 column 4.6  $\times$  50 mm, 4 min gradient 0 to 100% MeOH (90% in water, 0.2% PPA) UV detection at 220nm. LCMS Rt 1.48 min, [M+1] 504.08 YMC S5 column 4.6  $\times$  30 mm, 2 min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.86 – 1.96 ppm, 2H, m; 2.14 – 2.17 ppm, 2H, m; 2.56 – 2.62 ppm, 2H, m; 3.07 ppm, 2H, d, J = 6.9 Hz; 3.51 – 3.53 ppm, 2H, m; 3.90 ppm, 2H, t, J = 7.8 Hz; 4.32 ppm, 2H, t, J = 7.8 Hz; 5.17 ppm, 1H, t, J = 6.9 Hz; 6.75 ppm, 1H, d, J = 3.4 Hz; 6.88 – 6.90 ppm, 1H, m; 7.07 – 7.09 ppm, 2H, m; 7.11 – 7.18 ppm, 1H, m; 7.63 – 7.67 ppm, 2H, m.


**Title Compound:** Cyclohexyl amine (7.7 mg, 0.078 mmol), TEA (130  $\mu$ L) and compound **2** (26 mg, 0.052 mmol) were dissolved in acetonitrile (1.0 mL). The reaction was heated at 85 °C for 18 h and concentrated. The title compound was purified as an orange oil (24.1 mg, 90% yield) by Prep-HPLC YMC ODS S5 20x100mm, 16 min gradient 40 to 100% MeOH (90% in water 0.1%TFA) at 20mL/min UV detection at 220nm. HPLC Rt 3.51 min, Purity 100%, YMC S5 column 4.6  $\times$  50 mm, 4 min gradient 0 to 100% MeOH (90% in water, 0.2% PPA) UV detection at 220nm. LCMS Rt 1.82 min, [M+1] 516.16 YMC S5 column 4.6  $\times$  30 mm, 2 min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.05 – 1.11 ppm, 3H, m; 1.16 – 1.21 ppm, 2H, m; 1.49 – 1.51 ppm, 1H, m; 1.59 – 1.63 ppm, 2H, m; 1.77 – 1.80 ppm, 2H, m; 1.88 – 1.92 ppm, 2H, m; 2.15 – 2.18 ppm, 2H, m; 2.62 – 2.68 ppm, 2H, m; 2.94 – 2.98 ppm, 3H, m; 3.43 – 3.45 ppm, 2H, m; 3.84 ppm, 1H, m; 6.73 – 6.74 ppm, 1H, m; 6.88 – 6.89 ppm, 1H, m; 7.01 – 7.11 ppm, 2H, m; 7.17 ppm, 1H, dd, J = 1.1 Hz and 5.5 Hz; 7.65 – 7.67 ppm, 2H, m.

EXAMPLES 65 TO 73


Examples 65 to 73 were prepared using methodology described in Example 64.

Example	Structure	Name	[M+1]
65			447
66			489
67			515
68			487
69			539
70			509

Example	Structure	Name	[M+1]
71			553
72			523
73			503

EXAMPLE 74

5 4-benzylcarbamoyl-4-phenylpiperidine-1-carboxylic acid tert-butyl ester

Synthesis

10

**Compound 1:** Compound 1 is commercially available.

**Compound 2:** A solution of compound 1 (12.6 g; 33.4 mmol) in 100 mL of 1 N sodium hydroxide and 25 mL of tetrahydrofuran was treated with di-*tert*-butyl







dicarbonate (10.3g; 47.2 mmol) at room temperature. After stirring for 20 h, 10% aqueous hydrochloric acid was slowly added to neutralize the reaction mixture to pH = 7. Ethyl acetate (approximately 300 mL) was added and the organic layer was separated, washed with saturated aqueous sodium chloride, dried (anhydrous sodium sulfate), filtered and concentrated to give compound **2** as a white solid (10.2 g) that was used without further purification. LCMS  $m/z = 306$  (M+H)<sup>+</sup>





**Title Compound:** A solution of compound **2** (0.92 g; 3.0 mmol) in tetrahydrofuran (30 mL) was treated with benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (1.77 g; 4.0 mmol) and triethylamine (0.63 mL; 4.5 mmol) at room temperature. The reaction mixture was allowed to stir for 0.5 h at which time benzyl amine (0.39 mL; 3.6 mmol) was added and the reaction mixture was heated to 45°C for 2 h. The tetrahydrofuran was removed by evaporation and the residue was treated with ethyl acetate (approximately 150 mL) and 5% aqueous hydrochloric acid (approximately 100 mL). The organic layer was separated, washed with saturated aqueous sodium chloride, dried (sodium sulfate), filtered and concentrated. Column chromatography on silica gel using 7:3 hexane:ethyl acetate as the eluent gave 0.93 g of 4-benzylcarbamoyl-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester as a white solid. LCMS  $m/z = 396$  (M+H)<sup>+</sup>


20

EXAMPLES 75 TO 84

Examples 75 to 84 were prepared as described in Example 74.


Example	Structure	Name	M+H
75		4-(2-Methoxy-benzylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester	425
76		4-[2-(4-Methoxy-phenyl)-ethylcarbamoyl]-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester	439
77		4-(2,4-Dimethoxy-benzylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester	455
78		4-[2-(2,6-Dichloro-phenyl)-ethylcarbamoyl]-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester	478
79		4-Phenyl-4-(3-phenyl-propylcarbamoyl)-piperidine-1-carboxylic acid tert-butyl ester	423
80		4-Pentylcarbamoyl-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester	375

Example	Structure	Name	M+H
81		4-(3-Methoxy-propylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester	377
82		4-(4-Methoxy-benzylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester	425
83		4-(3,4-Difluoro-benzylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester	431
84		4-(4-Fluoro-benzylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester	413

EXAMPLE 85

5

4-phenyl-piperidine-4-carboxylic acid benzylamide

Synthesis







**Compound 1:** Compound 1 was prepared as described in Example 74.



**Title Compound:** A solution of compound 1 (0.93 g; 2.4 mmol) in 30 mL of dichloromethane was treated with 4 mL trifluoroacetic acid at room temperature. After stirring for 20 h, an additional 100 mL of dichloromethane was added followed by 100 mL 1 N sodium hydroxide. The organic layer was separated, washed with water and saturated aqueous sodium chloride, dried (anhydrous sodium sulfate), filtered and concentrated to give 0.67 g of 4-phenyl-piperidine-4-carboxylic acid benzylamide as a colorless oil that was used without further purification. LCMS  $m/z = 295$  (M+H)<sup>+</sup>


10

### EXAMPLES 86 TO 93

Examples 86 to 93 were prepared as described in Example 85.


Example	Structure	Name	M+H
86		4-Phenyl-piperidine-4-carboxylic acid 2,4-dimethoxy-benzylamide	355
87		(4-Phenyl-piperazin-1-yl)-(4-phenyl- piperidin-4-yl)-methanone	350
88		4-Phenyl-piperidine-4-carboxylic acid 2-methoxy-benzylamide	325
89		4-Phenyl-piperidine-4-carboxylic acid 4-methoxy-benzylamide	325
90		4-Phenyl-piperidine-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide	339
91		4-Phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	323

Example	Structure	Name	M+H
92		4-Phenyl-piperidine-4-carboxylic acid pentylamide	275
93		4-Phenyl-piperidine-4-carboxylic acid (3-methoxy-propyl)-amide	277

EXAMPLE 94

5

4-Phenyl-1-(2-phenyl-cyclopropanecarbonyl)-piperidine-4-carboxylic acid benzylamide

Synthesis

10


**Compound 1:** Compound 1 was prepared as described in Example 85.

15

**Title Compound:** Compound 1 (0.015 g; 0.05 mmol) was dissolved in 1 mL acetonitrile. Polystyrene-diisopropylethylamine (PS-DIEA) resin (0.1g) was added and the resulting suspension was treated with *trans*- 2-phenyl-cyclopropanecarbonyl chloride (0.02 g; 0.1 mmol) and shaken at room temperature. After 21 h, polystyrene-


trisamine (PS-trisamine) resin (0.1 g) was added and the reaction mixture was allowed to shake an additional 24 h. The reaction mixture was filtered and concentrated to give 0.015 g of 4-phenyl-1-(2-phenyl-cyclopropanecarbonyl)-piperidine-4-carboxylic acid benzylamide as a colorless oil. LCMS  $m/z = 440$  (M+H)<sup>+</sup>.

5

EXAMPLE 95

1-(3-Fluoro-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide


10

Synthesis

15


**Compound 1:** Compound **1** was prepared as described in Example 85.

**Title Compound:** Compound **1** (0.015 g; 0.05 mmol) was dissolved in 1 mL anhydrous acetonitrile. Polystyrene-diisopropylethylamine (PS-DIEA) resin (0.1g) was added and the resulting suspension was treated 3-fluoro-benzenesulfonyl chloride (0.02 g; 0.1 mmol) and shaken at room temperature. After 21 h, polystyrene-  
 20 trisamine (PS-trisamine) resin (0.1 g) was added and the reaction mixture was allowed to shake an additional 24 h. The reaction mixture was filtered and concentrated to give 0.012 g of 1-(3-fluoro-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide as a colorless oil. LCMS  $m/z = 454$  (M+H)<sup>+</sup>

EXAMPLE 96

4-Benzylcarbamoyl-4-phenyl-piperidine-1-carboxylic acid ethyl ester

5








Synthesis









10      **Compound 1:** Compound **1** was prepared as described in Example 85.









**Title Compound:** 4-Benzylcarbamoyl-4-phenyl-piperidine-1-carboxylic acid ethyl ester was prepared using methodology described in Example 2. LCMS  $m/z$  = 367 ( $M+H$ )<sup>+</sup>







EXAMPLES 97 TO 269







Examples 97 to 269 were synthesized using methodology described in Example 94, Example 95 and Example 96.








Example	Structure	Name	M+H
97		4-phenyl-1-(2-phenyl-cyclopropanecarbonyl)-piperidine-4-carboxylic acid 2,4-dimethoxy-benzylamide	500
98		1-(4-Methoxy-benzoyl)-4-phenyl-piperidine-4-carboxylic acid 2,4-dimethoxy-benzylamide	490
99		1-[2-(4-Fluoro-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid 2,4-dimethoxy-benzylamide	492
100		1-[2-(3-Methoxy-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid 2,4-dimethoxy-benzylamide	504
101		1-[2-(4-Chloro-phenoxy)-acetyl]-4-phenyl-piperidine-4-carboxylic acid 2,4-dimethoxy-benzylamide	524
102		1-(3-Cyclopentyl-propionyl)-4-phenyl-piperidine-4-carboxylic acid 2,4-dimethoxy-benzylamide	480
103		1-Butyryl-4-phenyl-piperidine-4-carboxylic acid 2,4-dimethoxy-benzylamide	426







Example	Structure	Name	M+H
104		1-(2-Fluoro-benzoyl)-4-phenyl-piperidine-4-carboxylic acid 2,4-dimethoxy-benzylamide	479
105		1-Cyclohexanecarbonyl-4-phenyl-piperidine-4-carboxylic acid 2,4-dimethoxy-benzylamide	466
106		(2-Phenyl-cyclopropyl)-[4-phenyl-4-(4-phenyl-piperazine-1-carbonyl)-piperidin-1-yl]-methanone	495
107		[1-(4-Methoxy-benzoyl)-4-phenyl-piperidin-4-yl]-(4-phenyl-piperazin-1-yl)-methanone	485
108		2-(4-Fluoro-phenyl)-1-[4-phenyl-4-(4-phenyl-piperazine-1-carbonyl)-piperidin-1-yl]-ethanone	487
109		2-(3-Methoxy-phenyl)-1-[4-phenyl-4-(4-phenyl-piperazine-1-carbonyl)-piperidin-1-yl]-ethanone	499
110		2-(4-Chloro-phenoxy)-1-[4-phenyl-4-(4-phenyl-piperazine-1-carbonyl)-piperidin-1-yl]-ethanone	519
111		3-Cyclopentyl-1-[4-phenyl-4-(4-phenyl-piperazine-1-carbonyl)-piperidin-1-yl]-propan-1-one	475







Example	Structure	Name	M+H
112		1-[4-Phenyl-4-(4-phenyl-piperazine-1-carbonyl)-piperidin-1-yl]-butan-1-one	421
113		(2-Fluoro-phenyl)-[4-phenyl-4-(4-phenyl-piperazine-1-carbonyl)-piperidin-1-yl]-methanone	473
114		(1-Cyclohexanecarbonyl-4-phenyl-piperidin-4-yl)-(4-phenyl-piperazin-1-yl)-methanone	461
115		4-Phenyl-1-(2-phenyl-cyclopropanecarbonyl)-piperidine-4-carboxylic acid 4-methoxy-benzylamide	470
116		1-(4-Methoxy-benzoyl)-4-phenyl-piperidine-4-carboxylic acid 4-methoxy-benzylamide	460
117		1-[2-(4-Fluoro-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid 4-methoxy-benzylamide	462
118		1-[2-(3-Methoxy-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid 4-methoxy-benzylamide	474
119		1-[2-(4-Chloro-phenoxy)-acetyl]-4-phenyl-piperidine-4-carboxylic acid 4-methoxy-benzylamide	494

Example	Structure	Name	M+H
120		1-(3-Cyclopentyl-propionyl)-4-phenyl-piperidine-4-carboxylic acid 4-methoxy-benzylamide	450
121		1-Butyryl-4-phenyl-piperidine-4-carboxylic acid 4-methoxy-benzylamide	396
122		1-(2-Fluoro-benzoyl)-4-phenyl-piperidine-4-carboxylic acid 4-methoxy-benzylamide	448
123		1-Cyclohexanecarbonyl-4-phenyl-piperidine-4-carboxylic acid 4-methoxy-benzylamide	436
124		4-Phenyl-1-(2-phenyl-cyclopropanecarbonyl)-piperidine-4-carboxylic acid 2-methoxy-benzylamide	470
125		1-(4-Methoxy-benzoyl)-4-phenyl-piperidine-4-carboxylic acid 2-methoxy-benzylamide	460

Example	Structure	Name	M+H
126		1-[2-(4-Fluoro-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid 2-methoxy-benzylamide	462
127		1-[2-(3-Methoxy-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid 2-methoxy-benzylamide	474
128		1-[2-(4-Chloro-phenoxy)-acetyl]-4-phenyl-piperidine-4-carboxylic acid 2-methoxy-benzylamide	494
129		1-(3-Cyclopentyl-propionyl)-4-phenyl-piperidine-4-carboxylic acid 2-methoxy-benzylamide	450
130		1-Butyryl-4-phenyl-piperidine-4-carboxylic acid 2-methoxy-benzylamide	396
131		1-(2-Fluoro-benzoyl)-4-phenyl-piperidine-4-carboxylic acid 2-methoxy-benzylamide	448






Example	Structure	Name	M+H
132		1-Cyclohexanecarbonyl-4-phenyl-piperidine-4-carboxylic acid 2-methoxy-benzylamide	436
133		1-(4-Methoxy-benzoyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide	440
134		1-[2-(4-Fluoro-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid benzylamide	430
135		1-[2-(3-Methoxy-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid benzylamide	444
136		1-[2-(4-Chloro-phenoxy)-acetyl]-4-phenyl-piperidine-4-carboxylic acid benzylamide	464
137		1-(3-Cyclopentyl-propionyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide	420
138		1-Butyryl-4-phenyl-piperidine-4-carboxylic acid benzylamide	365

Example	Structure	Name	M+H
139		1-(2-Fluoro-benzoyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide	417
140		1-Cyclohexanecarbonyl-4-phenyl-piperidine-4-carboxylic acid benzylamide	406
141		4-Phenyl-1-(2-phenyl-cyclopropanecarbonyl)-piperidine-4-carboxylic acid (3-methoxy-propyl)-amide	422
142		1-(4-Methoxy-benzoyl)-4-phenyl-piperidine-4-carboxylic acid (3-methoxy-propyl)-amide	412
143		1-[2-(4-Fluoro-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid (3-methoxy-propyl)-amide	414
144		1-[2-(3-Methoxy-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid (3-methoxy-propyl)-amide	426







Example	Structure	Name	M+H
145		1-[2-(4-Chloro-phenoxy)-acetyl]-4-phenyl-piperidine-4-carboxylic acid (3-methoxy-propyl)-amide	446
146		1-(3-Cyclopentyl-propionyl)-4-phenyl-piperidine-4-carboxylic acid (3-methoxy-propyl)-amide	402
147		1-Butyryl-4-phenyl-piperidine-4-carboxylic acid (3-methoxy-propyl)-amide	347
148		1-(2-Fluoro-benzoyl)-4-phenyl-piperidine-4-carboxylic acid (3-methoxy-propyl)-amide	399
149		1-Cyclohexanecarbonyl-4-phenyl-piperidine-4-carboxylic acid (3-methoxy-propyl)-amide	388
150		4-Phenyl-1-(2-phenyl-cyclopropanecarbonyl)-piperidine-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide	484







Example	Structure	Name	M+H
151		1-(4-Methoxy-benzoyl)-4-phenyl-piperidine-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide	474
152		1-[2-(4-Fluoro-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide	476
153		1-[2-(3-Methoxy-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide	488
154		1-[2-(4-Chloro-phenoxy)-acetyl]-4-phenyl-piperidine-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide	508
155		1-(3-Cyclopentyl-propionyl)-4-phenyl-piperidine-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide	464
156		1-Butyryl-4-phenyl-piperidine-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide	410







Example	Structure	Name	M+H
157		1-(2-Fluoro-benzoyl)-4-phenyl-piperidine-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide	462
158		1-Cyclohexanecarbonyl-4-phenyl-piperidine-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide	450
159		4-Phenyl-1-(2-phenyl-cyclopropanecarbonyl)-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	468
160		1-(4-Methoxy-benzoyl)-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	458
161		1-[2-(4-Fluoro-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	460







Example	Structure	Name	M+H
162		1-[2-(3-Methoxy-phenyl)-acetyl]-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	472
163		1-[2-(4-Chloro-phenoxy)-acetyl]-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	492
164		1-(3-Cyclopentyl-propionyl)-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	448
165		1-Butyryl-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	394
166		1-(2-Fluoro-benzoyl)-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	446








Example	Structure	Name	M+H
167		1-Cyclohexanecarbonyl-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	434
168		1-But-2-enoyl-4-phenyl-piperidine-4-carboxylic acid 2,4-dimethoxy-benzylamide	424
169		1-(3-Fluoro-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid 2,4-dimethoxy-benzylamide	514
170		1-[4-Phenyl-4-(4-phenyl-piperazine-1-carbonyl)-piperidin-1-yl]-but-2-en-1-one	419
171		[1-(3-Fluoro-benzenesulfonyl)-4-phenyl-piperidin-4-yl]-(4-phenyl-piperazin-1-yl)-methanone	509
172		1-But-2-enoyl-4-phenyl-piperidine-4-carboxylic acid 4-methoxy-benzylamide	394







Example	Structure	Name	M+H
173		1-(3-Fluoro-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid 4-methoxy-benzylamide	484
174		1-But-2-enoyl-4-phenyl-piperidine-4-carboxylic acid 2-methoxy-benzylamide	394
175		1-(3-Fluoro-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid 2-methoxy-benzylamide	484
176		1-But-2-enoyl-4-phenyl-piperidine-4-carboxylic acid benzylamide	363
177		1-But-2-enoyl-4-phenyl-piperidine-4-carboxylic acid (3-methoxy-propyl)-amide	345
178		1-(3-Fluoro-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid (3-methoxy-propyl)-amide	436







Example	Structure	Name	M+H
179		1-But-2-enoyl-4-phenyl-piperidine-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide	408
180		1-(3-Fluoro-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid [2-(4-methoxy-phenyl)-ethyl]-amide	498
181		1-But-2-enoyl-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	392
182		1-(3-Fluoro-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	482
183		4-Benzylcarbamoyl-4-phenyl-piperidine-1-carboxylic acid benzyl ester	430
184		4-Phenyl-piperidine-1,4-dicarboxylic acid 4-benzylamide 1-[(1-phenyl-ethyl)-amide]	443






Example	Structure	Name	M+H
185		1-(4-Ethyl-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide	464
186		4-Phenyl-1-(thiophene-2-sulfonyl)-piperidine-4-carboxylic acid benzylamide	442
187		1-(3-Cyano-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide	461
188		1-(2-Fluoro-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide	454
189		1-(4-Fluoro-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide	454
190		1-(4-Methoxy-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide	466






Example	Structure	Name	M+H
191		4-Phenyl-1-(toluene-3-sulfonyl)-piperidine-4-carboxylic acid benzylamide	450
192		1-(2-Phenoxy-acetyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide	430
193		1-(2-Phenoxy-acetyl)-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	458
194		4-(3,4-Difluoro-benzylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	466
195		1-(2-Phenoxy-acetyl)-4-phenyl-piperidine-4-carboxylic acid 3,4-difluoro-benzylamide	466
196		1-(4-Fluoro-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid 3,4-difluoro-benzylamide	490







Example	Structure	Name	M+H
197		4-Phenyl-piperidine-1,4-dicarboxylic acid 1-benzylamide 4-(3,4-difluoro-benzylamide)	465
198		4-Phenyl-1-(3-phenyl-acryloyl)-piperidine-4-carboxylic acid 3,4-difluoro-benzylamide	462
199		4-Phenyl-1-phenylacetyl-piperidine-4-carboxylic acid 3,4-difluoro-benzylamide	450
200		1-Benzoyl-4-phenyl-piperidine-4-carboxylic acid 3,4-difluoro-benzylamide	435
201		4-Phenyl-1-propionyl-piperidine-4-carboxylic acid 3,4-difluoro-benzylamide	387
202		1-(2-Benzyloxy-acetyl)-4-phenyl-piperidine-4-carboxylic acid 3,4-difluoro-benzylamide	480
203		4-(1-Benzyl-pyrrolidin-3-ylcarbonyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	499







Example	Structure	Name	M+H
204		4-(4-Methanesulfonyl-benzylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	508
205		4-(4-Fluoro-benzylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	448
206		4-[2-(3-Chloro-phenyl)-ethylcarbamoyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	478
207		4-Phenyl-4-[2-(3-trifluoromethyl-phenyl)-ethylcarbamoyl]-piperidine-1-carboxylic acid benzyl ester	512
208		4-[(Naphthalen-1-ylmethyl)-carbamoyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	480
209		4-Phenyl-4-(4-trifluoromethyl-benzylcarbamoyl)-piperidine-1-carboxylic acid benzyl ester	498

Example	Structure	Name	M+H
210		4-[(3-Methyl-benzo[b]thiophen-2-ylmethyl)-carbamoyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	500
211		4-(1-Benzyl-piperidin-4-ylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	513
212		4-[2-(1-Methyl-pyrrolidin-2-yl)-ethylcarbamoyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	451
213		4-(Cyclopropylmethyl-carbamoyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	394
214		4-Phenyl-4-(2-pyridin-2-yl-ethylcarbamoyl)-piperidine-1-carboxylic acid benzyl ester	445
215		4-(Indan-1-ylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	456

Example	Structure	Name	M+H
216		4-(2-Morpholin-4-yl-ethylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	453
217		1-[3-(2-Chloro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid 4-fluoro-benzylamide	480
218		1-[3-(2-Chloro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	490
219		4-Phenyl-1-[3-(4-trifluoromethyl-phenyl)-propionyl]-piperidine-4-carboxylic acid 4-fluoro-benzylamide	514
220		4-Phenyl-1-(3-phenyl-propynoyl)-piperidine-4-carboxylic acid 4-fluoro-benzylamide	458






Example	Structure	Name	M+H
221		4-Phenyl-1-[3-(4-trifluoromethyl-phenyl)-propionyl]-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	524
222		1-[3-(3,4-Difluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	492
223		4-Phenyl-1-(3-phenyl-propynoyl)-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	452
224		1-(4-Methoxy-benzenesulfonyl)-4-phenyl-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	494
225		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (biphenyl-2-ylmethyl)-amide	522






Example	Structure	Name	M+H
226		1-[3-(4-Fluoro-phenyl)-propionyl]- 4-phenyl-piperidine-4-carboxylic acid 4-trifluoromethyl-benzylamide	514
227		1-[3-(4-Fluoro-phenyl)-propionyl]- 4-phenyl-piperidine-4-carboxylic acid 4-fluoro-benzylamide	464
228		1-[3-(4-Fluoro-phenyl)-propionyl]- 4-phenyl-piperidine-4-carboxylic acid 4-chloro-benzylamide	480
229		1-[3-(4-Fluoro-phenyl)-propionyl]- 4-phenyl-piperidine-4-carboxylic acid 3-trifluoromethyl-benzylamide	514
230		1-[3-(4-Fluoro-phenyl)-propionyl]- 4-phenyl-piperidine-4-carboxylic acid 3,5-bis-trifluoromethyl- benzylamide	582
231		1-[3-(4-Fluoro-phenyl)-propionyl]- 4-phenyl-piperidine-4-carboxylic acid (thiophen-2-ylmethyl)-amide	452






Example	Structure	Name	M+H
232		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid benzylamide	446
233		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid 3-methyl-benzylamide	460
234		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid 4-methyl-benzylamide	460
235		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid 2-chloro-benzylamide	480
236		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid indan-1-ylamide	472
237		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (1,2,3,4-tetrahydronaphthalen-1-yl)-amide	486






Example	Structure	Name	M+H
238		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-amide	494
239		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(3-chloro-phenyl)-ethyl]-amide	494
240		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(3-trifluoromethyl-phenyl)-ethyl]-amide	528
241		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	488
242		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (2-thiophen-2-yl-ethyl)-amide	466


Example	Structure	Name	M+H
243		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(1H-indol-3-yl)-ethyl]-amide	499
244		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (4-phenyl-butyl)-amide	488
245		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(3-fluoro-phenyl)-ethyl]-amide	478
246		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(2-fluoro-phenyl)-ethyl]-amide	478
247		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (2-phenoxy-ethyl)-amide	476
248		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid cyclohexylmethyl-amide	452


Example	Structure	Name	M+H
249		3-(4-Fluoro-phenyl)-1-{4-[2-(4-fluoro-phenyl)-piperidine-1-carbonyl]-4-phenyl-piperidin-1-yl}-propan-1-one	518
250		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (5-chloro-benzooxazol-2-yl)-amide	507
251		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (2-methyl-5-phenyl-2H-pyrazol-3-yl)-amide	512
252		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (4-phenyl-thiazol-2-yl)-amide	515
253		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (1H-benzoimidazol-2-ylmethyl)-amide	486

Example	Structure	Name	M+H
254		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid methyl-pyridin-2-ylmethyl-amide	461
255		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid methyl-pyridin-3-ylmethyl-amide	461
256		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(2-methoxy-phenyl)-ethyl]-amide	490
257		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(4-bromo-phenyl)-ethyl]-amide	538
258		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (2-p-tolyl-ethyl)-amide	474


Example	Structure	Name	M+H
259		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(2,5-dimethoxy-phenyl)-ethyl]-amide	520
260		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (2-benzo[1,3]dioxol-5-yl-ethyl)-amide	504
261		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(3,4-dichloro-phenyl)-ethyl]-amide	528
262		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(2,4-dimethyl-phenyl)-ethyl]-amide	488
263		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid [2-(3,4-dimethyl-phenyl)-ethyl]-amide	488

Example	Structure	Name	M+H
264		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (2-o-tolyl-ethyl)-amide	474
265		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (2-m-tolyl-ethyl)-amide	474
266		1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (3-methyl-benzo[b]thiophen-2-ylmethyl)-amide	516
267		4-Phenyl-4-[(pyridin-2-ylmethyl)-carbamoyl]-piperidine-1-carboxylic acid benzyl ester	431
268		4-Phenyl-4-(pyridin-2-ylcarbamoyl)-piperidine-1-carboxylic acid benzyl ester	416

Example	Structure	Name	M+H
269		4-(2-Methoxy-pyridin-3-ylcarbamoyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	447

EXAMPLE 270

5

Synthesis



10 **Compound 1:** Compound 1 were prepared using methodology described in Example 85.



**Compound 2:** A solution of compound 1 (1.26 g; 4.03 mmol) in anhydrous acetonitrile (25 mL) was treated with diphenyl *N*-cyanocarbonimidate (1.0g; 4.2 mmol) and heated at 85°C for 1.5 h. The acetonitrile was removed by evaporation and the crude residue was purified by column chromatography on silica gel using a 7:3  
 5 hexane:ethyl acetate to 1:1 hexane:ethyl acetate gradient as the eluent to give 0.51 g of compound 2 as a white solid. LCMS  $m/z = 338$  (M+H)<sup>+</sup>


**Title Compound:** A solution of compound 2 (0.081 g; 0.18 mol) in isopropanol (5 mL) was treated with benzyl amine (0.04 mL; 0.37 mmol) and heated at 90°C for 15 h. The isopropanol was removed by evaporation and the crude residue  
 10 was purified by recrystallization from ethyl acetate/hexane to give 0.061 g of the title compound as white crystals. LCMS  $m/z = 471$  (M+H)<sup>+</sup>

#### EXAMPLES 271 TO 274


Examples 271 to 274 were prepared using methodology described in Example  
 15 270.

Example	Structure	Name	M+H
271			408
272			423

Example	Structure	Name	M+H
273			408
274			380

EXAMPLE 275

5                      4-Phenyl-1-sulfamoyl-piperidine-4-carboxylic acid benzylamide

Synthesis


10

**Compound 1:** Compound **1** was prepared as described in Example 85.

**Title Compound:** 4-Phenyl-1-sulfamoyl-piperidine-4-carboxylic acid benzylamide was prepared using methodology described in Example 15. <sup>1</sup>H NMR


(CDCl<sub>3</sub>, rt):  $\delta$  ppm) 2.15-2.24 (2 H, m), 2.50-2.55 (2 H, m), 3.21-3.27 (2 H, m), 3.42-3.49 (2 H, m), 4.28 (2 H, s), 4.36 (2 H, d, J = 5.9 Hz), 5.47 (1 H, bs), 7.02 (2 H, d, J = 7.5 Hz), 7.23 (2 H, d, J = 1.9 Hz), 7.30-7.41 (6 H, m). LCMS Rt 1.45 min, [M+1] 374.0.

5

EXAMPLE 276

1-Dimethylsulfamoyl-4-phenyl-piperidine-4-carboxylic acid benzylamide

10


Scheme

15

**Compound 1:** Compound **1** was prepared as described in Example 85.


**Title Compound:** 1-Dimethylsulfamoyl-4-phenyl-piperidine-4-carboxylic acid benzylamide was prepared using methodology described in Example 16. <sup>1</sup>H NMR (CDCl<sub>3</sub>, rt):  $\delta$  ppm) 2.13-2.22 (2 H, m), 2.42-2.49 (2 H, m), 2.80 (6 H, s), 3.31-3.45 (4 H, m), 4.34 (2 H, d, J = 5.7 Hz), 4.28 (2 H, s), 5.47 (1 H, bs), 7.02 (2 H, t, J = 5.0, Hz), 7.21-7.25 (2 H, m), 7.27-7.41 (6 H, m). LCMS Rt 1.59 min, [M+1] 402.0.

20

EXAMPLE 277

1-(2-Methoxy-ethylsulfamoyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide

5

Scheme





10 **Compound 1:** Compound 1 was prepared as described in Example 85.





**Compound 2:** Compound 2 was prepared as described in Example 17.


**Title Compound:** 1-(2-Methoxy-ethylsulfamoyl)-4-phenyl-piperidine-4-carboxylic acid benzylamide was prepared using methodology described in Example 17. <sup>1</sup>H NMR (CDCl<sub>3</sub>, rt): δ ppm 2.15-2.24 (2 H, m), 2.44-2.51 (2 H, m), 3.11-3.25 (2 H, m), 3.30-3.43 (6 H, m), 3.48 (2 H, t, J = 5.2 Hz), 4.34 (2 H, d, J = 5.7 Hz), 4.51 (1 H, bs), 5.46 (1 H, s), 7.01-7.25 (4 H, m), 7.29-7.41 (7 H, m). LCMS Rt 1.47 min, [M+1] 432.3.

EXAMPLES 278 TO 285

Example 278 to 285 were prepared using methodology described in Example 277.

Example	Structure	Name	M+H
278		1-(4-Fluoro-benzylsulfamoyl)-4-phenyl-piperidine-4-carboxylic acid pyridin-2-ylamide	470
279		1-(2-Methoxy-ethylsulfamoyl)-4-phenyl-piperidine-4-carboxylic acid pyridin-2-ylamide	420
280		1-(4-Fluoro-benzylsulfamoyl)-4-phenyl-piperidine-4-carboxylic acid (2-methoxy-pyridin-3-yl)-amide	500
281		1-(2-Methoxy-ethylsulfamoyl)-4-phenyl-piperidine-4-carboxylic acid (2-methoxy-pyridin-3-yl)-amide	450


Example	Structure	Name	M+H
282		N-[1-(4-Fluorobenzylsulfamoyl)-4-phenylpiperidin-4-ylmethyl]-2-methoxybenzamide	482
283		N-(1-Benzylsulfamoyl-4-phenylpiperidin-4-ylmethyl)-2-methoxybenzamide	464
284		2-Methoxy-N-(4-phenyl-1-propylsulfamoyl-piperidin-4-ylmethyl)-benzamide	416
285		N-(1-Dimethylsulfamoyl-4-phenylpiperidin-4-ylmethyl)-2-methoxybenzamide	432

EXAMPLE 286

5

4-phenyl-1-(3-phenylpropionyl)-piperidine-4-carboxylic acid  
(biphenyl-3-ylmethyl)-amide

# Synthesis



5        **Compound 1:** Compound 1 is commercially available.

10        **Compound 2:** A suspension of compound 1 (3.53 g; 9.35 mmol) in 50 mL of tetrahydrofuran was treated with triethylamine (2.9 mL; 20.8 mmol) and the reaction mixture was cooled to 0°C. Hydrocinnamoyl chloride (1.92 g; 11.4 mmol) was added as a solution in 5 mL of tetrahydrofuran. The reaction mixture was allowed to slowly

15        warm to room temperature and stirred for 18 h. The tetrahydrofuran was removed by evaporation and the residue was treated with ethyl acetate (approximately 150 mL) and 10% aqueous hydrochloric acid (approximately 100 mL). The organic layer was separated, washed with saturated aqueous sodium chloride, dried (sodium sulfate), filtered and concentrated. The crude product was purified by recrystallization from

20        ethanol to give 1.02 g of compound 2 as a white solid. LCMS  $m/z = 338$  ( $\text{M}+\text{H}$ )<sup>+</sup>


**Compound 3:** A suspension of compound 2 (0.23 g; 0.67 mmol) in 12 mL of dichloromethane was treated with triethylamine (0.14 mL; 1.0 mmol) followed by fluoro-*N,N,N'*,*N'*-tetramethylformamidinium hexafluorophosphate (0.22 g; 0.83 mmol) at room temperature. After stirring for 1 h, the dichloromethane was removed by

20        evaporation to give 0.23 g of compound 3 that was used in the next step without further purification. LCMS  $m/z = 340$  ( $\text{M}+\text{H}$ )<sup>+</sup>

**Title Compound:** A solution of compound 3 (0.13 g; 0.39 mmol) in 15 mL of dichloromethane was treated with triethylamine (0.071 mL; 0.51 mmol) followed by


2-phenylbenzyl amine (0.074 mL; 0.043 mmol) at room temperature. After stirring for 8 h, an additional 50 mL of dichloromethane and 40 mL of 10% aqueous hydrochloric acid was added. The organic layer was separated, washed with saturated aqueous sodium chloride, dried (sodium sulfate), filtered and concentrated. Column chromatography on silica gel using 1:1 hexane:ethyl acetate as the eluent gave 0.16 g of 4-phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (biphenyl-3-ylmethyl)-amide as a white solid.  $m/z = 504$  (M+H)<sup>+</sup>.

### EXAMPLE 287



4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid  
methyl-phenethyl-amide

### Synthesis



**Compound 1:** Compound 1 was prepared as described in Example 286.





**Title Compound:** Compound 1 (0.015 g; 0.05 mmol) was dissolved in 1 mL acetonitrile. Polystyrene-diisopropylethylamine (PS-DIEA) resin (0.1g) was added and the resulting suspension was treated with *N*-methylphenethylamine 0.02 g; 0.1 mmol) and shaken at room temperature. After 24 h, polystyrene-tosyl chloride, high

loading (PS-TsCl) resin (0.2 g) was added and the reaction mixture was allowed to shake an additional 24 h. The reaction mixture was filtered and concentrated to give 0.010 g of 4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid methyl-phenethyl-amide as a colorless oil. LCMS  $m/z = 456$  (M+H)<sup>+</sup>

5







### EXAMPLES 288 TO 322







Examples 288 to 322 were prepared using methodology described in Example 287.







Example	Structure	Name	M+H
288		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (biphenyl-3-ylmethyl)-amide	504
289		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 3-methyl-benzylamide	442
290		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 4-chloro-benzylamide	462
291		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (1-phenyl-ethyl)-amide	442


Example	Structure	Name	M+H
292		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (2-phenyl-propyl)-amide	456
293		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid sec-butylamide	394
294		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid indan-2-ylamide	454
295		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 2,6-dimethoxy-benzylamide	488
296		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 3-trifluoromethyl-benzylamide	496
297		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid cyclopentylamide	406


Example	Structure	Name	M+H
298		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid benzyl-methyl-amide	442
299		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (2-hydroxy-indan-1-yl)-amide	470
300		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid benzyl-(1-methyl-1H-imidazol-2-ylmethyl)-amide	522
301		3-Phenyl-1-[4-phenyl-4-(4-pyridin-2-yl-piperazine-1-carbonyl)-piperidin-1-yl]-propan-1-one	484
302		3-Phenyl-1-[4-phenyl-4-(4-pyrimidin-2-yl-piperazine-1-carbonyl)-piperidin-1-yl]-propan-1-one	485
303		1-[4-[4-(4-Chloro-phenyl)-piperazine-1-carbonyl]-4-phenyl-piperidin-1-yl]-3-phenyl-propan-1-one	517

Example	Structure	Name	M+H
304		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(1H-indol-3-yl)-ethyl]-amide	481
305		1-[4-(3,4-Dihydro-1H-isoquinoline-2-carbonyl)-4-phenyl-piperidin-1-yl]-3-phenyl-propan-1-one	454
306		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (5-phenyl-1H-pyrazol-3-yl)-amide	480
307		1-[4-[4-(2-Chloro-phenyl)-piperazine-1-carbonyl]-4-phenyl-piperidin-1-yl]-3-phenyl-propan-1-one	517
308		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 3,4-difluoro-benzylamide	464
309		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (1-benzyl-pyrrolidin-3-yl)-amide	497


Example	Structure	Name	M+H
310		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 4-methanesulfonyl-benzylamide	506
311		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 4-fluoro-benzylamide	446
312		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(3-chloro-phenyl)-ethyl]-amide	476
313		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(3-trifluoromethyl-phenyl)-ethyl]-amide	510
314		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (naphthalen-1-ylmethyl)-amide	478
315		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 4-trifluoromethyl-benzylamide	496

Example	Structure	Name	M+H
316		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (3-methyl-benzo[b]thiophen-2-ylmethyl)-amide	498
317		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (1-benzyl-piperidin-4-yl)-amide	511
318		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(1-methyl-pyrrolidin-2-yl)-ethyl]-amide	449
319		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid cyclopropylmethyl-amide	392
320		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (2-pyridin-2-yl-ethyl)-amide	443
321		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid indan-1-ylamide	454

Example	Structure	Name	M+H
322		4-Phenyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (2-morpholin-4-yl-ethyl)-amide	451

EXAMPLE 323

5 4-(Benzylcarbamoyl-methyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester

Synthesis

**Compound 1:** Compound 1 is commercially available.

**Compound 2:** Benzyl chloroformate (4.75 mL, 33.1 mmol) was added dropwise to a solution of 4-phenyl-4-piperidinecarboxylic acid p-methylbenzenesulfonate (10.0 g, 26.5 mmol) in 1M sodium hydroxide (200 mL) / dichloromethane (100 mL). After 2 hours the reaction mixture was made acidic with 1M hydrochloric acid (pH = 3), the organic layer was separated and the aqueous layer extracted with ethyl acetate (3 x 100 mL). The organic layers were collected, concentrated and crude product washed with water (3 x 50 mL) to give 8.52 g of 4-phenyl-piperidine-1, 4-dicarboxylic acid monobenzyl ester. LRMS  $m/z$  340.2 (M+H)<sup>+</sup>.


**Compound 3:** Thionyl chloride (4.29 mL, 58.9 mmol) was added to 4-Phenyl-piperidine-1,4-dicarboxylic acid monobenzyl ester (2.00 g, 5.89 mmol) and heated to reflux for 2 hours. The reaction mixture was concentrated under reduced pressure, taken up in ethyl ether (25 mL), cooled to 0 °C and diazomethane in ethyl ether was (30.0 mmole, 100 mL) added. After completion of the reaction, as monitored by thin layer chromatography, the excess diazomethane was quenched with acetic acid (5 mL). The reaction mixture was concentrated under reduced pressure and crude product purified by column chromatography to give 1.27 g of 4-(2-diazo-acetyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester.

**Compound 4:** 4-(2-Diazo-acetyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester (1.50 g, 4.12 mmol) in methanol (40 mL) was irradiated under UV( $\lambda=365$  nm) for 36 hours. The reaction mixture was concentrated under reduced pressure, crude product taken up in 3M lithium hydroxide (20 mL) / dioxane (20 mL) and heated to 60 °C for 1 hour. The reaction mixture was concentrated under reduced pressure to remove dioxane, made acidic with 6M hydrochloric acid, and extracted with ethyl acetate (6 x 50 mL). Organic layers were collected, concentrated and crude product purified by column chromatography to give 1.16 g of 4-carboxymethyl-4-phenyl-piperidine-1-carboxylic acid benzyl ester. <sup>1</sup>H NMR (CD<sub>3</sub>Cl<sub>3</sub>, 300 MHz)  $\delta$  7.32 (m, 10H), 5.10 (s, 2H), 3.76(d, 2H, J = 13.4 Hz), 3.20(t, 2H, J = 10.7 Hz ), 2.56 (s, 2H), 2.31(d, 2H, J = 13.6 Hz), 1.91(t, 2H, J = 13.0 Hz).

**Title Compound:** Benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (0.120 g, 0.271 mmol) was added to a solution of benzyl amine


(0.030 mL, 0.271 mmol), 4-carboxymethyl-4-phenyl-piperidine-1-carboxylic acid benzyl ester (0.100 g, 0.247 mmol), triethylamine (0.103 mL, 0.741 mmol) in tetrahydrofuran (5 mL). After 1 hour the mixture was diluted with ethyl ether (20 mL), washed with saturated aqueous sodium bicarbonate (20 mL) followed by water (2 x 10 mL). Organic layer collected, concentrated under reduced pressure and crude product purified by column chromatography to give 0.099 g of 4-(benzylcarbamoyl-methyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester. LRMS  $m/z$  443.2 (M+H)<sup>+</sup>.

10

EXAMPLE 324

N-[2-(2-Fluoro-phenyl)-ethyl]-2-{1-[3-(4-fluoro-phenyl)-propionyl]-4-phenyl-piperidin-4-yl}-acetamide

15

Synthesis

**Compound 1:** Compound 1 were prepared using methodology described in Example 323.

**Compound 2:** Ammonium formate (1.00 g) was added to a solution of 4-{[2-(2-fluoro-phenyl)-ethylcarbamoyl]-methyl}-4-phenyl-piperidine-1-carboxylic acid  
5 benzyl ester (0.520 g, 1.10 mmol) in methanol (50 mL) containing 10% palladium/carbon (0.500 g) and stirred for 4 h. The reaction mixture was filtered through celite and concentrated under reduced pressure. The crude product was taken up in 1M sodium hydroxide (100 mL) and extracted with ethyl acetate (3 x 50 mL). The organic layers were collected and concentrated under reduced pressure to give  
10 0.328 g (87%) of N-[2-(2-fluoro-phenyl)-ethyl]-2-(4-phenyl-piperidin-4-yl)-acetamide. LRMS  $m/z$  341.1 (M+H)<sup>+</sup>.

**Title Compound:** Benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (0.311 g, 0.704 mmol) was added to a solution of N-[2-(2-fluoro-phenyl)-ethyl]-2-(4-phenyl-piperidin-4-yl)-acetamide (0.200 g, 0.587 mmol), 3-  
15 (4-Fluoro-phenyl)-propionic acid (0.118 g, 0.704 mmol) and triethylamine (0.245 mL, 1.76 mmol) in tetrahydrofuran (5 mL). After 1 hour the mixture was diluted with ethyl ether (20 mL) and washed with saturated aqueous sodium bicarbonate (20 mL) followed by water (2 x 10 mL). The organic layer was collected, concentrated under reduced pressure and crude product purified by column chromatography on silica gel  
20 to give 0.098 g (34 %) of N-[2-(2-fluoro-phenyl)-ethyl]-2-{1-[3-(4-fluoro-phenyl)-propionyl]-4-phenyl-piperidin-4-yl}-acetamide. LRMS  $m/z$  491.1 (M+H)<sup>+</sup>.

EXAMPLES 325 TO 380

Examples 325 to 380 were synthesized using methodology described in Example 324.






Example	Structure	Name	M+H
325		4-Phenyl-4-([(thiophen-2-ylmethyl)-carbamoyl]-methyl)-piperidine-1-carboxylic acid benzyl ester	450
326		4-Phenyl-4-[(1-phenyl-ethylcarbamoyl)-methyl]-piperidine-1-carboxylic acid benzyl ester	458
327		4-[(2-Methoxy-benzylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	474
328		4-[(3-Methoxy-benzylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	474
329		4-[(4-Methoxy-benzylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	474
330		4-[(2,3-Dimethoxy-benzylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	504

Example	Structure	Name	M+H
331		4-[(2,4-Dimethoxy-benzylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	504
332		4-[(3-Methyl-benzylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	458
333		4-[(4-Methyl-benzylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	458
334		4-[(4-Fluoro-benzylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	462
335		4-[(2-Chloro-benzylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	478
336		4-[(4-Chloro-benzylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	478






Example	Structure	Name	M+H
337		4-Phenyl-4-[(3-trifluoromethyl-benzylcarbamoyl)-methyl]-piperidine-1-carboxylic acid benzyl ester	512
338		4-Phenyl-4-[(4-trifluoromethyl-benzylcarbamoyl)-methyl]-piperidine-1-carboxylic acid benzyl ester	512
339		4-(Phenethylcarbamoyl-methyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	458
340		4-{[2-(2-Fluoro-phenyl)-ethylcarbamoyl]-methyl}-4-phenyl-piperidine-1-carboxylic acid benzyl ester	476
341		4-{[2-(3-Fluoro-phenyl)-ethylcarbamoyl]-methyl}-4-phenyl-piperidine-1-carboxylic acid benzyl ester	476
342		4-{[2-(4-Fluoro-phenyl)-ethylcarbamoyl]-methyl}-4-phenyl-piperidine-1-carboxylic acid benzyl ester	476






Example	Structure	Name	M+H
343		4-Phenyl-4-([2-(3-trifluoromethyl-phenyl)-ethylcarbamoyl]-methyl)-piperidine-1-carboxylic acid benzyl ester	526
344		4-([2-(4-Ethyl-phenyl)-ethylcarbamoyl]-methyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	486
345		4-([2-(2,5-Dimethoxy-phenyl)-ethylcarbamoyl]-methyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	518
346		4-Phenyl-4-([(pyridin-3-ylmethyl)-carbamoyl]-methyl)-piperidine-1-carboxylic acid benzyl ester	445
347		4-Phenyl-4-([(pyridin-4-ylmethyl)-carbamoyl]-methyl)-piperidine-1-carboxylic acid benzyl ester	445
348		4-Phenyl-4-[(2-pyridin-4-yl-ethylcarbamoyl)-methyl]-piperidine-1-carboxylic acid benzyl ester	459
349		2-[1-[3-(4-Fluoro-phenyl)-propionyl]-4-phenyl-piperidin-4-yl]-N-(1-phenyl-ethyl)-acetamide	474

Example	Structure	Name	M+H
350		N-(1-Phenyl-ethyl)-2-[4-phenyl-1-(3-phenyl-propionyl)-piperidin-4-yl]-acetamide	456
351		2-{1-[2-(4-Fluoro-phenyl)-acetyl]-4-phenyl-piperidin-4-yl}-N-(1-phenyl-ethyl)-acetamide	460
352		2-[1-(4-Fluoro-benzoyl)-4-phenyl-piperidin-4-yl]-N-(1-phenyl-ethyl)-acetamide	446
353		2-[1-(2,3-Difluoro-benzoyl)-4-phenyl-piperidin-4-yl]-N-(1-phenyl-ethyl)-acetamide	464
354		N-(1-Phenyl-ethyl)-2-[4-phenyl-1-(2,4,5-trifluoro-benzoyl)-piperidin-4-yl]-acetamide	482


Example	Structure	Name	M+H
355		N-[2-(2-Fluoro-phenyl)-ethyl]-2-[4-phenyl-1-(3-phenyl-propionyl)-piperidin-4-yl]-acetamide	474
356		2-{1-[2-(4-Fluoro-phenyl)-acetyl]-4-phenyl-piperidin-4-yl}-N-[2-(2-fluoro-phenyl)-ethyl]-acetamide	478
357		2-[1-(4-Fluoro-benzoyl)-4-phenyl-piperidin-4-yl]-N-[2-(2-fluoro-phenyl)-ethyl]-acetamide	464
358		2-[1-(2,3-Difluoro-benzoyl)-4-phenyl-piperidin-4-yl]-N-[2-(2-fluoro-phenyl)-ethyl]-acetamide	482
359		N-[2-(2-Fluoro-phenyl)-ethyl]-2-[4-phenyl-1-(2,4,5-trifluoro-benzoyl)-piperidin-4-yl]-acetamide	500

Example	Structure	Name	M+H
360		4-[(2-Methoxy-phenylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	460
361		4-Phenyl-4-[(pyridin-2-ylmethyl)-carbamoyl]-methyl-piperidine-1-carboxylic acid benzyl ester	445
362		4-Phenyl-4-[(1H-pyrazol-3-ylcarbamoyl)-methyl]-piperidine-1-carboxylic acid benzyl ester	419
363		4-(Isoxazol-3-ylcarbamoylmethyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	420
364		4-[(3-Methyl-isoxazol-5-ylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	435
365		4-[(5-Methyl-isoxazol-3-ylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	435


Example	Structure	Name	M+H
366		4-Phenyl-4-(thiazol-2-ylcarbamoylmethyl)-piperidine-1-carboxylic acid benzyl ester	437
367		4-Phenyl-4-([1,3,4]thiadiazol-2-ylcarbamoylmethyl)-piperidine-1-carboxylic acid benzyl ester	438
368		4-Phenyl-4-[(1H-tetrazol-5-ylcarbamoyl)-methyl]-piperidine-1-carboxylic acid benzyl ester	421
369		4-[(2-Ethyl-2H-pyrazol-3-ylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	448
370		4-[(2,5-Dimethyl-2H-pyrazol-3-ylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	448

Example	Structure	Name	M+H
371		4-(Benzothiazol-2-ylcarbamoylmethyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	487
372		4-[(3-Methyl-isothiazol-5-ylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	451
373		4-Phenyl-4-[(5-phenyl-2H-pyrazol-3-ylcarbamoyl)-methyl]-piperidine-1-carboxylic acid benzyl ester	496
374		4-Phenyl-4-[(5-phenyl-oxazol-2-ylcarbamoyl)-methyl]-piperidine-1-carboxylic acid benzyl ester	497
375		4-[(5-Chloro-benzooxazol-2-ylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	505

Example	Structure	Name	M+H
376		4-Phenyl-4-[(5-trifluoromethyl-[1,3,4]thiadiazol-2-ylcarbamoyl)-methyl]-piperidine-1-carboxylic acid benzyl ester	506
377		4-[(2-Methyl-5-phenyl-2H-pyrazol-3-ylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	510
378		4-[(5-Oxo-1-phenyl-4,5-dihydro-1H-pyrazol-3-ylcarbamoyl)-methyl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	512
379		4-Phenyl-4-[(4-phenyl-thiazol-2-ylcarbamoyl)-methyl]-piperidine-1-carboxylic acid benzyl ester	513
380		N-(2-methoxy-phenyl)-2-[4-phenyl-1-(3-phenyl-propionyl)-piperidin-4-yl]-acetamide	458

EXAMPLE 381

2-[1-(4-Fluorobenzenesulfonyl)-4-phenylpiperidin-4-yl]-  
N-(2-methoxy-phenyl)-acetamide


Synthesis

10

**Compound 1:** Compound 1 was prepared using methodology described in Example 324. LRMS  $m/z$  325 (M+H)<sup>+</sup>.


**Title compound:** 2-[1-(4-Fluorobenzenesulfonyl)-4-phenylpiperidin-4-yl]-N-(2-methoxy-phenyl)-acetamide was prepared using methodology described in Example 95. LRMS  $m/z$  484 (M+H)<sup>+</sup>.

15

EXAMPLE 382

N-(2-Methoxy-phenyl)-2-(4-phenyl-1-pyrimidin-2-yl-piperidin-4-yl)-acetamide


5


Synthesis

10      **Compound 1:** Compound 1 was prepared using methodology described in Example 324. LRMS  $m/z$  325 (M+H)<sup>+</sup>.

**Title Compound:** N-(2-Methoxy-phenyl)-2-(4-phenyl-1-pyrimidin-2-yl-piperidin-4-yl)-acetamide was prepared using methodology described in Example 521. LRMS  $m/z$  403 (M+H)<sup>+</sup>.

15

EXAMPLE 383

Synthesis



5           **Compound 1:** Compound 1 was prepared using methodology described in Example 324. LRMS  $m/z$  325 ( $\text{M}+\text{H}$ )<sup>+</sup>.

**Compound 2:** Compound 2 was prepared using methodology described in Example 25. LRMS  $m/z$  470 ( $\text{M}+\text{H}$ )<sup>+</sup>.


**Title Compound:** The title compound was prepared using methodology  
 10   described in Example 25. LRMS  $m/z$  421 ( $\text{M}+\text{H}$ )<sup>+</sup>.

EXAMPLES 384 AND 385

Examples 384 and 385 was prepared using methodology described in Example 383.


Example	Structure	Name	(M+H)
384			483
385			497

5

EXAMPLE 386

2-(1-Dimethylsulfamoyl-4-phenyl-piperidin-4-yl)-  
N-(2-methoxy-phenyl)-acetamide


10

Synthesis

5        **Compound 1:** Compound 1 was prepared using methodology described in Example 324. LRMS  $m/z$  325 ( $\text{M}+\text{H}$ )<sup>+</sup>.


**Title Compound:** 2-(1-Dimethylsulfamoyl-4-phenyl-piperidin-4-yl)-N-(2-methoxy-phenyl)-acetamide was prepared using methodology described in Example 16. LRMS  $m/z$  432 ( $\text{M}+\text{H}$ )<sup>+</sup>.

10

EXAMPLE 387

2-(1-Cyano-4-phenyl-piperidin-4-yl)-N-(2-methoxy-phenyl)-acetamide


15

Synthesis

20        **Compound 1:** Compound 1 was prepared using methodology described in Example 324. LRMS  $m/z$  325 ( $\text{M}+\text{H}$ )<sup>+</sup>.


**Title Compound:** 2-(1-Cyano-4-phenyl-piperidin-4-yl)-N-(2-methoxy-phenyl)-acetamide may prepared using methodology described in Example 521 using cyanogen bromide instead of 2-chloropyrimidine. . LRMS  $m/z$  350 (M+H)<sup>+</sup>.

5

EXAMPLE 388

2-[1-(2-Methoxy-ethylsulfamoyl)-4-phenyl-piperidin-4-yl]-  
N-(2-methoxy-phenyl)-acetamide

10

Synthesis

15


**Compound 1:** Compound 1 was prepared using methodology described in Example 324. LRMS  $m/z$  325 (M+H)<sup>+</sup>.

**Title Compound:** 2-[1-(2-Methoxy-ethylsulfamoyl)-4-phenyl-piperidin-4-yl]-N-(2-methoxy-phenyl)-acetamide was prepared using methodology described in Example 17. LRMS  $m/z$  462 (M+H)<sup>+</sup>.


20

EXAMPLE 389

Example 389 was prepared using methodology described in Example 388.


Example	Structure	Name	M+H
389		N-(2-Methoxy-phenyl)-2-(4-phenyl-1-propylsulfamoyl-piperidin-4-yl)-acetamide	447

5

EXAMPLE 390

4-benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid benzylamide

10


Synthesis

**Compound 1:** Compound **1** is commercially available.


**Compound 2:** A solution of compound **1** (0.86 g; 2.7 mmol) in tetrahydrofuran (25 mL) was treated with triethylamine (0.49 mL; 3.7 mmol) and benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (1.3 g; 2.9 mmol). After 0.5 h benzylamine (0.33 mL; 3.0 mmol) was added and the reaction mixture was heated to 50°C for 15 h. The tetrahydrofuran was removed by evaporation and the residue was portioned between ethyl acetate and 5% aqueous hydrochloric acid. The organic layer was separated, washed with saturated aqueous sodium chloride, dried (anhydrous sodium sulfate), filtered and concentrated. Column chromatography on silica gel using 1:1 hexane:ethyl acetate as the eluent gave 0.81 g of compound **2** as a colorless oil. LRMS  $m/z$  410 (M+H)<sup>+</sup>.

**Compound 3:** A solution of compound **2** (0.41 g; 1.0 mmol) in dichloromethane (15 mL) was treated with trifluoroacetic acid (2 mL) at room temperature. After 24 h additional dichloromethane (50 mL) and 1 N sodium hydroxide (40 mL) was added. The organic layer was separated, washed with saturated aqueous sodium chloride, dried (anhydrous sodium sulfate), filtered and concentrated to give compound **3** that was used in the next step without additional purification. LRMS  $m/z$  309 (M+H)<sup>+</sup>.

**Title Compound:** A solution of compound **3** (0.16 g; 0.52 mmol) in tetrahydrofuran (20 mL) was treated with triethylamine (0.09 mL; 0.65 mmol) and hydrocinnamoyl chloride (0.1 g; 0.59 mmol) at room temperature. After 16 h the tetrahydrofuran was removed by evaporation and the residue was portioned between ethyl acetate and 5% aqueous hydrochloric acid. The organic layer was separated, washed with saturated sodium chloride, dried (anhydrous sodium sulfate) and concentrated. Column chromatography on silica gel using 1:1 hexane:ethyl acetate as the eluent gave 0.15 g of 4-benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid benzylamide as a white solid. LRMS  $m/z$  442 (M+H)<sup>+</sup>.

EXAMPLE 391

4-(2-Fluoro-benzyl)-1-[3-(4-fluoro-phenyl)-propionyl]-piperidine-  
4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide

Synthesis

**Compound 1:** Compound 1 is commercially available.

**Compound 2:** Di-*tert*-butyl dicarbonate (12.66 g, 58.0 mmol) was added to a solution of piperidine-4-carboxylic acid (5.00 g, 38.7 mmol) in dioxane (100 mL) and saturated aqueous sodium bicarbonate (100 mL) and stirred at room temperature for 48 h. The reaction mixture was concentrated to 100 mL under reduced pressure and ethyl acetate (200 mL) was added. The solution was made acidic with 6M hydrochloric acid (pH = 3), the organic layer collected and concentrated under reduced pressure to give 7.22 g of piperidine-1,4-dicarboxylic acid mono-*tert*-butyl ester. LRMS  $m/z$  228.1 (M-H)<sup>-</sup>.


**Compound 3:** Lithium diisopropylamide (10.9 mmol, 2M tetrahydrofuran) was added to a solution of piperidine-1,4-dicarboxylic acid mono-*tert*-butyl ester (1.00 g, 4.36 mmol) in tetrahydrofuran (25 mL) at 0 °C. After 1.5 h 2-fluorobenzyl bromide (0.788 mL, 6.54 mmol) was added, the reaction mixture was allowed to warm to room temperature and stirred overnight. The reaction was quenched with the addition of water (10 mL) followed by 1M hydrochloric acid (10 mL). The aqueous layer was extracted with ethyl acetate (3 x 50 mL), the organic layers collected, concentrated under reduced pressure and crude product purified by column chromatography to give 0.330 g, (22 %) of 4-(2-fluoro-benzyl)-piperidine-1,4-dicarboxylic acid mono-*tert*-butyl ester. <sup>1</sup>H-NMR (CD<sub>3</sub>Cl<sub>3</sub>, 300 MHz) δ 7.11-7.00 (m, 5H), 4.00 (br., 2H), 2.92 (s, 2H), 2.90 (br., 2H), 2.05 (br., 2H), 1.46 (br., 2H), 1.44 (s, 9H).

**Compound 4:** Benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (0.650 g, 1.47 mmol) was added to a solution of 2-(4-ethyl-phenyl)-ethylamine (0.235 mL, 1.47 mmol), 4-(2-fluoro-benzyl)-piperidine-1,4-dicarboxylic acid mono-*tert*-butyl ester (0.330 g, 0.978 mmol) and triethylamine (0.409 mL, 2.93 mmol) in tetrahydrofuran (5 mL). After 1 hour the mixture was diluted with ethyl ether (20 mL) and washed with saturated aqueous sodium bicarbonate (20 mL) followed by water (2 x 10 mL). The organic layer was collected, concentrated under reduced pressure and crude product purified by column chromatography to give 0.301 g (65 %) of 4-[2-(4-ethyl-phenyl)-ethylcarbamoyl]-4-(2-fluoro-benzyl)-piperidine-1-carboxylic acid *tert*-butyl ester. LRMS  $m/z$  469.1 (M+H)<sup>+</sup>.


**Compound 5:** Trifluoroacetic acid (20 mL) was added to a solution 4-[2-(4-ethyl-phenyl)-ethylcarbamoyl]-4-(2-fluoro-benzyl)-piperidine-1-carboxylic acid tert-butyl ester (0.300 g, 0.64 mmol) in dichloromethane (50 mL). After 1 h the reaction mixture was concentrated under reduced pressure, the crude product taken up in 1M sodium hydroxide (50 mL) and extracted with ethyl acetate (3 x 50 mL). The organic layers were collected, concentrated under reduced pressure and crude product purified by column chromatography to give 0.188 g (80 %) of 4-(2-fluoro-benzyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide LRMS  $m/z$  369.1 (M+H)<sup>+</sup>.

**Title Compound:** Benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (0.266 g, 0.602 mmol) was added to a solution of 4-(2-fluoro-benzyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide (0.185 g, 0.502 mmol), 3-(4-fluoro-phenyl)-propionic acid (0.101 g, 0.602 mmol) and triethylamine (0.210 mL, 1.51 mmol) in tetrahydrofuran (5 mL). After 1 h the mixture was diluted with ethyl ether (20 mL), washed with saturated aqueous sodium bicarbonate (20 mL) followed by water (2 x 10 mL). The organic layer was collected, concentrated under reduced pressure and the crude product purified by column chromatography to give 0.105 g (40 %) of 4-(2-fluoro-benzyl)-1-[3-(4-fluoro-phenyl)-propionyl]-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide LRMS  $m/z$  519.2 (M+H)<sup>+</sup>.

20

EXAMPLE 392

4-Benzyl-1-(4-fluoro-benzenesulfonyl)-piperidine-4-carboxylic acid benzylamide

Synthesis








5 **Compound 1:** Compound 1 was prepared as described in Example 390.







**Title Compound:** 4-Benzyl-1-(4-fluoro-benzenesulfonyl)-piperidine-4-carboxylic acid benzylamide was prepared using methodology described in Example 95. LRMS  $m/z$  467 ( $\text{M}+\text{H}$ )<sup>+</sup>.

10 EXAMPLES 393 TO 520

Examples 393 to 530 were synthesized using methodology described in Example 391 and Example 392.

Example	Structure	Name	M+H
393		4-Benzyl-4-benzylcarbamoyl-piperidine-1-carboxylic acid benzyl ester	444
394		4-Benzyl-1-(2-phenoxy-acetyl)-piperidine-4-carboxylic acid benzylamide	444
395		4-Benzyl-1-(3-phenyl-acryloyl)-piperidine-4-carboxylic acid benzylamide	440

Example	Structure	Name	M+H
396		4-Benzyl-1-phenylacetyl-piperidine-4-carboxylic acid benzylamide	428
397		1-Benzoyl-4-benzyl-piperidine-4-carboxylic acid benzylamide	414
398		4-Benzyl-1-propionyl-piperidine-4-carboxylic acid benzylamide	365
399		4-Benzyl-1-(2-benzyloxy-acetyl)-piperidine-4-carboxylic acid benzylamide	458
400		4-Benzyl-4-(4-fluoro-benzylcarbamoyl)-piperidine-1-carboxylic acid tert-butyl ester	428
401		4-Benzyl-4-(3-trifluoromethyl-benzylcarbamoyl)-piperidine-1-carboxylic acid tert-butyl ester	478
402		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 3-trifluoromethyl-benzylamide	510

Example	Structure	Name	M+H
403		4-Benzyl-1-[3-(2-chloro-phenyl)-propionyl]-piperidine-4-carboxylic acid 3-trifluoromethyl-benzylamide	544
404		4-Benzyl-1-(4-fluoro-benzyl)-piperidine-4-carboxylic acid 3-trifluoromethyl-benzylamide	486
405		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 4-fluoro-benzylamide	460
406		4-Benzyl-1-[3-(2-chloro-phenyl)-propionyl]-piperidine-4-carboxylic acid 4-fluoro-benzylamide	494
407		4-Benzyl-1-[3-(4-methoxy-phenyl)-propionyl]-piperidine-4-carboxylic acid 3-trifluoromethyl-benzylamide	540
408		4-Benzyl-1-[3-(4-trifluoromethyl-phenyl)-propionyl]-piperidine-4-carboxylic acid 3-trifluoromethyl-benzylamide	578






Example	Structure	Name	M+H
409		4-Benzyl-1-[3-(3,4-difluorophenyl)-propionyl]-piperidine-4-carboxylic acid 3-trifluoromethyl-benzylamide	546
410		4-Benzyl-1-[3-(4-fluorophenyl)-acryloyl]-piperidine-4-carboxylic acid 3-trifluoromethyl-benzylamide	526
411		4-Benzyl-1-(4-fluorobenzoyl)-piperidine-4-carboxylic acid 3-trifluoromethyl-benzylamide	500
412		4-Benzyl-1-[2-(4-chlorophenoxy)-acetyl]-piperidine-4-carboxylic acid 3-trifluoromethyl-benzylamide	546
413		4-Benzyl-1-[3-(4-methoxyphenyl)-propionyl]-piperidine-4-carboxylic acid 4-fluorobenzylamide	490
414		4-Benzyl-1-[3-(4-trifluoromethylphenyl)-propionyl]-piperidine-4-carboxylic acid 4-fluorobenzylamide	528







Example	Structure	Name	M+H
415		4-Benzyl-1-[3-(3,4-difluorophenyl)-propionyl]-piperidine-4-carboxylic acid 4-fluoro-benzylamide	496
416		4-Benzyl-1-(3-phenyl-propynoyl)-piperidine-4-carboxylic acid 4-fluoro-benzylamide	456
417		4-Benzyl-1-(4-fluoro-benzoyl)-piperidine-4-carboxylic acid 4-fluoro-benzylamide	450
418		4-Benzyl-1-[2-(4-chloro-phenoxy)-acetyl]-piperidine-4-carboxylic acid 4-fluoro-benzylamide	496
419		4-Benzyl-1-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid 4-fluoro-benzylamide	498
420		4-(4-Fluoro-benzyl)-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 3-trifluoromethyl-benzylamide	528







Example	Structure	Name	M+H
421		1-(4-fluorobenzenesulfonyl)-4-(4-fluorobenzyl)-piperidine-4-carboxylic acid 3-(trifluoromethyl)benzylamide	554
422		4-(4-Fluoro-benzyl)-4-phenethylcarbamoyl-piperidine-1-carboxylic acid tert-butyl ester	442
423		4-[2-(4-Ethyl-phenyl)-ethylcarbamoyl]-4-(4-fluorobenzyl)-piperidine-1-carboxylic acid tert-butyl ester	470
424		4-(4-Fluoro-benzyl)-1-(3-phenylpropionyl)-piperidine-4-carboxylic acid phenethyl-amide	474
425		4-(4-Fluoro-benzyl)-1-(2-phenoxyacetyl)-piperidine-4-carboxylic acid phenethyl-amide	476
426		1-(4-Fluoro-benzenesulfonyl)-4-(4-fluorobenzyl)-piperidine-4-carboxylic acid phenethyl-amide	500
427		4-(4-Fluoro-benzyl)-1-(3-phenylpropionyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	502
428		4-(4-Fluoro-benzyl)-1-(2-phenoxyacetyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	504








Example	Structure	Name	M+H
429		1-(4-Fluoro-benzenesulfonyl)-4-(4-fluoro-benzyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	528
430		1-Benzenesulfonyl-4-benzyl-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	492
431		4-Benzyl-1-(toluene-4-sulfonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	506
432		4-Benzyl-1-(4-fluoro-benzenesulfonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	510
433		4-Benzyl-1-(4-methoxy-benzenesulfonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	522
434		4-Benzyl-1-(4-chloro-benzenesulfonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	526
435		4-Benzyl-1-(4-(trifluoromethoxy)-benzenesulfonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	576

Example	Structure	Name	M+H
436		4-Benzyl-1-[2-(4-fluoro-phenyl)-acetyl]-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	488
437		4-Benzyl-1-(4-chloro-benzoyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	490
438		4-Benzyl-1-(2-phenyl-cyclopropanecarbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	496
439		4-Benzyl-1-[2-(4-methoxy-phenyl)-acetyl]-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	500
440		4-Benzyl-1-[2-(4-chloro-phenyl)-acetyl]-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	504
441		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(2-methoxy-phenyl)-ethyl]-amide	486
442		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(4-bromo-phenyl)-ethyl]-amide	535








Example	Structure	Name	M+H
443		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (2-p-tolyl-ethyl)-amide	470
444		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(2,5-dimethoxy-phenyl)-ethyl]-amide	516
445		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (2-benzo[1,3]dioxol-5-yl-ethyl)-amide	500
446		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(3,4-dichloro-phenyl)-ethyl]-amide	525
447		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(2,4-dimethyl-phenyl)-ethyl]-amide	484

Example	Structure	Name	M+H
448		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(3,4-dimethyl-phenyl)-ethyl]-amide	484
449		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (2-o-tolyl-ethyl)-amide	470
450		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (2-m-tolyl-ethyl)-amide	470
451		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (3-methyl-benzo[b]thiophen-2-ylmethyl)-amide	512
452		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-amide	490
453		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	484







Example	Structure	Name	M+H
454		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(3-trifluoromethyl-phenyl)-ethyl]-amide	524
455		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 2-chloro-benzylamide	476
456		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid phenethyl-amide	456
457		4-Benzyl-1-(5-methyl-3-phenyl-isoxazole-4-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	537
458		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(4-fluoro-phenyl)-ethyl]-amide	474
459		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid (3-phenyl-propyl)-amide	470

Example	Structure	Name	M+H
460		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 3-methyl-benzylamide	456
461		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid 4-chloro-benzylamide	476
462		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(2-fluoro-phenyl)-ethyl]-amide	474
463		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(3-fluoro-phenyl)-ethyl]-amide	474
464		4-Benzyl-1-(2-methoxy-acetyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	424
465		1-Benzoyl-4-benzyl-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	456
466		4-Benzyl-1-phenylacetyl-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	470

Example	Structure	Name	M+H
467		4-Benzyl-1-(4-methyl-benzoyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	470
468		4-Benzyl-1-(4-fluoro-benzoyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	474
469		4-Benzyl-1-(2-phenoxy-acetyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	486
470		4-Benzyl-1-cyclohexanecarbonyl-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	462
471		4-Benzyl-1-(isoxazole-5-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	447
472		4-Benzyl-1-(2,4,5-trifluoro-benzoyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	510
473		4-Benzyl-1-but-2-enoyl-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	420

Example	Structure	Name	M+H
474		4-Benzyl-1-pentanoyl-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	436
475		4-Benzyl-1-(3-methyl-butyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	436
476		1-Acetyl-4-benzyl-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	394
477		4-Benzyl-1-(pyridine-4-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	457
478		4-Benzyl-1-(pyridine-2-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	457
479		4-Benzyl-1-(pyridine-3-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	457
480		4-Benzyl-1-(2-chloro-pyridine-3-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	491


Example	Structure	Name	M+H
481		4-Benzyl-1-(3-piperidin-1-yl-propionyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	491
482		4-Benzyl-1-(1H-indole-2-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	495
483		4-Benzyl-1-(3-phenyl-propynoyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	480
484		4-Benzyl-1-(3-pyridin-3-yl-propionyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	485
485		4-Benzyl-1-[2-(1-methyl-1H-imidazol-4-yl)-acetyl]-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	474
486		4-Benzyl-1-(6-methyl-pyridine-3-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	471


Example	Structure	Name	M+H
487		4-Benzyl-1-(2-dimethylaminoacetyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	437
488		4-Benzyl-1-(1H-indazole-3-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	496
489		4-Benzyl-1-(pyrazine-2-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	458
490		4-Benzyl-1-(morpholine-4-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	465
491		4-Benzyl-1-(2-oxo-imidazolidine-4-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	464
492		4-Benzyl-1-(1H-pyrazole-4-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	446

Example	Structure	Name	M+H
493		4-Benzyl-1-([1,2,3]thiadiazole-4-carbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	464
494		4-Benzyl-1-[3-(4-chloro-phenyl)-propionyl]-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	518
495		1-(1H-Benzoimidazole-5-carbonyl)-4-benzyl-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	496
496		4-Benzyl-1-(1-cyano-cyclopropanecarbonyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	445
497		4-Benzyl-1-(3-phenyl-propionyl)-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	584
498		4-Benzyl-1-(4-fluoro-benzoyl)-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	574


Example	Structure	Name	M+H
499		4-Benzyl-1-phenylacetyl-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	570
500		4-Benzyl-1-(2-phenoxy-acetyl)-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	586
501		4-Benzyl-1-[2-(4-chloro-phenoxy)-acetyl]-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	621
502		1-Acetyl-4-benzyl-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	494
503		4-Benzyl-1-cyclohexanecarbonyl-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	562

Example	Structure	Name	M+H
504		4-Benzyl-1-(3-methyl-butyl)-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	536
505		4-Benzyl-1-(isoxazole-5-carbonyl)-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	547
506		4-Benzyl-1-(2-methoxy-acetyl)-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	524
507		1-Benzenesulfonyl-4-benzyl-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	592
508		4-Benzyl-1-(4-fluoro-benzenesulfonyl)-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	610
509		1-[3-(4-Fluoro-phenyl)-propionyl]-4-(4-methoxy-benzyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	532

Example	Structure	Name	M+H
510		4-(3-Chloro-benzyl)-1-[3-(4-fluoro-phenyl)-propionyl]-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	536

EXAMPLE 521

5 4-Benzyl-1-pyrimidin-2-yl-piperidine-4-carboxylic acid  
[2-(4-ethyl-phenyl)-ethyl]-amide

Synthesis

10


**Compound 1:** Compound **1** was prepared using methodology described in Example 390. LRMS  $m/z$  352 (M+H)<sup>+</sup>.

**Title Compound:** A solution of compound **1** (0.05 g; 0.14 mmol) in anhydrous acetonitrile (1 mL) was treated with 2-chloropyrimidine (0.024 g; 0.21 mmol) and diisopropyl ethylamine (0.036 mL; 0.21 mmol) and heated at 90°C for 1 h. The acetonitrile was removed by evaporation and the crude residue was purified by

preparative HPLC to give 4-benzyl-1-pyrimidin-2-yl-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide as a white solid. LRMS  $m/z$  430 (M+H)<sup>+</sup>.

### EXAMPLE 522


5



4-Benzyl-1-dimethylsulfamoyl-piperidine-4-carboxylic acid  
[2-(4-ethyl-phenyl)-ethyl]-amide

10

### Synthesis




**Compound 1:** Compound 1 was prepared using methodology described in  
15 Example 390. LRMS  $m/z$  352 (M+H)<sup>+</sup>.


**Title Compound:** 4-Benzyl-1-dimethylsulfamoyl-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide was prepared using methodology described in Example 16. LRMS  $m/z$  458 (M+H)<sup>+</sup>.

EXAMPLE 523

Example 523 was prepared using methodology described in Example 522.


Example	Structure	Name	M+H
523		4-Benzyl-1-dimethylsulfamoyl-piperidine-4-carboxylic acid [2-(4-chloro-phenyl)-ethyl]-(1-methyl-1H-imidazol-2-ylmethyl)-amide	559

5

EXAMPLE 524

4-Benzyl-1-(2-methoxy-ethylsulfamoyl)-piperidine-4-carboxylic acid  
[2-(4-ethyl-phenyl)-ethyl]-amide

10





**Compound 1:** Compound 1 was prepared using methodology described in Example 390. LRMS  $m/z$  352 (M+H)<sup>+</sup>.

15


**Title Compound:** 4-Benzyl-1-(2-methoxy-ethylsulfamoyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide was prepared using methodology described in Example 17. LRMS  $m/z$  488 (M+H)<sup>+</sup>.

EXAMPLES 525 TO 526

Examples 525 to 526 were prepared using methodology described in Example 524.


Example	Structure	Name	M+H
525		4-Benzyl-1-benzylsulfamoyl-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	521
526		4-Benzyl-1-(4-fluoro-benzylsulfamoyl)-piperidine-4-carboxylic acid [2-(4-ethyl-phenyl)-ethyl]-amide	539

5

EXAMPLE 527

1-[4-(isoquinolin-1-ylaminomethyl)-4-phenyl-piperidin-1-yl]-  
3-phenyl-propan-1-one

10

Synthesis

5        **Compound 1:** A suspension of 1-benzyl-4-phenyl-piperidine-4-carbonitrile (6.24 g; 20 mmol) in tetrahydrofuran was cooled to 0°C and treated with lithium aluminum hydride (3.04g; 80 mmol). The reaction mixture was allowed slowly warm to room temperature overnight. The reaction was cooled in an ice-acetone bath and quenched with water (12 mL) and 15% aqueous sodium hydroxide (3 mL). The  
10        resulting slurry was filtered through celite eluting with ethyl ether and evaporated to give 4.51 g of compound **1** as a colorless oil that was used in the next step without additional purification. LRMS  $m/z$  281 (M+H)<sup>+</sup>.


15        **Compound 2:** Pd(OAc)<sub>2</sub> (4.5 mg, 0.02 mmol), 2-(di-*t*-butylphosphino)biphenyl (11.9 mg, 0.04 mmol) and sodium *t*-butoxide (0.54 g, 5.6 mmol) were added to toluene (8 mL) and the mixture was sparged with argon. 1-Chloroisoquinoline (0.65 g, 4 mmol) and compound **1** (1.35 g, 4.8 mmol) were added and the reaction was heated at reflux under an argon atmosphere. After 48 hours a second portion of Pd(OAc)<sub>2</sub> (4.5 mg, 0.02 mmol) and -(di-*t*-butylphosphino)biphenyl (11.9 mg, 0.04 mmol) was added. The reaction was allowed to reflux another 24 hours  
20        then diluted with water (8 mL). The mixture was filtered through celite pad and

portioned between ethyl acetate and water. The organic phase was separated, dried (anhydrous magnesium sulfate), filtered and concentrated. Column chromatography on silica gel using a dichloromethane to ethyl acetate gradient as the eluent gave 0.29 g compound **2**. LRMS  $m/z$  408 (M+H)<sup>+</sup>.

- 5           **Compound 3:** A solution of compound **2** (0.250g) in methanol (20 mL) was treated with Pd(OH)<sub>2</sub> (50 mg), placed under an atmosphere of hydrogen (60 psi) and heated to 40°C until no starting material remained as judged by LCMS. The reaction mixture was filtered through celite using ethyl acetate as the eluent and evaporated to give 0.19 g compound **3** that was used in the next step without further purification.
- 10   LRMS  $m/z$  318 (M+H)<sup>+</sup>.


**Title Compound:** 1-[4-(isoquinolin-1-ylaminomethyl)-4-phenyl-piperidin-1-yl]-3-phenyl-propan-1-one was prepared using methodology described in Example 390. LRMS  $m/z$  451 (M+H)<sup>+</sup>.

15

EXAMPLE 528

[1-(4-Fluorobenzenesulfonyl)-4-phenylpiperidin-4-ylmethyl]-  
isoquinolin-1-yl-amine


20

Synthesis

**Compound 1:** Compound 1 was prepared as described in Example 527.


**Title Compound:** [1-(4-Fluoro-benzenesulfonyl)-4-phenyl-piperidin-4-ylmethyl]-isoquinolin-1-yl-amine was prepared using methodology described in Example 95. . LRMS  $m/z$  477 (M+H)<sup>+</sup>.

### EXAMPLE 529



Isoquinolin-1-yl-(4-phenyl-1-pyrimidin-2-yl-piperidin-4-ylmethyl)-amine

### Synthesis




15


**Compound 1:** Compound 1 was prepared as described in Example 527.

**Title Compound:** Isoquinolin-1-yl-(4-phenyl-1-pyrimidin-2-yl-piperidin-4-ylmethyl)-amine was prepared using methodology described in Example 521. LRMS  $m/z$  396 (M+H)<sup>+</sup>.

20

EXAMPLE 530

5


Synthesis

**Compound 1:** Compound **1** was prepared as described in Example 527.

10 **Compound 2:** Compound **2** was prepared using methodology described in Example 25. LRMS  $m/z$  463 (M+H)<sup>+</sup>.


**Title Compound:** The title compound was prepared using methodology described in Example 25. LRMS  $m/z$  413 (M+H)<sup>+</sup>.

15

EXAMPLE 531

4-(Isoquinolin-1-ylaminomethyl)-4-phenyl-piperidine-  
1-sulfonic acid dimethylamide


20

Synthesis

5 **Compound 1:** Compound 1 was prepared as described in Example 527.


**Title Compound:** 4-(Isoquinolin-1-ylaminomethyl)-4-phenyl-piperidine-1-sulfonic acid dimethylamide was prepared using methodology described in Example 16. LRMS  $m/z$  426 (M+H)<sup>+</sup>.

10

EXAMPLE 532

4-(Isoquinolin-1-ylaminomethyl)-4-phenyl-piperidine-1-sulfonic acid benzylamide

15


Synthesis

**Compound 1:** Compound 1 was prepared as described in Example 527.


**Title Compound:** 4-(Isoquinolin-1-ylaminomethyl)-4-phenyl-piperidine-1-sulfonic acid benzylamide was prepared using methodology described in Example 17.

5 LRMS  $m/z$  488 (M+H)<sup>+</sup>.

EXAMPLE 533



10 1-[4-phenyl-1-(3-phenyl-propionyl)-piperidin-4-ylmethyl]-  
1,3-dihydro-benzoimidazol-2-one

Synthesis

5        **Compound 1:** Compound **1** was prepared as described in Example 527.

10        **Compound 2:** A mixture of compound **1** (0.60 g, 2.14 mmol), 2-fluoronitrobenzene (0.25 mL, 2.35 mmol) and potassium carbonate (excess) in *N,N*-dimethylformamide, was stirred overnight at 70°C. The reaction was diluted with water (4 mL) then extracted with ethyl ether (4x15 mL). The combined extracts were  
 10        evaporated and the residue was purified by column chromatography on silica gel using a dichloromethane to 1:1 dichloromethane/ethyl acetate gradient. Product fractions were combined and evaporated to give 0.75 g (87%) compound **2**. LRMS  $m/z$  403 (M+H)<sup>+</sup>.


15        **Compound 3:** Compound **2** (0.75 g, 1.87 mmol) was dissolved in methanol and a catalytic amount of 10% palladium/carbon was added. The mixture was stirred under  $H_2$  (balloon pressure) until the color disappeared. The reaction was filtered through a celite pad and evaporated to give 0.58 g (84%) of compound **3**. LRMS  $m/z$  373 (M+H)<sup>+</sup>.

**Compound 4:** A solution of compound **3** (76.3 mg, 0.206 mmol) in dichloromethane (4 mL) was treated with triethylamine (63  $\mu$ L, 0.45 mmol) and a solution of diphosgene (13.7  $\mu$ L, 0.113 mmol) in dichloromethane (2 mL). After stirring overnight, the reaction was quenched with saturated aqueous sodium bicarbonate and separated. The organic phase was dried (anhydrous magnesium sulfate), filtered and concentrated. Column chromatography on silica gel using dichloromethane as the eluent gave 0.077g of compound **4**. LRMS  $m/z$  399 (M+H)<sup>+</sup>.

**Compound 5:** Compound **4** (63.1 mg, 0.159 mmol) was dissolved in methanol and Pd(OH)<sub>2</sub>/C (wet) was added. The mixture was hydrogenated at 60 psi and 50°C overnight. The reaction was purified directly by column chromatography on silica gel using 59:40:1 methanol:chloroform:ammonium hydroxide as the eluent to give 10 mg (20%) of compound **5**. LRMS  $m/z$  508 (M+H)<sup>+</sup>.


**Title Compound:** 1-[4-phenyl-1-(3-phenyl-propionyl)-piperidin-4-ylmethyl]-1,3-dihydro-benzoimidazol-2-one was prepared using methodology described in Example 390. LRMS  $m/z$  441 (M+H)<sup>+</sup>.

#### EXAMPLE 534



1-[1-(4-Fluoro-benzenesulfonyl)-4-phenyl-piperidin-4-ylmethyl]-  
2-methyl-1H-benzimidazole

# Synthesis




**Compound 1:** Compound 1 was prepared as described in Example 533.


**Compound 2:** A solution of compound 1 (0.2508 g, 0.676 mmol) in acetic acid (2 mL) was treated with EEDQ (0.170 g, 0.678 mmol) and heated to 120°C. After 12 h, the reaction mixture was evaporated to dryness and the residue was portioned between ethyl acetate (4 mL) and saturated aqueous sodium bicarbonate (3 mL). The aqueous layers was separated and washed with dichloromethane (2 x 4 mL). The combined organic layers were combined, dried (anhydrous magnesium sulfate), filtered and concentrated. Column chromatography on silica gel using an ethyl acetate to 10% methanol:ethyl acetate gradient as the eluent gave 0.21 g of compound 2. LRMS  $m/z$  397 ( $\text{M}+\text{H}$ )<sup>+</sup>

**Compound 3:** Compound 3 was prepared using methodology described in Example 533. LRMS  $m/z$  306 ( $\text{M}+\text{H}$ )<sup>+</sup>

**Title Compound:** 1-[1-(4-Fluoro-benzenesulfonyl)-4-phenyl-piperidin-4-ylmethyl]-2-methyl-1H-benzimidazole was prepared using methodology described in Example 95. LRMS  $m/z$  464 ( $\text{M}+\text{H}$ )<sup>+</sup>


EXAMPLE 535


1-[4-(2-Methyl-benzoimidazol-1-ylmethyl)-4-phenyl-piperidin-1-yl]-  
3-phenyl-propan-1-one

Synthesis

**Compound 1:** Compound **1** was prepared as described in Example 534.

**Title Compound:** 1-[4-(2-Methyl-benzoimidazol-1-ylmethyl)-4-phenyl-piperidin-1-yl]-3-phenyl-propan-1-one was prepared using methodology described in Example 94. LRMS  $m/z$  439 (M+H)<sup>+</sup>.

EXAMPLE 536

Synthesis

5

**Compound 1:** Compound 1 was prepared as described in Example 534.

**Compound 2:** Compound 2 was prepared using methodology described in Example 25. LRMS  $m/z$  451 ( $M+H$ )<sup>+</sup>.

**Title Compound:** The title compound was prepared using methodology described in Example 25. LRMS  $m/z$  402 ( $M+H$ )<sup>+</sup>.


10

EXAMPLE 537

Example 537 was prepared using methodology described in Example 536.


Example	Structure	Name	M+H
537			464

15

EXAMPLE 538

N-(4-Benzyl-piperidin-4-ylmethyl)-2-methoxy-benzamide

5

Synthesis

10 **Compound 1:** Compound **1** is commercially available.


**Compound 2:** To compound **1** (1.0 g, 9.08 mmol) in dichloromethane (30 mL) at 0 ° C was added triethylamine (1.5 mL, 10.7 mmol) followed by benzoylchloride (1.1 mL, 9.08 mmol). The reaction mixture was stirred at room temperature for 3 h, diluted with ethyl ether (100 mL), washed with saturated 1 N HCl, 1 N NaOH, water and dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by silica gel chromatography using Hexanes/EtOAc (2/1) as eluent to give a white solid compound **2**, 1.41 g (73% yield).  
 15 LCMS Rt 1.06 min, [M+1] 215.2.

**Compound 3:** To a solution of diisopropylamine (0.35 mL, 2.52 mmol) in dry tetrahydrofuran (2.5 mL) at 0 °C was added 1.58 mL (2.52 mmol) of 1.6 M n-BuLi in hexanes. After 30 minutes, the reaction mixture was cooled to -78 °C. Compound 2 (360 mg, 1.68 mmol) in dry tetrahydrofuran (2.5 mL) was added dropwise. After 1 h, BnBr (0.22 mL, 1.85 mmol) was added. After stirring from -78 °C to room temperature over 2 h the reaction mixture was quenched with water and diluted with Et<sub>2</sub>O (30 mL). The organic phase was washed with 1 N HCl, water and brine. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic phase was dried over Anhydrous sodium sulfate, filtered and concentrated. The residue was purified by silica gel chromatography using Hexanes/EtOAc (1/4) as eluent to give a white solid compound 3, 0.472 g (92% yield). NMR <sup>1</sup>H (CDCl<sub>3</sub>) δ (ppm) 1.40-2.05 (4 H, m), 2.89 (2 H, s), 3.00-3.42 (2 H, m), 3.70-3.90 (1 H, bs), 4.70-4.90 (1 H, bs), 7.26-7.43 (10 H, m).


**Compound 4:** To compound 3 in dry tetrahydrofuran (5 mL) was added 231 mg (mmol) of LAH then stirred at 65 °C for 20 h. The solution was cooled to 0 °C and quenched with water (0.23 mL), 15% NaOH (0.69 mL) and water (0.23 mL). The quenched reaction mixture was stirred at ambient temperature for 0.5 h. The solid was filtered and washed with Et<sub>2</sub>O. The organic solution was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give a colorless oil compound 4 sufficiently pure to be taken on to the next step. LCMS [M+1] 295.2.

**Compound 5:** To compound 4 (1.52 mmol) in dichloromethane (5 mL) at 0 °C was added 0.43 mL (3.06 mmol) of Et<sub>3</sub>N followed by 0.25 mL (1.68 mmol) of *o*-anisoyl chloride. After stirring 2 hours from 0 °C to room temperature, the reaction mixture was diluted with Et<sub>2</sub>O (30 mL), washed with 1N NaOH, water, sat. sodium chloride, dried over anh. sodium sulfate and concentrated. The residue was purified by silica gel chromatography using EtOAc as eluent to give a light yellow syrup compound 5, 0.551 g (85% yield). NMR <sup>1</sup>H (CDCl<sub>3</sub>) δ (ppm) 1.45-2.0 (4 H, m), 2.02-2.70 (4 H, m), 2.69 (2 H, s), 3.40 (2 H, d, J= 5.9 Hz), 3.55 (2 H, s), 4.00 (2 H, s), 7.00 (1 H, d, J= 8.2 Hz), 7.08-7.40 (12 H, m), 7.45 (1 H, dt, J= 1.7, 11.1 Hz), 8.05 (1 H, bs), 8.23 (1 H, dd, J=1.8, 7.8 Hz). LCMS Rt 1.72 min, [M+1] 429.2.

**Title Compound:** N-(4-Benzyl-piperidin-4-ylmethyl)-2-methoxy-benzamide was prepared using methodology described in Example 1. LCMS [M+1] 339.1.

EXAMPLE 539


5                    4-Benzyl-4-[(2-methoxy-benzoylamino)-methyl]-piperidine-  
1-carboxylic acid ethyl ester

Synthesis

10


**Compound 1:** Compound **1** was prepared as described in Example 538.

**Title Compound:** 4-Benzyl-4-[(2-methoxy-benzoylamino)-methyl]-  
piperidine-1-carboxylic acid ethyl ester was prepared using methodology described in  
15    Example 2. NMR  $^1\text{H}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm) 1.24 (3 H, t,  $J=7.1$  Hz), 1.47-1.51- (4 H, m),  
2.70 (2 H, s), 3.41-3.50 (4 H, m), 3.65-3.72 (2 H, m), 4.01 (3 H, s), 4.11 (2 H, q,  $J=$   
7.1 Hz), 7.02 (1 H, d,  $J=8.2$  Hz), 7.09-7.33 (6 H, m), 8.06 (1 H, bs), 7.48 (1 H, td,  $J=$   
1.7, 7.5 Hz), 8.22 (1 H, dd,  $J=2.2, 7.8$  Hz). LCMS Rt 1.72 min,  $[\text{M}+1]$  411.1.

EXAMPLE 540


N-[4-Benzyl-1-(3-phenyl-propionyl)-piperidin-4-ylmethyl]-2-methoxy-benzamide

5


Synthesis

10 **Compound 1:** Compound 1 was prepared as described in Example 538.

**Title Compound:** N-[4-Benzyl-1-(3-phenyl-propionyl)-piperidin-4-ylmethyl]-2-methoxy-benzamide was prepared using methodology described in Example 94. NMR  $^1\text{H}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm) 1.26-1.38 (2 H, m), 1.43-1.50 (2 H, m), 2.59-2.64 (2 H, m), 2.65 (2 H, s), 2.93-2.98 (2 H, m), 3.23-3.62 (5 H, m), 3.95-4.05 (1 H, m), 4.01 (3 H, s), 7.02 (1 H, d,  $J=8.2$  Hz), 7.09-7.34 (11 H, m), 7.45-7.55 (1 H, m), 8.07 (1 H, t,  $J=5.8$  Hz), 8.21 (1 H, dd,  $J=1.8, 7.8$  Hz).  
 15 LCMS Rt 1.77 min,  $[\text{M}+1]$  471.1.

EXAMPLE 541

N-[4-Benzyl-1-(4-fluoro-benzylsulfamoyl)-piperidin-4-ylmethyl]-  
2-methoxy-benzamide


Synthesis

**Compound 1:** Compound **1** was prepared as described in Example 538.


**Title Compound:** N-[4-Benzyl-1-(4-fluoro-benzylsulfamoyl)-piperidin-4-ylmethyl]-2-methoxy-benzamide was prepared using methodology described in Example 17. NMR  $^1\text{H}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm) 1.50-1.70 (4 H, m), 2.66 (2 H, s), 3.20-3.50 (6 H, m), 4.02 (3 H, s), 4.19 (2 H, d,  $J=5.0$  Hz), 4.58 (1 H, bs), 6.99-7.33 (10 H, m), 7.48 (1 H, td,  $J=1.7, 8.0$  Hz), 8.08 (1 H, bt,  $J=5.7$  Hz), 8.20 (1 H, dd,  $J=1.8, 7.8$  Hz).

EXAMPLE 542

Example 542 was prepared using methodology described in Example 541.


Example	Structure	Name	(M+1)+
542		N-[4-Benzyl-1-(2-methoxyethylsulfamoyl)-piperidin-4-ylmethyl]-2-methoxy-benzamide	476.2

5

EXAMPLE 543

N-(4-Benzyl-1-dimethylsulfamoyl-piperidin-4-ylmethyl)-2-methoxy-benzamide


10

Synthesis

**Compound 1:** Compound 1 was prepared as described in Example 538.

15

**Title Compound:** N-(4-Benzyl-1-dimethylsulfamoyl-piperidin-4-ylmethyl)-2-methoxy-benzamide was prepared using methodology described in Example 16.  
LRMS  $m/z$  446 (M+H)<sup>+</sup>.


EXAMPLES 544 AND 545

4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-  
1-carboxylic acid tert-butyl ester

and

4-[1-(4-Chloro-benzyl)-1H-tetrazol-5-yl]-4-phenyl-piperidine-  
1-carboxylic acid tert-butyl ester

10

Synthesis


**Compound 1:** Compound **1** was prepared using methodology described in  
Example 74. LRMS  $m/z$  287 ( $M+H$ )<sup>+</sup>.

**Compound 2:** Compound **1** (286 mg, 1.0 mmol), sodium azide (325 mg, 5.0 mmol) and ammonium chloride (268 mg, 5.0 mmol) in DMF (1.5 mL) was heated at 145 °C for 24 hours. After cooling to room temperature the reaction mixture was diluted with water (15 mL) and extracted with ethyl acetate (3 x 10 mL). The combined organic fractions were washed with water, saturated sodium chloride, dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by

silica gel chromatography using  $\text{CHCl}_3/\text{MeOH}$  (95:5) as eluent to give a white solid compound **2**, 250 mg (76% yield). NMR  $^1\text{H}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm) 1.42 (9 H, s), 2.05-2.20 (2 H, m), 2.60-3.30 (4 H, m), 3.90 (2 H, bd,  $J=13.6$  Hz), 7.15-7.30 (5 H, m).


**Title Compounds:** Compound **2** (240 mg, 0.73 mmol),  $\text{K}_2\text{CO}_3$  (111 mg, 0.8 mmol), 4-chlorobenzyl bromide (164 mg, 0.8 mmol) in acetonitrile (6 mL) was heated at 70 °C for 2 hours. The reaction mixture was cooled to room temperature, diluted with ethyl acetate (30 mL), washed with water, saturated sodium chloride, dried over anhydrous sodium chloride, filtered and concentrated. The residue was purified by silica gel chromatography using Hexanes/Ethyl acetate (3:1) as eluent to give a white solid 4-[2-(4-chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester, 244 mg (74% yield) NMR  $^1\text{H}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm) 1.44 (9 H, s), 2.10-2.25 (2 H, m), 2.75-2.90 (4 H, m), 3.95 (2 H, bs), 5.68 (2 H, s), 7.16-7.35 (9 H, m). LCMS Rt 1.38 min,  $[\text{M}+1]$  454.0 and white solid 4-[1-(4-chloro-benzyl)-1H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester, 38 mg (11% yield) NMR  $^1\text{H}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm) 1.43 (9 H, s), 2.00-2.55 (4 H, m), 2.75-2.90 (2 H, m), 3.00 (1 H, bs), 3.35 (1 H, bs), 3.82 (2 H, bs), 5.92 (2 H, bs), 6.71 (2 H, d,  $J=8.6$  Hz), 7.09 (2 H, d,  $J=1.4$  Hz), 7.11 (1 H, d,  $J=1.9$  Hz), 7.22-7.38 (4 H, m). LCMS Rt 1.38 min,  $[\text{M}+1]$  454.2

20

EXAMPLE 546

4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester


25

Synthesis

5        **Compound 1:** Compound 1 was prepared as described in Example 544.


**Compound 2:** Compound 1 (236 mg, 0.52 mmol) in dichloromethane (3 mL) and TFA (1 mL) was stirred at room temperature for 1 hour then concentrated. The residue was dissolved in ethyl acetate (20 mL), washed with 1N NaOH (2 x 5 mL), water, saturated aqueous sodium chloride, dried over anhydrous sodium sulfate and  
 10 concentrated to give an oil compound 2, 181 mg (98%). NMR H (CDCl<sub>3</sub>) δ (ppm) 1.24 (3 H, t, J=7.0 Hz), 2.11 (2 H, bs), 2.43 (2 H, bs), 3.05 (1 H, bs), 3.35 (1 H, bs), 3.85-3.95 (2 H, bs), 4.11 (2 H, q, J=7.0 Hz), 4.91 (2 H, s), 6.72 (2 H, d, J=8.4 Hz), 7.21 (2 H, d, J=8.4 Hz), 7.10 (2 H, d, J=6.1 Hz), 7.29-7.38 (3 H, m). [M+1] 426.0

**Title Compound:** 4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester was prepared using methodology described in  
 15 Example 2. LRMS *m/z* 455 (M+H)<sup>+</sup>.

EXAMPLE 547

20


{4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl}-  
 phenyl-methanone

Synthesis

5        **Compound 1:** Compound **1** was prepared as described in Example 544.


**Title Compound:** {4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl}-phenyl-methanone was prepared using methodology described in Example 94. LRMS  $m/z$  459 (M+H)<sup>+</sup>.

10

EXAMPLE 548

4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-  
1-carboxylic acid phenylamide

15

Synthesis

**Compound 1:** Compound 1 was prepared as described in Example 544.

**Title Compound:** 4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid phenylamide was prepared using methodology as described in Example 390. LRMS  $m/z$  474 (M+H)<sup>+</sup>.

5

### EXAMPLES 549 TO 598








Examples 549 to 598 were prepared using methodology described in Example 546, Example 547 and Example 548.







Example	Structure	Name	(M+1)+
549		4-(2-Benzyl-2H-tetrazol-5-yl)-4-phenyl-piperidine-1-carboxylic acid ethyl ester	392.2
550		4-Phenyl-4-[2-(3-trifluoromethoxy-benzyl)-2H-tetrazol-5-yl]-piperidine-1-carboxylic acid ethyl ester	476.2
551		4-[2-(3-Methyl-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	406.2
552		4-[2-(3-Bromo-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	470.1
553		4-{2-[2-(3-Methoxy-phenyl)-2-oxo-ethyl]-2H-tetrazol-5-yl}-4-phenyl-piperidine-1-carboxylic acid ethyl ester	450.2
554		4-[2-(3,4-Dichloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	460.1

Example	Structure	Name	(M+1)+
555		4-[2-(4-Fluoro-3-trifluoromethyl-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	478.2
556		4-[2-(4-Methyl-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	406.2
557		4-[2-(2-Oxo-2-phenyl-ethyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	420.2
558		4-[2-(4-tert-Butyl-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	448.2
559		4-Phenyl-4-[2-(4-trifluoromethoxy-benzyl)-2H-tetrazol-5-yl]-piperidine-1-carboxylic acid ethyl ester	476.2
560		4-(2-Naphthalen-2-ylmethyl-2H-tetrazol-5-yl)-4-phenyl-piperidine-1-carboxylic acid ethyl ester	442.2
561		4-[2-(3-Fluoro-4-trifluoromethyl-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	478.2
562		4-[2-(4-Bromo-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	470.1
563		4-[2-(3,5-Difluoro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	428.2


Example	Structure	Name	(M+1)+
564		4-[2-(2,4-Dichloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	460.1
565		4-[2-(2-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	426.1
566		4-[2-(3-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	426.1
567		4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	426.1
568		4-[2-(2-Oxo-propyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	358.2
569		4-(2-Allyl-2H-tetrazol-5-yl)-4-phenyl-piperidine-1-carboxylic acid ethyl ester	342.2
570		4-[2-(3-Methoxy-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	422.2
571		4-[2-(4-Methoxy-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	422.2
572		4-[2-(2-Methoxy-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	422.2

Example	Structure	Name	(M+1)+
573		4-[2-(3,5-Dimethoxy-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	452.2
574		4-[2-(2-Methyl-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	406.2
575		4-Phenyl-4-[2-(2-trifluoromethoxy-benzyl)-2H-tetrazol-5-yl]-piperidine-1-carboxylic acid ethyl ester	476.1
576		4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid tert-butyl ester	454.2
577		4-[2-(2-Fluoro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	410.2
578		4-[2-(3-Fluoro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	410.2
579		4-[2-(4-Fluoro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	410.2
580		4-Phenyl-4-[2-(2-trifluoromethyl-benzyl)-2H-tetrazol-5-yl]-piperidine-1-carboxylic acid ethyl ester	460.2
581		4-Phenyl-4-[2-(3-trifluoromethyl-benzyl)-2H-tetrazol-5-yl]-piperidine-1-carboxylic acid ethyl ester	460.2

Example	Structure	Name	(M+1)+
582		4-Phenyl-4-[2-(4-trifluoromethyl-benzyl)-2H-tetrazol-5-yl]-piperidine-1-carboxylic acid ethyl ester	460.2
583		{4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl}-(4-fluoro-phenyl)-methanone	476.1
584		{4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl}-p-tolyl-methanone	472.2
585		1-{4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl}-2-phenyl-ethanone	472.2
586		1-{4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl}-2-(4-fluoro-phenyl)-ethanone	490.2
587		1-{4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl}-2-p-tolyl-ethanone	486.2
588		1-{4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl}-3-phenyl-propan-1-one	486.2


Example	Structure	Name	(M+1)+
589		1-[4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl]-3-(4-fluoro-phenyl)-propan-1-one	504.2
590		1-[4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl]-3-p-tolyl-propan-1-one	500.2
591		1-[4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl]-2-phenoxy-ethanone	488.1
592		4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	488.1
593		1-[4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl]-2-methoxy-ethanone	426.2
594		4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid (4-fluoro-phenyl)-amide	491.2

Example	Structure	Name	(M+1)+
595		4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidine-1-carbothioic acid phenylamide	489.1
596		{4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl}-pyridin-4-yl-methanone	459.2
597		{4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl}-pyridin-2-yl-methanone	459.2
598		1-{4-[2-(4-Chloro-benzyl)-2H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl}-2-thiophen-2-yl-ethanone	478.1

EXAMPLE 599

5

4-[1-(4-Chloro-benzyl)-1H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester

Synthesis

5 **Compound 1:** Compound 1 was prepared as described in Example 545.


**Compound 2:** Compound 1 (31 mg, 0.068 mmol) in dichloromethane (0.75 mL) and TFA (0.25 mL) was stirred at room temperature for 1.5 hour then concentrated. The residue was dissolved in ethyl acetate (10 mL), washed with 1N NaOH (2 x 5 mL), water, saturated sodium chloride, dried over anhydrous sodium sulfate and concentrated to give an oil compound 6, 20 mg (83%). NMR H (CDCl<sub>3</sub>) δ (ppm) 1.72 (1 H, bs), 2.12-2.21 (2 H, m), 2.46 (2 H, bd, J= 13.7 Hz), 2.94 (4 H, bs), 4.92 (2 H, s), 6.71 (2 H, d, J=8.4 Hz), 7.12 (2 H, d, J= 6.4 Hz), 7.16 (2 H, d, J= 11.3 Hz), 7.21-7.36 (3 H, m).

10

**Title Compound:** 4-[1-(4-Chloro-benzyl)-1H-tetrazol-5-yl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester was prepared using methodology described in Example 2. NMR H (CDCl<sub>3</sub>) δ (ppm) 1.24 (3 H, t, J=7.0 Hz), 2.11 (2 H, bs), 2.43 (2 H, bs), 3.05 (1 H, bs), 3.35 (1 H, bs), 3.85-3.95 (2 H, bs), 4.11 (2 H, q, J=7.0 Hz), 4.91 (2 H, s), 6.72 (2 H, d, J=8.4 Hz), 7.21 (2 H, d, J=8.4 Hz), 7.10 (2 H, d, J=6.1 Hz), 7.29-7.38 (3 H, m). [M+1] 426.0

15


20

EXAMPLE 600

1-{4-[1-(4-Chloro-benzyl)-1H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl}-

3-phenyl-propan-1-one

25


Synthesis

5

**Compound 1:** Compound 1 was prepared as described in Example 599.


**Title Compound:** 1-[4-[1-(4-Chloro-benzyl)-1H-tetrazol-5-yl]-4-phenyl-piperidin-1-yl]-3-phenylpropan-1-one was prepared using methodology described in Example 390. NMR H (CDCl<sub>3</sub>) δ (ppm) 1.72-1.77 (1 H, m), 2.09-2.27 (2 H, m), 2.48-2.62 (3 H, m), 2.69-2.77 (1 H, m), 2.92-2.96 (2 H, m), 3.63 (2 H, dd, J= 3.0, 8.0 Hz), 4.31-4.35 (1 H, m), 4.81 (1 H, d, J=15.8 Hz), 4.98 (1 H, d, J=15.8 Hz), 6.71 (2 H, d, J=8.5 Hz), 6.71 (2 H, d, J=8.5 Hz), 7.06-7.38 (8 H, m). [M+1] 486.3

15

EXAMPLE 601

1-[4-(5-Phenethyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-  
3-phenylpropan-1-one

# Synthesis



**Compound 1:** Compound 1 was prepared as described in Example 74.

**Compound 2:** To hydroxylamine hydrochloride (4.25 g, 61.2 mmol) and sodium methoxide (3.30 g, 61.2 mmol) in  $\text{PrOH}$  (100 mL) was added compound **1** (5.8 g, 20.4 mmol). The reaction mixture was heated at reflux overnight, diluted with Ethyl acetate (200 mL), washed with water, sat. sodium chloride, dried over anhydrous sodium sulfate and concentrated to give a white solid compound **2**, 4.60 g (72% yield). LCMS  $R_t$  1.06 min,  $[\text{M}+1]$  320.2.

**Compound 3:** Compound **2** (31 mg, 0.10 mmol), Et<sub>3</sub>N (15 mg, 0.15 mmol) and hydrocinnamoyl chloride (19 mg, 0.11 mmol) in tetrahydrofuran (0.5 mL) was stirred overnight. The reaction mixture was diluted with ethyl acetate (5 mL), washed with saturated sodium bicarbonate, water, brine, dried over anhydrous sodium sulfate, filtered and concentrated to give a white solid compound **3**, 0.03 g (66% yield).

**Compound 4:** To compound **3** (30 mg, 0.066 mmol) in dry tetrahydrofuran (0.5 mL) was added 0.13 mL (0.132 mmol) of tetrabutyl ammonium fluoride. After stirring for two hours at room temperature the reaction mixture was diluted with Ethyl acetate (5 mL), washed with water, brine, dried over anhydrous sodium sulfate, filtered and concentrated to give a white solid compound **4** sufficiently pure to be taken on to the next step.

**Compound 5:** Compound **5** were prepared using methodology described in Example 85 using hydrochloric acid instead of trifluoroacetic acid to accomplish nitrogen deprotection. LRMS *m/z* 334 (M+H)<sup>+</sup>.









**Title Compound:** 1-[4-(5-Phenethyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-3-phenyl-propan-1-one was prepared using methodology described in Example 94. LRMS *m/z* 466 (M+H)<sup>+</sup>.

#### EXAMPLES 602 TO 626


Examples 602 to 626 were prepared as described in Example 601.

Example	Structure	Name	(M+H) <sup>+</sup>
602		1-{4-[5-(2-Methoxy-phenyl)-[1,2,4]oxadiazol-3-yl]-4-phenyl-piperidin-1-yl}-3-phenyl-propan-1-one	468.2
603		1-[4-(5-Phenethyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-2-phenyl-ethanone	452.2

Example	Structure	Name	(M+H) <sup>+</sup>
604		2-(4-Fluoro-phenyl)-1-[4-(5-phenethyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-ethanone	470.2
605		1-[4-(5-Phenethyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-2-p-tolyl-ethanone	466.2
606		3-(4-Fluoro-phenyl)-1-[4-(5-phenethyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-propan-1-one	484.2
607		1-[4-(5-Phenethyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-3-p-tolyl-propan-1-one	480.3
608		1-{4-[5-(4-Chloro-benzyl)-[1,2,4]oxadiazol-3-yl]-4-phenyl-piperidin-1-yl}-2-phenyl-ethanone	472.2
609		1-{4-[5-(4-Chloro-benzyl)-[1,2,4]oxadiazol-3-yl]-4-phenyl-piperidin-1-yl}-2-(4-fluoro-phenyl)-ethanone	490.2
610		1-{4-[5-(4-Chloro-benzyl)-[1,2,4]oxadiazol-3-yl]-4-phenyl-piperidin-1-yl}-2-p-tolyl-ethanone	486.2
611		1-{4-[5-(4-Chloro-benzyl)-[1,2,4]oxadiazol-3-yl]-4-phenyl-piperidin-1-yl}-3-phenyl-propan-1-one	486.2


Example	Structure	Name	(M+H) <sup>+</sup>
612		1-{4-[5-(4-Chloro-benzyl)-[1,2,4]oxadiazol-3-yl]-4-phenyl-piperidin-1-yl}-3-(4-fluoro-phenyl)-propan-1-one	504.2
613		1-{4-[5-(4-Chloro-benzyl)-[1,2,4]oxadiazol-3-yl]-4-phenyl-piperidin-1-yl}-3-p-tolyl-propan-1-one	500.2
614		1-{4-[5-(4-Chloro-benzyl)-[1,2,4]oxadiazol-3-yl]-4-phenyl-piperidin-1-yl}-2-phenoxy-ethanone	488.2
615		[4-(5-Benzyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-(4-fluoro-phenyl)-methanone	442.2
616		1-[4-(5-Benzyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-2-phenyl-ethanone	438.2
617		1-[4-(5-Benzyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-2-(4-fluoro-phenyl)-ethanone	456.2
618		1-[4-(5-Benzyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-2-p-tolyl-ethanone	452.2
619		1-[4-(5-Benzyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-3-phenyl-propan-1-one	452.2

Example	Structure	Name	(M+H) <sup>+</sup>
620		1-[4-(5-Benzyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-3-(4-fluoro-phenyl)-propan-1-one	470.2
621		1-[4-(5-Benzyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-3-p-tolyl-propan-1-one	466.2
622		1-[4-(5-Phenethyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-2-thiophen-2-yl-ethanone	458.2
623		4-[5-(4-Chloro-benzyl)-[1,2,4]oxadiazol-3-yl]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	488.2
624		1-[4-(5-Benzyl-[1,2,4]oxadiazol-3-yl)-4-phenyl-piperidin-1-yl]-2-thiophen-2-yl-ethanone	444.2
625		1-{4-[5-(4-Methoxy-benzyl)-[1,2,4]oxadiazol-3-yl]-4-phenyl-piperidin-1-yl}-3-phenyl-propan-1-one	482.2
626		3-(4-Fluoro-phenyl)-1-{4-[5-(4-methoxy-benzyl)-[1,2,4]oxadiazol-3-yl]-4-phenyl-piperidin-1-yl}-propan-1-one	500.2

EXAMPLE 627

4-(5-Fluoro-1-methyl-1H-benzimidazol-2-yl)-4-phenyl-piperidine-  
1-carboxylic acid benzyl ester

5

Synthesis

10

**Compound 1:** Compound **1** is commercially available.

**Compound 2:** 2,5-Difluoronitrobenzene (1.0 g, 6.29 mmol), methylamine hydrochloride (2.12 g, 31.43 mmol) and potassium carbonate (4.34 g, 31.43 mmol) in DMSO (10 mL) was stirred for 24 hours, diluted with Et<sub>2</sub>O (200 mL), washed with  
15 water (3 times), brine, dried over anhydrous sodium sulfate and concentrated to give compound **2** as an orange solid 1.1 g (100 %).

**Compound 3:** Compound 2 (1.1 g, 6.29 mmol) and 10% Pd on charcoal (300 mg) in methanol (20 mL) was stirred under hydrogen for 24 hours, filtered over celite and concentrated. The residue was purified by silica gel chromatography using Hexanes/Ethyl acetate (2:1) as eluent to give a white solid compound 3, 672 mg (74%  
5 yield). NMR  $^1\text{H}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm) 2.83 (3 H, s), 3.28 (3 H, bs), 6.40-6.60 (3 H, m).

**Compound 4:** Compound 4 was prepared as described in Example 323.







**Compounds 5 and 6:** Compound 4 (700 mg, 2.06 mmol) in thionyl chloride was heated at reflux for 2 hours then concentrated. The residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (3.5 mL) and cooled to 0 °C. Compound 3 (289 mg, 2.06 mmol) in  $\text{CH}_2\text{Cl}_2$   
10 (3.5 mL) followed by  $\text{Et}_3\text{N}$  (1.44 mL, 10.3 mmol) were added. After 1 hour the reaction mixture was diluted with Ethyl acetate (30 mL), washed with 1N NaOH, water, brine, dried over anhydrous sodium sulfate and concentrated. The residue was purified by silica gel chromatography using Hexanes/Ethyl acetate (2:1) as eluent to give a white solid compound 5 and compound 6 521 mg (55% yield) as an inseparable  
15 mixture. LCMS Rt 1.42 min and 1.52,  $[\text{M}+1]$  462.2.








**Title Compound:** The mixture of compound 5 and compound 6 (521 mg, 1.12 mmol) in glacial acetic acid (5 mL) was heated at reflux for 2 hours then concentrated. The residue was purified by silica gel chromatography using Hexanes/Ethyl acetate (2:1) as eluent to give a white solid 4-(5-fluoro-1-methyl-1H-  
20 benzoimidazol-2-yl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester, 365 mg (73% yield). NMR  $^1\text{H}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm) 2.05-2.40 (2 H, m), 2.55-2.75 (2 H, m), 3.22 (3 H, s), 3.25-3.40 (1 H, m), 3.50-3.70 (1 H, m), 5.14 (2 H, s), 7.01 (1 H, dt,  $J=2.4$ , 9.1 Hz), 7.09-7.35 (13 H, m), 7.48 (1 H, dd,  $J=2.2$ , 9.4 Hz).  
LCMS Rt 1.41 min,  $[\text{M}+1]$  442.2.







25


EXAMPLES 628 TO 646

Examples 628 to 646 were prepared as described in Example 627.

Example	Structure	Name	(M+1)+
628		4-(1H-Benzimidazol-2-ylmethyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	426.2
629		4-(1-Methyl-1H-benzimidazol-2-ylmethyl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	440.2
630		4-(1H-Benzimidazol-2-yl)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	412.2
631		1-[4-(5-Fluoro-1-methyl-1H-benzimidazol-2-yl)-4-phenyl-piperidin-1-yl]-3-phenyl-propan-1-one	442.2
632		1-[4-(5-Fluoro-1-methyl-1H-benzimidazol-2-yl)-4-phenyl-piperidin-1-yl]-2-phenoxy-ethanone	444.2
633		5-Fluoro-2-[1-(4-fluoro-benzenesulfonyl)-4-phenyl-piperidin-4-yl]-1-methyl-1H-benzimidazole	468.2


Example	Structure	Name	(M+1)+
634			485.2
635		1-[4-(5-Fluoro-1-methyl-1H-benzimidazol-2-yl)-4-phenyl-piperidin-1-yl]-2-(4-fluoro-phenoxy)-ethanone	462.2
636		1-[4-(5-Fluoro-1H-benzimidazol-2-yl)-4-phenyl-piperidin-1-yl]-3-(4-methoxy-phenyl)-propan-1-one	458.2
637		1-[4-(5-Fluoro-1H-benzimidazol-2-yl)-4-phenyl-piperidin-1-yl]-4-phenyl-butan-1-one	442.2
638		1-[4-(5-Fluoro-1H-benzimidazol-2-yl)-4-phenyl-piperidin-1-yl]-2-(3-trifluoromethyl-phenyl)-ethanone	482.2
639		5-Chloro-2-[1-(3-fluoro-benzenesulfonyl)-4-phenyl-piperidin-4-yl]-1H-benzimidazole	470.1
640		5-Chloro-2-[1-(4-fluoro-benzenesulfonyl)-4-phenyl-piperidin-4-yl]-1H-benzimidazole	470.1

Example	Structure	Name	(M+1)+
641			467.2
642			495.2
643			481.2
644		1-[4-(1-Methyl-1H-benzimidazol-2-ylmethyl)-4-phenyl-piperidin-1-yl]-3-phenyl-propan-1-one	438.3
645		2-[1-(4-Methoxy-benzenesulfonyl)-4-phenyl-piperidin-4-ylmethyl]-1-methyl-1H-benzimidazole	476.2
646		4-(5-Chloro-1H-benzimidazol-2-yl)-4-phenyl-piperidine-1-carboxylic acid ethyl ester	384.1

EXAMPLE 647

4-tert-Butoxycarbonylamino-4-phenyl-piperidine-1-carboxylic acid benzyl ester

5

Synthesis


10 **Compound 1:** Compound **1** was prepared as described in Example 323.

**Compound 2:** A solution of compound **1** (0.62 g; 1.8 mmol) in tetrahydrofuran (40 mL) was cooled to 0°C and treated with triethylamine (0.38 mL; 2.7 mmol) and diphenylphosphoryl azide (0.44 mL; 2.0 mmol). The cooling bath was removed and the reaction mixture was allowed to stir at room temperature 18 h. The tetrahydrofuran was removed by evaporation and the crude residue was purified by column chromatography on silica gel using 8:2 hexane:ethyl acetate as the eluent to give 0.67g of compound **2**. LRMS  $m/z$  365 (M+H)<sup>+</sup>.

**Title Compound:** A solution of compound **2** (0.67 g; 1.8 mmol) in chlorobenzene (15 mL) was treated with t-butanol (25 mL) and heated at 115°C for 20


- h. The chlorobenzene was removed by evaporation and the residue was subjected to column chromatography on silica gel using an 8:2 hexane:ethyl acetate to 1:1 hexane:ethyl acetate gradient as the eluent to give 0.05 g of 4-tert-butoxycarbonylamino-4-phenyl-piperidine-1-carboxylic acid benzyl ester as a colorless oil LRMS  $m/z$  411 (M+H)<sup>+</sup> and 0.2 g compound **3** as a colorless oil LRMS  $m/z$  337 (M+H)<sup>+</sup>.

### EXAMPLE 648



4-(2-Methoxy-benzoylamino)-4-phenyl-piperidine-1-carboxylic acid benzyl ester

### Synthesis




**Compound 1:** Compound **1** was prepared as described in Example 626.

**Compound 2:** A solution of compound **1** (0.05g; 0.1 mmol) in dichloromethane (2 mL) was treated with trifluoroacetic acid (1 mL) and allowed to stir at room temperature for 1 h at which time LCMS indicated complete consumption of compound **1**. Additional dichloromethane (20 mL) and 1 N sodium hydroxide (20 mL) were added. The organic layer was separated, washed with saturated aqueous sodium chloride, dried (anhydrous sodium sulfate), filtered and concentrated to give

0.03 g of compound 2 that was used in the next step without additional purification.  
LCMS  $m/z = 311$  (M+H)<sup>+</sup>


**Title Compound:** A solution of compound 2 (0.03g; 0.009 mmol) in tetrahydrofuran (2 mL) was treated with triethylamine (0.1 mL; 0.7 mmol) and o-anisoyl chloride (0.05 g; 0.3 mmol) at room temperature. After stirring 15 h the tetrahydrofuran was removed by evaporation and the residue was purified by prep LC to give 0.011 g of compound 4-(2-methoxy-benzoylamino)-4-phenyl-piperidine-1-carboxylic acid benzyl ester as a white solid. LCMS  $m/z = 461$  (M+H)<sup>+</sup>

10

EXAMPLE 649

4-[3-(2-Methoxy-benzyl)-ureido]-4-phenyl-piperidine-1-carboxylic acid benzyl ester

15

Synthesis

**Compound 1:** Compound 1 was prepared as described in Example 626.




20

**Title Compound:** A solution of compound 1 (0.05g; 0.15 mmol) in anhydrous tetrahydrofuran (2 mL) was treated with 2-methoxybenzyl amine (0.041g; 0.30 mmol) at room temperature. After stirring 2 h the tetrahydrofuran was removed by evaporation and the residue was purified by prep LC to give 0.015 g of 4-[3-(2-


methoxy-benzyl)-ureido]-4-phenyl-piperidine-1-carboxylic acid benzyl ester as a white solid. LCMS  $m/z = 475$  (M+H)<sup>+</sup>.

### EXAMPLES 650 TO 652


5 Examples 650 to 652 were prepared as described in Example 649.

Example	Structure	Name	M+H
650		4-[3-(2-Methoxy-phenyl)-ureido]-4-phenyl-piperidine-1-carboxylic acid benzyl ester	461
651		4-(3-Phenethyl-ureido)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	459
652		4-(3-Isoquinolin-1-yl-ureido)-4-phenyl-piperidine-1-carboxylic acid benzyl ester	482

### EXAMPLE 653



3-Amino-pyrazine-2-carboxylic acid (4-phenyl-piperidin-4-ylmethyl)-amide

Synthesis

5 **Compound 1:** Compound 1 is commercially available.

**Compound 2:** A solution of compound 1 (20 g; 90 mmol) in 1 N sodium hydroxide (200 mL) and tetrahydrofuran (50 mL) was treated with di-*tert*-butyl dicarbonate (19.7 g; 90.3 mmol) at room temperature. After stirring 24 h the reaction mixture was adjusted to pH = 7 by the addition of 1 N hydrochloric acid and washed with ethyl acetate (2 x 100 mL). The organic layers were combined, washed with saturated aqueous sodium chloride, dried (sodium sulfate), filtered and concentrated. The crude product was recrystallized from the minimum amount of ethanol to give 20 g compound 2 as a white solid. LRMS  $m/z$  = 287 (M+H)<sup>+</sup>


15 **Compound 3:** A solution of compound 2 (4.5 g; 15.7 mmol) in ethanol (42.5 mL) and acetic acid (7.5 mL) was treated with platinum oxide (250 mg) and hydrogenated at 60 psi for 12h. Analysis by thin layer chromatography indicated the reaction was not complete. Additional platinum oxide (124 mg) was added and the reaction mixture hydrogenated at 60 psi for another 12 h. The reaction mixture was filtered through celite and concentrated to give 4.6 g of compound 3 as a colorless oil that was used without additional purification. LRMS  $m/z$  = 291 (M+H)<sup>+</sup>

**Compound 4:** A suspension of 3-amino-pyrazine-2-carboxylic acid (2.9 g; 20.8 mmol) in tetrahydrofuran (30 mL) was treated with triethylamine (7 mL; 50 mmol) followed by benzotriazole-1-yl-oxy-tris-(dimethylamino)-phosphoniumhexafluorophosphate (8.8 g; 19.9 mmol) at room temperature. After 15

minutes of stirring the reaction mixture became homogeneous and compound **3** (4.6 g; 15.8 mmol) was added as a solution in tetrahydrofuran (20 mL). The reaction was heated at 60°C for 12 h. The solvent was removed by rotary evaporation and the crude product was purified directly by column chromatography on silica gel using 1:1 ethyl acetate:hexane as the eluent to give 2.0 g of compound **4** as a white foam. LRMS  $m/z = 412$  (M+H)<sup>+</sup>


**Title Compound:** A solution of compound **4** (1.9g; 4.6 mmol) in anhydrous dichloromethane (50 mL) was treated with trifluoroacetic acid (50 mL) at room temperature. After stirring for 19 h the reaction mixture was made basic (pH = 12) with 6 N NaOH and additional dichloromethane (100 mL) was added. The organic layer was separated, washed with saturated aqueous sodium chloride, dried (sodium sulfate), filtered and concentrated to give 1.4 g of the title compound as a white foam that was used without additional purification. LRMS  $m/z = 312$  (M+H)<sup>+</sup>

15

EXAMPLE 654

3-Amino-pyrazine-2-carboxylic acid [4-(3-fluoro-phenyl)-  
piperidin-4-ylmethyl]-amide

20

Synthesis

5 **Compound 1:** Compound 1 is commercially available.

**Compound 2:** Compound 1 (5.0 g, 28.0 mmol) was suspended in 100 mL of dichloromethane. Di-*tert*-butyl dicarbonate (6.1 g, 28.0 mmol) and diisopropylethylamine (10.7 mL, 61.6 mmol) was added and the reaction mixture was stirred at room temperature overnight. The reaction mixture was diluted with diethyl  
 10 ether (300 mL), washed with 1N hydrochloric acid (2 x 100 mL), saturated sodium bicarbonate (100 mL), water (100 mL) and saturated aqueous sodium chloride (100 mL). The organic layer was dried (sodium sulfate) and concentrated. The crude residue was purified by silica gel chromatography using 9:1 hexane:diethyl ether as eluent to give 3.46 g of compound 2 as a colorless oil.

15 **Compound 3:** Sodium hydride (1.9 g, 79.0 mmol) was suspended in dimethylformamide (30 mL) and cooled to 0°C. 3-Fluorophenylacetonitrile (2.61 g, 19.3 mmol) was slowly added followed by the addition of compound 2 (3.68 g, 15.2 mmol) as a solution in dimethylformamide (30 mL). The reaction was stirred at 0°C

for 0.5 hours and was allowed to warm to room temperature and stirred overnight. The reaction mixture was poured into ice and extracted with ethyl acetate (200 mL). The aqueous layer was extracted with ethyl acetate (2 x 200 mL). The organic layers were combined and washed with 10% lithium chloride (2 x 100 mL), dried (sodium sulfate), filtered and concentrated. The crude residue was purified by flash chromatography on silica gel using 9:1 hexane:ethyl acetate as the eluent to give 4.04 g of compound **3** as a yellow oil.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.48 (9H, s), 1.92 (2H, td,  $J = 4.3, 13.1$ ), 2.05 (2H, t,  $J = 10.2$ ), 3.19 (2H, t,  $J = 12.4$ ), 4.29 (2H, d,  $J = 13.7$ ), 7.01-7.45 (4H, m). LRMS  $m/z$  305 ( $\text{M}+\text{H}$ ) $^+$ .

**Compound 4:** Compound **3** (4.04 g, 13.27 mmol) in 50 mL of 15% acetic acid in ethanol was treated with platinum (IV) oxide (0.210 g, 0.925 mmol) and hydrogenated at 60 psi (Parr Apparatus) for 48 h. The reaction mixture was filtered through a thick pad of celite and concentrated. To the concentrate was added 200 mL of diethyl ether and the organic solution was washed with 1N sodium hydroxide (2 x 50 mL) and dried (sodium sulfate), filtered and concentrated to give 4.07 g of compound **4**. LRMS  $m/z$  309 ( $\text{M}+\text{H}$ ) $^+$ .

**Compound 5:** To compound **4** (0.562 g, 1.94 mmol) in 15 mL of tetrahydrofuran was added 3-aminopyrazine-2-carboxylic acid (0.270 g, 1.94 mmol), benzotriazole-1-yl-oxy-tris-(dimethylamino)-phosphoniumhexafluorophosphate (0.858 g, 1.94 mmol) and triethylamine (0.271 mL, 1.94 mmol). The reaction mixture was stirred for 2 hours at room temperature then diluted with diethyl ether (50 mL), washed with water (3 x 50 mL), saturated sodium chloride (50 mL) and dried over sodium sulfate. After filtration, the solvent was removed by rotary evaporation and the crude residue was purified using flash chromatography (1:1 hexane/diethyl ether) to give 0.69 g of compound **5**. LRMS  $m/z$  430 ( $\text{M}+\text{H}$ ) $^+$ .

**Title Compound:** Compound **5** (0.691 g, 1.61 mmol) was suspended in tetrahydrofuran (5 mL) and 4 N hydrochloric acid in 1,4-dioxane (5 mL) and the reaction mixture was stirred overnight at room temperature. The reaction mixture was concentrated, dissolved in water (15 mL) and the aqueous layer was washed with diethyl ether (2 x 15 mL). 6 N Sodium hydroxide was added to basify the aqueous layer that was then extracted with dichloromethane (3 x 20 mL). The organic layers were combined, dried (sodium sulfate), filtered and concentrated to give 0.47 g of the







title compound as a yellow solid which was used without additional purification.








LRMS  $m/z$  330 (M+H)<sup>+</sup>.








### EXAMPLES 655 TO 695

- 5 Examples 655-695 were synthesized using methodology described in Example 1 and Example 654.

Example	Structure	Name	M+H
655		3-Amino-pyrazine-2-carboxylic acid [1-benzyl-4-(4-fluoro-phenyl)-piperidin-4-ylmethyl]-amide	422
656		N-[1-Benzyl-4-(2-fluoro-phenyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	450
657		N-[4-(2-Fluoro-phenyl)-piperidin-4-ylmethyl]-2-methoxy-benzamide	343
658		N-[4-(2-Fluoro-phenyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	359
659		1-Benzoyl-4-(3-chloro-phenyl)-piperidine-4-carbonitrile	326

Example	Structure	Name	M+H
660		1-Benzoyl-4-(2,5-difluorophenyl)-piperidine-4-carbonitrile	327
661		C-[1-Benzyl-4-(3-chlorophenyl)-piperidin-4-yl]-methylamine	316
662		N-[1-Benzyl-4-(3-chlorophenyl)-piperidin-4-ylmethyl]-2-methoxy-benzamide	450
663		N-[1-Benzyl-4-(3-chlorophenyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	466
664		3-Amino-pyrazine-2-carboxylic acid [1-benzyl-4-(3-chlorophenyl)-piperidin-4-ylmethyl]-amide	437
665		N-[4-(3-Chloro-phenyl)-piperidin-4-ylmethyl]-2-methoxy-benzamide	360


Example	Structure	Name	M+H
666		N-[4-(3-Chloro-phenyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	376
667		3-Amino-pyrazine-2-carboxylic acid [4-(3-chloro-phenyl)-piperidin-4-ylmethyl]-amide	347
668		4-Cyano-4-(2,5-difluoro-phenyl)-piperidine-1-carboxylic acid tert-butyl ester	323
669		4-Cyano-4-(3-fluoro-phenyl)-piperidine-1-carboxylic acid tert-butyl ester	305
670		4-Aminomethyl-4-(3-fluoro-phenyl)-piperidine-1-carboxylic acid tert-butyl ester	309
671		4-(3-Fluoro-phenyl)-4-[(2-methoxy-benzoylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester	444
672		N-[4-(3-Fluoro-phenyl)-piperidin-4-ylmethyl]-2-methoxy-benzamide	343

Example	Structure	Name	M+H
673		3-Amino-pyrazine-2-carboxylic acid [1-benzyl-4-(3-methoxy-phenyl)-piperidin-4-ylmethyl]-amide	433
674		4-[[[(3-Amino-pyrazine-2-carbonyl)-amino]-methyl]-4-(3-fluoro-phenyl)-piperidine-1-carboxylic acid tert-butyl ester	430
675		3-Amino-pyrazine-2-carboxylic acid [4-(3-methoxy-phenyl)-piperidin-4-ylmethyl]-amide	342
676		3-Amino-pyrazine-2-carboxylic acid [4-(3-fluoro-phenyl)-piperidin-4-ylmethyl]-amide	330
677		4-Aminomethyl-4-(2,5-difluoro-phenyl)-piperidine-1-carboxylic acid tert-butyl ester	327
678		4-(2,5-Difluoro-phenyl)-4-[(2-methoxy-benzoylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester	462
679		4-(2,5-Difluoro-phenyl)-4-[(2-hydroxy-6-methoxy-benzoylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester	478


Example	Structure	Name	M+H
680		4-[[[3-Amino-pyrazine-2-carbonyl)-amino]-methyl]-4-(2,5-difluoro-phenyl)-piperidine-1-carboxylic acid tert-butyl ester	448
681		N-[4-(2,5-Difluoro-phenyl)-piperidin-4-ylmethyl]-2-methoxy-benzamide	361
682		3-Amino-pyrazine-2-carboxylic acid [4-(2,5-difluoro-phenyl)-piperidin-4-ylmethyl]-amide	348
683		2,3-Dihydro-benzofuran-7-carboxylic acid (1-benzyl-4-phenyl-piperidin-4-ylmethyl)-amide	428
684		N-(1-Benzyl-4-phenyl-piperidin-4-ylmethyl)-5-chloro-2-methoxy-benzamide	450
685		N-[1-Benzyl-4-(3-methoxy-phenyl)-piperidin-4-ylmethyl]-2-methoxy-benzamide	446
686		2-Methoxy-N-[4-(3-methoxy-phenyl)-piperidin-4-ylmethyl]-benzamide	355

Example	Structure	Name	M+H
687		N-(1-Benzyl-4-phenyl-piperidin-4-ylmethyl)-2-hydroxy-6-methoxy-benzamide	432
688		2-Hydroxy-6-methoxy-N-(4-phenyl-piperidin-4-ylmethyl)-benzamide	341
689		4-(3-Fluoro-phenyl)-4-[(2-hydroxy-6-methoxy-benzoylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester	460
690		N-[1-Benzyl-4-(2-fluoro-phenyl)-piperidin-4-ylmethyl]-2-methoxy-benzamide	434
691		4-(3,5-Difluoro-phenyl)-4-[(2-hydroxy-6-methoxy-benzoylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester	478
692		4-[[[3-Amino-pyrazine-2-carbonyl)-amino]-methyl]-4-(3,5-difluoro-phenyl)-piperidine-1-carboxylic acid tert-butyl ester	448
693		4-[(2-Hydroxy-6-methoxy-benzoylamino)-methyl]-4-(3-trifluoromethyl-phenyl)-piperidine-1-carboxylic acid tert-butyl ester	510

Example	Structure	Name	M+H
694		4-[[[(3-Amino-pyrazine-2-carbonyl)-amino]-methyl]-4-(3-trifluoromethyl-phenyl)-piperidine-1-carboxylic acid tert-butyl ester	481
695		4-[(2-Methoxy-benzoylamino)-methyl]-4-(3-trifluoromethyl-phenyl)-piperidine-1-carboxylic acid tert-butyl ester	494

EXAMPLE 696

5 *N*-(1-Cyclohexylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-methoxy-benzamide

Synthesis

**Compound 1:** Compound 1 was prepared as described in Example 15.

**Compound 2:** Compound 2 was prepared as described in *J. Org. Chem.* **2003**, 68, 115-119.

**Compound 3:** Compound 2 (3.34 g, 8.6 mmol) was suspended in 15 mL of acetonitrile and cooled to 0°C. Compound **1** (2.14 g, 6.6 mmol) was slowly added. The reaction was allowed to warm to room temperature overnight. The reaction mixture was concentrated and purified directly by flash chromatography on silica gel using 3:1 ethylacetate/hexane as the eluent to give 1.92 g of compound **3**. LRMS  $m/z$  469 (M+H)<sup>+</sup>.

**Compound 4:** Compound **3** (1.9 g, 4.05 mmol) was suspended in 16 mL of dichloromethane and cooled to 0°C. A solution of methyl triflate (0.505 mL, 4.46 mmol) in dichloromethane (16 mL) was slowly added. The reaction was allowed to warm to room temperature, stirred for 2 h and concentrated. Compound **4** was used in the next step without further purification. LRMS  $m/z$  483 (M+H)<sup>+</sup>.

**Title Compound:** To compound **4** (0.075 g, 0.118 mmol) and cyclohexylamine (0.016 mL, 0.142 mmol) was added 0.695 mL of acetonitrile. The reaction mixture was heated to 80°C for 4 h and concentrated. The crude material was purified using preparative reverse phase HPLC to give 0.026 g of the title compound as a white solid. LRMS  $m/z$  488 (M+H)<sup>+</sup>.

20






EXAMPLES 697 TO 843








Examples 697 to 843 were prepared using methodology described in Example 15, Example 16, Example 17 and Example 696.








Example	Structure	Name	M+H
697		4-Phenyl-1-sulfamoyl-piperidine-4-carboxylic acid benzyl-methyl-amide	390
698		1-(4-Fluorobenzylsulfamoyl)-4-phenyl-piperidine-4-carboxylic acid benzyl-methyl-amide	498
699		4-Phenyl-1-sulfamoyl-piperidine-4-carboxylic acid [2-(3-trifluoromethylphenyl)-ethyl]-amide	457
700		1-(4-Fluorobenzylsulfamoyl)-4-phenyl-piperidine-4-carboxylic acid [2-(3-trifluoromethylphenyl)-ethyl]-amide	566
701		N-[1-(Imidazole-1-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxybenzamide	457







Example	Structure	Name	M+H
702		4-Phenyl-1-sulfamoyl-piperidine-4-carboxylic acid (biphenyl-3-ylmethyl)-amide	452
703		1-(4-Fluorobenzylsulfamoyl)-4-phenyl-piperidine-4-carboxylic acid (biphenyl-3-ylmethyl)-amide	560
704		N-(4-Cyclohex-1-enyl-1-dimethylsulfamoyl)-piperidin-4-ylmethyl)-2-methoxy-benzamide	438
705		N-(4-Cyclohex-1-enyl-1-dimethylsulfamoyl)-piperidin-4-ylmethyl)-2,4-dimethoxy-benzamide	468
706		N-(4-Cyclohex-1-enyl-1-dimethylsulfamoyl)-piperidin-4-ylmethyl)-2,6-dimethoxy-benzamide	468
707		N-(4-Cyclohex-1-enyl-1-dimethylsulfamoyl)-piperidin-4-ylmethyl)-2-trifluoromethoxy-benzamide	492

Example	Structure	Name	M+H
708		N-(4-Cyclohex-1-enyl-1-dimethylsulfamoyl-piperidin-4-ylmethyl)-2-fluoro-benzamide	426
709		4-Cyclohex-1-enyl-4-[(2-trifluoromethoxy-benzenesulfonylamino)-methyl]-piperidine-1-sulfonic acid dimethylamide	528
710		4-Phenyl-4-[(2-trifluoromethoxy-benzenesulfonylamino)-methyl]-piperidine-1-sulfonic acid amide	496
711		4-Phenyl-4-[(2-trifluoromethoxy-benzenesulfonylamino)-methyl]-piperidine-1-sulfonic acid 4-fluorobenzamide	604
712		5-Chloro-2-methoxy-N-(4-phenyl-1-sulfamoyl-piperidin-4-ylmethyl)-benzamide	440






Example	Structure	Name	M+H
713		5-Chloro-N-[1-(4-fluorobenzylsulfamoyl)-4-phenylpiperidin-4-ylmethyl]-2-methoxybenzamide	548
714		4-Phenyl-1-sulfamoyl-piperidine-4-carboxylic acid methyl-[2-(3-trifluoromethyl-phenyl)-ethyl]-amide	472
715		1-(4-Fluorobenzylsulfamoyl)-4-phenylpiperidine-4-carboxylic acid methyl-[2-(3-trifluoromethyl-phenyl)-ethyl]-amide	580
716		2,3-Dihydro-benzofuran-7-carboxylic acid (4-phenyl-1-sulfamoyl-piperidin-4-ylmethyl)-amide	418
717		2,3-Dihydro-benzofuran-7-carboxylic acid [1-(4-fluorobenzylsulfamoyl)-4-phenylpiperidin-4-ylmethyl]-amide	526






Example	Structure	Name	M+H
718		N-[4-(3,6-Dihydro-2H-pyran-4-yl)-1-dimethylsulfamoyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	440
719		2-Methoxy-N-(1-sulfamoyl-4-thiophen-3-yl-piperidin-4-ylmethyl)-benzamide	412
720		N-[4-(4-Fluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	423
721		4-[(4-Methyl-pyridin-2-ylamino)-methyl]-4-phenyl-piperidine-1-sulfonic acid dimethylamide	391
722		N-[4-(3-Fluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	423
723		N-(1-Dimethylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-fluoro-benzamide	422
724		2,3-Dimethoxy-N-(4-phenyl-1-sulfamoyl-piperidin-4-ylmethyl)-benzamide	436






Example	Structure	Name	M+H
725		N-[4-(2-Fluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	423
726		2-Methoxy-N-{4-phenyl-1-[(pyridin-2-ylmethyl)-sulfamoyl]-piperidin-4-ylmethyl}-benzamide	497
727		2-Methoxy-N-[4-(3-methoxy-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-benzamide	436
728		Carbamic acid 2-{4-[(2-methoxy-benzoylamino)-methyl]-4-phenyl-piperidine-1-sulfonylamino}-ethyl ester	493
729		Ethyl-carbamic acid 2-{4-[(2-methoxy-benzoylamino)-methyl]-4-phenyl-piperidine-1-sulfonylamino}-ethyl ester	521
730		2-Methoxy-N-[4-phenyl-1-(2-pyridin-4-ylethylsulfamoyl)-piperidin-4-ylmethyl]-benzamide	511
731		N-(4-Phenyl-1-sulfamoyl-piperidin-4-ylmethyl)-2-trifluoromethoxy-benzamide	459






Example	Structure	Name	M+H
732		2,6-Dimethoxy-N-(4-phenyl-1-sulfamoyl-piperidin-4-ylmethyl)-benzamide	436
733		Cyclopropyl-carbamic acid 2-{4-[(2-methoxybenzoylamino)-methyl]-4-phenyl-piperidine-1-sulfonylamino}-ethyl ester	533
734		2-Methoxy-N-[1-(2-methylimidazole-1-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-benzamide	471
735		2-Methoxy-N-(1-methylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-benzamide	420
736		N-(1-Ethylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-methoxybenzamide	434
737		N-(1-Cyclopropylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-methoxybenzamide	446







Example	Structure	Name	M+H
738		2-Methoxy-N-{4-phenyl-1-[(tetrahydro-furan-2-ylmethyl)-sulfamoyl]-piperidin-4-ylmethyl}-benzamide	490
739		N-[1-(Isopropyl-methyl-sulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	462
740		2-Methoxy-N-{4-phenyl-1-[(pyridin-3-ylmethyl)-sulfamoyl]-piperidin-4-ylmethyl}-benzamide	497
741		2-Methoxy-N-{4-phenyl-1-[(pyridin-4-ylmethyl)-sulfamoyl]-piperidin-4-ylmethyl}-benzamide	497
742		N-[1-(2-Hydroxy-propylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	464






Example	Structure	Name	M+H
743		(2-{4-[(2-Methoxy-benzoylamino)-methyl]-4-phenyl-piperidine-1-sulfonylamino}-ethyl)-carbamic acid tert-butyl ester	549
744		N-{1-[(2-Hydroxy-ethyl)-methyl-sulfamoyl]-4-phenyl-piperidin-4-ylmethyl}-2-methoxy-benzamide	464
745		N-(1-Dimethylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-methoxy-nicotinamide	435
746		Quinoline-8-carboxylic acid (1-dimethylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-amide	455
747		N-[1-(Cyclopropylmethyl-sulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	460

Example	Structure	Name	M+H
748		N-[1-(3-Hydroxy-pyrrolidine-1-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	476
749		N-[1-(2-Dimethylamino-ethylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	477
750		N-[1-(2-Fluoro-ethylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	452
751		N-[1-(Carbamoylmethyl-sulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	463
752		1-{4-[(2-Methoxy-benzoylamino)-methyl]-4-phenyl-piperidine-1-sulfonyl}-pyrrolidine-2-carboxylic acid amide	503

Example	Structure	Name	M+H
753		N-(1-Isopropylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-methoxy-benzamide	448
754		N-[1-(2-Aminoethylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	449
755		2-Methoxy-N-{4-phenyl-1-[2-(pyrimidin-2-ylamino)ethylsulfamoyl]-piperidin-4-ylmethyl}-benzamide	527
756		2-Methoxy-N-{4-phenyl-1-[3-(pyrimidin-2-ylamino)propylsulfamoyl]-piperidin-4-ylmethyl}-benzamide	541
757		2-Methoxy-N-[4-phenyl-1-(pyridin-4-ylsulfamoyl)-piperidin-4-ylmethyl]-benzamide	483







Example	Structure	Name	M+H
758		2-Methoxy-N-[4-phenyl-1-(pyridin-3-ylsulfamoyl)-piperidin-4-ylmethyl]-benzamide	483
759		2-Methoxy-N-[4-phenyl-1-(pyridin-2-ylsulfamoyl)-piperidin-4-ylmethyl]-benzamide	483
760		N-[1-(4-Hydroxy-piperidine-1-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	490
761		N-[1-(3,4-Difluoro-phenylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	518
762		N-[1-(2,4-Difluoro-phenylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	518






Example	Structure	Name	M+H
763		2-Methoxy-N-(4-phenyl-1-phenylsulfamoyl-piperidin-4-ylmethyl)-benzamide	482
764		N-[1-(2-Hydroxypropylsulfamoyl)-4-phenylpiperidin-4-ylmethyl]-2-methoxybenzamide	464
765		N-[1-(2-Hydroxy-1-methylethylsulfamoyl)-4-phenylpiperidin-4-ylmethyl]-2-methoxybenzamide	464
766		N-[1-(1-Hydroxymethylpropylsulfamoyl)-4-phenylpiperidin-4-ylmethyl]-2-methoxybenzamide	478
767		N-[1-(2-Hydroxypropylsulfamoyl)-4-phenylpiperidin-4-ylmethyl]-2-methoxybenzamide	464
768		N-[1-(1-Hydroxymethylpropylsulfamoyl)-4-phenylpiperidin-4-ylmethyl]-2-methoxybenzamide	478

Example	Structure	Name	M+H
769		N-[1-(2-Hydroxy-1-methyl-ethylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	464
770		2-Methoxy-N-[1-(2-phenoxy-ethylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-benzamide	526
771		{4-[(2-Methoxy-benzoylamino)-methyl]-4-phenyl-piperidine-1-sulfonylamino}-acetic acid methyl ester	478
772		2-{4-[(2-Methoxy-benzoylamino)-methyl]-4-phenyl-piperidine-1-sulfonylamino}-propionic acid methyl ester	492
773		2-{4-[(2-Methoxy-benzoylamino)-methyl]-4-phenyl-piperidine-1-sulfonylamino}-3-phenyl-propionic acid methyl ester	568







Example	Structure	Name	M+H
774		N-(1-Dimethylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-hydroxy-6-methoxy-benzamide	450
775		N-(1-Dimethylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-fluoro-6-methoxy-benzamide	452
776		2-Difluoromethoxy-N-(1-dimethylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-benzamide	470
777		3-Amino-pyrazine-2-carboxylic acid (1-dimethylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-amide	421
778		N-[1-(4-Fluoro-phenylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	500
779		2-Methoxy-N-[1-(methoxy-methyl-sulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-benzamide	450

Example	Structure	Name	M+H
780		N-(1-Hydroxysulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-methoxy-benzamide	422
781		N-[1-(2-Fluorophenyl)sulfamoyl]-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	500
782		N-[1-(3-Fluorophenyl)sulfamoyl]-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	500
783		N-[1-(1,1-Dioxo-tetrahydro-1lambda*6*-thiophen-3-yl)sulfamoyl]-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	524
784		N-[1-(4-Hydroxymethyl-piperidine-1-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	504

Example	Structure	Name	M+H
785		N-{1-[(2-Fluoro-phenyl)-methyl-sulfamoyl]-4-phenyl-piperidin-4-ylmethyl}-2-methoxy-benzamide	514
786		N-{1-[(3-Fluoro-phenyl)-methyl-sulfamoyl]-4-phenyl-piperidin-4-ylmethyl}-2-methoxy-benzamide	514
787		N-[1-(Hydroxy-methyl-sulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	436
788		2-Methoxy-N-(1-methylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-benzamide	419
789		N-(1-tert-Butylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-methoxy-benzamide	462
790		N-[1-(4,4-Dimethyl-oxazolidine-3-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	490

Example	Structure	Name	M+H
791		N-[1-(2,6-Dimethyl-morpholine-4-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	504
792		N-[1-(4,4-Dimethyl-4,5-dihydro-imidazole-1-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	487
793		N-[1-(2-Hydroxy-1,1-dimethyl-ethylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	478
794		N-[1-(4-Hydroxy-benzylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	512
795		N-[1-(3-Hydroxy-phenylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	498

Example	Structure	Name	M+H
796		N-[1-(2-Hydroxy-cyclohexylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	504
797		2-Methoxy-N-[1-(2-methoxymethyl-pyrrolidine-1-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-benzamide	504
798		N-[1-(1-Hydroxymethyl-2-methyl-propylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	492
799		N-[1-(1-Hydroxymethyl-2-methyl-propylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	492
800		N-[1-(2-Cyclohexyl-1-hydroxymethyl-ethylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	546
801		N-[1-(2-Hydroxy-indan-1-ylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	538

Example	Structure	Name	M+H
802		N-[1-(2-Hydroxy-indan-1-ylsulfamoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	538
803		N-[1-(3-Hydroxy-pyrrolidine-1-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	476
804		1-{4-[(2-Methoxy-benzoylamino)-methyl]-4-phenyl-piperidine-1-sulfonyl}-pyrrolidine-2-carboxylic acid methyl ester	518
805		2-{4-[(2-Methoxy-benzoylamino)-methyl]-4-phenyl-piperidine-1-sulfonylamino}-3-methyl-butyric acid methyl ester	520
806		2-Hydroxy-6-methoxy-N-{1-[(2-methoxy-ethyl)-methyl-sulfamoyl]-4-phenyl-piperidin-4-ylmethyl}-benzamide	494
807		2-Hydroxy-6-methoxy-N-(4-phenyl-1-sulfamoyl-piperidin-4-ylmethyl)-benzamide	422

Example	Structure	Name	M+H
808		N-[4-(3-Fluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	439
809		N-[4-(3-Fluoro-phenyl)-1-(2-methyl-imidazole-1-sulfonyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	505
810		N-[4-(3-Fluoro-phenyl)-1-(2-hydroxy-ethylsulfamoyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	484
811		N-[1-(Cyclopropylmethyl)-sulfamoyl]-4-(3-fluoro-phenyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	494
812		N-[4-(3-Fluoro-phenyl)-1-(4-fluoro-phenylsulfamoyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	534
813		N-[4-(3-Fluoro-phenyl)-1-(4-hydroxy-benzylsulfamoyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	546
814		N-[4-(3-Fluoro-phenyl)-1-(2-hydroxy-1-methyl-ethylsulfamoyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	498
815		N-[4-(3-Fluoro-phenyl)-1-(1-hydroxymethyl-propylsulfamoyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	512


Example	Structure	Name	M+H
816		N-[4-(3-Fluoro-phenyl)-1-(2-hydroxy-propylsulfamoyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	498
817		N-[4-(3-Fluoro-phenyl)-1-(3-hydroxy-pyrrolidine-1-sulfonyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	510
818		2-Hydroxy-6-methoxy-N-[1-(2-methyl-imidazole-1-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-benzamide	487
819		Trifluoro-methanesulfonate 3-[(4-(3-fluoro-phenyl)-4-[(2-hydroxy-6-methoxy-benzoylamino)-methyl]-piperidine-1-sulfonyl)-1,2-dimethyl-3H-imidazol-1-ium];	669
820		N-[4-(3-Fluoro-phenyl)-1-(2-phenoxy-ethylsulfamoyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	560
821		N-{4-(3-Fluoro-phenyl)-1-[(2-hydroxy-ethyl)-methyl-sulfamoyl]-piperidin-4-ylmethyl}-2-hydroxy-6-methoxy-benzamide	498
822		N-{4-(3-Fluoro-phenyl)-1-[(tetrahydro-furan-2-ylmethyl)-sulfamoyl]-piperidin-4-ylmethyl}-2-hydroxy-6-methoxy-benzamide	524

Example	Structure	Name	M+H
823		N-(1-Dimethylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-hydroxy-benzamide	420
824		N-[1-Dimethylsulfamoyl-4-(3-trifluoromethyl-phenyl)-piperidin-4-ylmethyl]-2-hydroxy-benzamide	488
825		N-[1-Dimethylsulfamoyl-4-(3-trifluoromethyl-phenyl)-piperidin-4-ylmethyl]-2-methoxy-benzamide	502
826		N-[4-(2-Fluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	439
827		N-[1-Dimethylsulfamoyl-4-(2-fluoro-phenyl)-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	468
828		N-[4-(3-Chloro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	440
829		N-[4-(3-Chloro-phenyl)-1-dimethylsulfamoyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	468


Example	Structure	Name	M+H
830		N-[4-(3,5-Difluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	441
831		N-[4-(3,5-Difluoro-phenyl)-1-dimethylsulfamoyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	470
832		N-[4-(3,5-Difluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-2-hydroxy-6-methoxy-benzamide	457
833		3-Amino-pyrazine-2-carboxylic acid [4-(3,5-difluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-amide	428
834		3-Amino-pyrazine-2-carboxylic acid (4-phenyl-1-sulfamoyl-piperidin-4-ylmethyl)-amide	392
835		2-Hydroxy-6-methoxy-N-[1-sulfamoyl-4-(3-trifluoromethyl-phenyl)-piperidin-4-ylmethyl]-benzamide	489
836		3-Amino-pyrazine-2-carboxylic acid [1-sulfamoyl-4-(3-trifluoromethyl-phenyl)-piperidin-4-ylmethyl]-amide	460

Example	Structure	Name	M+H
837		3-Amino-pyrazine-2-carboxylic acid [4-(3-chloro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-amide	427
838		3-Amino-pyrazine-2-carboxylic acid [4-(3-fluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-amide	410
839		N-[4-(2,5-Difluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	441
840		3-Amino-pyrazine-2-carboxylic acid [4-(2,5-difluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-amide	428
841		2-Methoxy-N-[1-sulfamoyl-4-(3-trifluoromethyl-phenyl)-piperidin-4-ylmethyl]-benzamide	472
842		N-[4-(2,3-Difluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	440

Example	Structure	Name	M+H
843		3-Amino-pyrazine-2-carboxylic acid [4-(2,3-difluoro-phenyl)-1-sulfamoyl-piperidin-4-ylmethyl]-amide	427

EXAMPLE 844

- 5      3-Amino-pyrazine-2-carboxylic acid [1-(amino-methanesulfonylimino-methyl)-4-phenyl-piperidin-4-ylmethyl]-amide

Synthesis

**Compound 1:** Compound 1 was prepared as described in Example 653.

- 15      **Compound 2:** A solution of compound 1 (0.15 g; 0.47 mmol) in anhydrous acetonitrile (8 mL) at room temperature was treated with *N*-diphenoxymethylmethanesulfonamide (for preparation see US patent 4871765) (0.17 g; 0.5 mmol). The reaction mixture was heated to 85°C for 2 h. The solvent was removed by rotary evaporation and the crude residue was purified directly by column chromatography on silica gel using 9:1 ethyl acetate:hexane as the eluent to give 0.2 g of compound 2 as a white solid. LRMS  $m/z$  = 510 (M+H)<sup>+</sup>


**Title Compound:** Compound 2 (0.060 g; 0.12 mmol) was treated with 7 N ammonia in methanol (2 mL) and heated to 85°C for 15 minutes in a sealed tube. The solvent was removed by rotary evaporation and the crude residue was purified directly by column chromatography on silica gel using 10:1 chloroform:methanol as the eluent to give 0.024 g of the title compound as a white solid. LRMS  $m/z$  = 433 (M+H)<sup>+</sup>

### EXAMPLES 845 TO 852


Examples 845 to 852 were prepared using methodology described in Example 844.

Example	Structure	Name	M+H
845		N-[1-(Methanesulfonylimino-phenoxy-methyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	523
846		N-[1-(Amino-methanesulfonylimino-methyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	446
847		N-[1-(Cyclopropylamino-methanesulfonylimino-methyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	486
848		N-{1-[(Cyclopropylmethyl-amino)-methanesulfonylimino-methyl]-4-phenyl-piperidin-4-ylmethyl}-2-methoxy-benzamide	500

Example	Structure	Name	M+H
849		N-(1-{[Methanesulfonylimino-[(pyridin-2-ylmethyl)-amino]-methyl]-4-phenyl-piperidin-4-ylmethyl}-2-methoxy-benzamide	537
850		N-{1-[Methanesulfonylimino-(2-pyridin-4-yl-ethylamino)-methyl]-4-phenyl-piperidin-4-ylmethyl}-2-methoxy-benzamide	551
851		N-{1-[(2-Hydroxy-ethylamino)-methanesulfonylimino-methyl]-4-phenyl-piperidin-4-ylmethyl}-2-methoxy-benzamide	490
852		N-{1-[(3-Imidazol-1-yl-propylamino)-methanesulfonylimino-methyl]-4-phenyl-piperidin-4-ylmethyl}-2-methoxy-benzamide	554

EXAMPLE 853

5

Synthesis

**Compound 1:** Compound 1 was prepared as described in Example 653.



**Compound 2:** A solution of compound 1 (0.211 mg; 0.68 mmol) in anhydrous acetonitrile (5 mL) was treated with diphenyl *N*-cyanocarbonimide (0.180 mg; 0.76 mmol) and heated to 85°C for 1 h. The solvent was removed by rotary evaporation and the crude residue was purified directly by column chromatography on silica gel using 7:3 ethyl acetate:hexane as the eluent to give 0.273 g of compound 2 as a white foam. LRMS  $m/z$  = 457 (M+H)<sup>+</sup>







**Title Compound:** Compound 2 (0.061 g; 0.13 mmol) was treated with 7 N ammonia in methanol (2 mL) and heated to 60°C for 1 h in a sealed tube. The solvent was removed by rotary evaporation and the crude residue was purified by preparative reverse phase HPLC to give 0.030 g of the title compound as a white solid. LRMS  $m/z$  = 379 (M+H)<sup>+</sup>







15







#### EXAMPLES 854 TO 915







Examples 854 to 915 were prepared using methodology described in Example 25 and Example 853.







Example	Structure	M+H
854		411
855		440








Example	Structure	M+H
856		503
857		520
858		478
859		477
860		499
861		434







Example	Structure	M+H
862		408
863		480
864		551
865		451
866		489
867		411







Example	Structure	M+H
868		440
869		503
870		503
871		508
872		478
873		501








Example	Structure	M+H
874		423
875		452
876		515
877		515
878		478
879		487





Example	Structure	M+H
880		409
881		454
882		438
883		505
884		427
885		512


Example	Structure	M+H
886		505
887		507
888		429
889		523
890		494
891		445
892		416

Example	Structure	M+H
893		458
894		408
895		428
896		465
897		408
898		492

Example	Structure	M+H
899		476
900		555
901		477
902		415
903		398
904		507


Example	Structure	M+H
905		494
906		428
907		415
908		447
909		393
910		423
911		438

Example	Structure	M+H
912		425
913		471
914		428
915		415

EXAMPLE 916

5

3-Amino-pyrazine-2-carboxylic acid (4-phenyl-1-pyrimidin-2-yl-piperidin-4-ylmethyl)-amide

Synthesis

5           **Compound 1:** Compound **1** was prepared using methodology described in Example **653**.






**Title Compound:** The title compound was prepared using methodology described in Example 521 and purified by preparative reverse phase HPLC to give a white solid. LRMS  $m/z$  390 (M+H)<sup>+</sup>.


10

EXAMPLES 917 TO 924


Examples 917 to 924 were prepared using methodology described in Example 916.

Example	Structure	Name	M+H
917		3-Amino-pyrazine-2-carboxylic acid [1-(4-(3-trifluoromethyl-phenyl)-piperidin-4-ylmethyl)]-amide	459
918		3-Amino-pyrazine-2-carboxylic acid [1-(4-methoxy-pyrimidin-2-yl)-4-phenyl-piperidin-4-ylmethyl]-amide	421
919		3-Amino-pyrazine-2-carboxylic acid [4-phenyl-1-(4-trifluoromethyl-pyrimidin-2-yl)-piperidin-4-ylmethyl]-amide	459

Example	Structure	Name	M+H
920		3-Amino-pyrazine-2-carboxylic acid [4-(3-fluoro-phenyl)-1-pyrimidin-2-yl-piperidin-4-ylmethyl]-amide	409
921		3-Amino-pyrazine-2-carboxylic acid [4-(3,5-difluoro-phenyl)-1-pyrimidin-2-ylpiperidin-4-ylmethyl]-amide	410
922		3-Amino-pyrazine-2-carboxylic acid [4-(2,5-difluoro-phenyl)-1-pyrimidin-2-yl-piperidin-4-ylmethyl]-amide	427
923		N-[4-(2,3-Difluoro-phenyl)-1-pyrimidin-2-yl-piperidin-4-ylmethyl]-2-methoxy-benzamide	439
924		3-Amino-pyrazine-2-carboxylic acid [4-(2,3-difluoro-phenyl)-1-pyrimidin-2-yl-piperidin-4-ylmethyl]-amide	426

EXAMPLE 925

3-Amino-pyrazine-2-carboxylic acid (4-phenyl-1-pyrimidin-  
2-yl-piperidin-4-ylmethyl)-amide

Synthesis

**Compound 1:** Compound **1** was prepared as described in Example **653**.


**Title Compound:** A solution of compound **1** (0.072 g; 0.23 mmol) in anhydrous acetonitrile (2 mL) was treated with polystyrene-diisopropylethylamine (300 mg) and hydrocinnamoyl chloride (0.045 g; 0.27 mmol) at room temperature. The reaction was shaken for 24 h, filtered and concentrated by rotary evaporation. The crude residue was purified by preparative reverse phase HPLC to give 0.054 g of the title compound as white solid. LRMS  $m/z = 444$  (M+H)<sup>+</sup>

EXAMPLES 926 TO 929

Examples 926 to 929 were prepared using methodology described in Example 925.


Example	Structure	Name	M+H
926		4-[[[(3-Amino-pyrazine-2-carbonyl)-amino]-methyl]-4-phenyl-piperidine-1-carboxylic acid ethyl ester	385
927		3-Amino-pyrazine-2-carboxylic acid (1-benzoyl-4-phenyl-piperidin-4-ylmethyl)-amide	417
928		3-Amino-pyrazine-2-carboxylic acid [4-phenyl-1-(pyridine-3-carbonyl)-piperidin-4-ylmethyl]-amide	418
929		3-Amino-pyrazine-2-carboxylic acid [4-phenyl-1-(pyridine-4-carbonyl)-piperidin-4-ylmethyl]-amide	418

5

EXAMPLE 930

3-Amino-pyrazine-2-carboxylic acid [1-(1-amino-2-nitro-vinyl)-  
4-phenyl-piperidin-4-ylmethyl]-amide


10

Synthesis

5           **Compound 1:** Compound **1** was prepared as described in Example **653**.


**Compound 2:** A solution of compound **1** (0.065 g; 0.21 mmol) in anhydrous acetonitrile (3 mL) was treated with 1,1-bis(methylthio)-2-nitroethylene (0.058 g; 0.35 mmol) and heated at 85°C for 3 h. The solvent was removed by rotary evaporation and the crude residue was purified by recrystallization from ethyl acetate to give 0.068 g of compound **2** as a bright yellow solid. LRMS  $m/z = 430$  (M+H)<sup>+</sup>

**Title Compound:** Compound **2** (0.042 g; 0.098 mmol) was treated with 7 N ammonia in methanol (1 mL) and heated at 85°C for 15 min in a sealed tube. The solvent was removed by rotary evaporation and the crude residue was purified directly by column chromatography on silica gel using 9:1 ethyl acetate:hexane as the eluent to give 0.018 g of the title compound as a white solid. LRMS  $m/z = 398$  (m+H)<sup>+</sup>

EXAMPLE 931

20           3-Amino-pyrazine-2-carboxylic acid [1-(3,5-dimethyl-isoxazole-4-sulfonyl)-4-phenyl-piperidin-4-ylmethyl]-amide

# Synthesis



**Compound 1:** Compound 1 was prepared using methodology described in Example 653.




**Title Compound:** A solution of compound 1 (0.043 mg; 0.14 mmol) in anhydrous acetonitrile (1 mL) was treated with triethylamine (0.1 mL; 0.7 mmol) and 3,5-dimethyl-isoxazole-4-sulfonyl chloride (0.040 mg; 0.2 mmol) and allowed to stir at room temperature for 0.5 h. The solvent was removed by rotary evaporation and the crude residue was purified directly by column chromatography on silica gel using 1:1 ethyl acetate:hexane as the eluent to give 0.012 g of the title compound as a white solid. LRMS  $m/z = 472$  (m+H)<sup>+</sup>

15


## EXAMPLES 932 TO 936

Examples 932 to 936 were prepared using methodology described in Example 931.

Example	Structure	Name	M+H
932		3-Amino-pyrazine-2-carboxylic acid [1-(3-fluorobenzenesulfonyl)-4-phenylpiperidin-4-ylmethyl]-amide	471
933		3-Amino-pyrazine-2-carboxylic acid [1-(4-fluorobenzenesulfonyl)-4-phenylpiperidin-4-ylmethyl]-amide	471


Example	Structure	Name	M+H
934		3-Amino-pyrazine-2-carboxylic acid [1-(3-cyanobenzenesulfonyl)-4-phenylpiperidin-4-ylmethyl]-amide	478
935		3-Amino-pyrazine-2-carboxylic acid [1-(2-methanesulfonylbenzenesulfonyl)-4-phenylpiperidin-4-ylmethyl]-amide	531
936		3-Amino-pyrazine-2-carboxylic acid [4-phenyl-1-(propane-2-sulfonyl)piperidin-4-ylmethyl]-amide	419

EXAMPLE 937



5 1-{4-[(1H-Indazol-3-ylamino)-methyl]-4-phenyl-piperidin-1-yl}-  
3-phenyl-propan-1-one

# Synthesis



**Compound 1:** Compound 1 was prepared as described in Example 653.

**Compound 2:** To compound 1 (0.89 g, 3.06 mmol) and triethylamine (0.95 g, 9.36 mmol) in dichloromethane (5 mL) was added 2-fluorobenzoyl chloride (0.53 g, 3.37 mmol). After 1 h, the reaction mixture was diluted with diethyl ether (20 mL) then washed with 1 N sodium hydroxide, water and saturated aqueous sodium chloride. The organic layer was separated, dried (sodium sulfate), filtered and concentrated. Purification by silica gel chromatography using 2:1 hexanes:ethyl acetate as the eluent gave compound 2 (344 mg, 27%) as a colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.43 (9H, s), 1.82-1.91 (2 H, m), 2.15-2.20 (2 H, m), 3.18-3.26 (2 H, m), 3.68-3.74 (4 H, m), 6.32-6.36 (1 H, m), 7.03 (1 H, dd, *J*= 6.1, 12.0 Hz), 7.20-7.46 (7 H, m), 8.03 (1 H, td, *J*= 1.8, 7.9 Hz). LRMS *m/z* 357 (M+H)<sup>+</sup>.


**Compound 3:** Compound 2 (340 mg, 0.82 mmol) and Lawesson's reagent (433 mg, 1.07 mmol) in toluene (3 mL) was heated at 100°C for 3 h then cooled to room temperature. Water (1 mL), saturated aqueous sodium bicarbonate (3 mL) and ethyl acetate (3 mL) were added. After stirring for 20 min, the reaction mixture was diluted with ethyl acetate then washed with water and saturated aqueous sodium chloride. The organic layer was separated, dried (sodium sulfate), filtered and

concentrated. Purification by silica gel chromatography using 3:1 hexanes:ethyl acetate as the eluent gave compound **3** (255 mg, 72%) as a yellow solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.44 (9 H, s), 1.94-1.98 (2 H, m), 2.17-2.25 (2 H, m), 3.27-3.34 (2 H, m), 3.67-3.75 (2 H, m), 4.15 (2 H, d, *J* = 5.0 Hz), 6.95-7.01 (1 H, m), 7.15 (1 H, td, *J* = 1.0, 7.4 Hz), 7.27-7.63 (7 H, m), 8.07 (1 H, td, *J* = 1.9, 8.0 Hz). LRMS *m/z* 429 (M+H)<sup>+</sup>.


**Compound 4:** Compound **3** (250 mg, 0.58 mmol) and hydrazine (187 mg, 5.80 mmol) in 1,4-dioxane (3 mL) was heated at 100°C for 3 days then cooled to room temperature. The reaction mixture was diluted with ethyl acetate (10 mL) then washed with saturated aqueous sodium bicarbonate, water and saturated aqueous sodium chloride. The organic layer was separated, dried (sodium sulfate), filtered and concentrated. Purification by silica gel chromatography using 3:1 hexanes:ethyl acetate as the eluent gave compound **4** (149 mg, 63%) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.43 (9 H, s), 1.90-1.97 (2 H, m), 2.20-2.25 (2 H, m), 3.16-3.25 (2 H, m), 3.64 (2 H, s), 3.64-3.69 (3 H, m), 6.98 (1 H, td, *J* = 1.5, 7.9 Hz), 7.29-7.35 (5 H, m), 7.42 (4 H, d, *J* = 4.4 Hz). LRMS *m/z* 407 (M+H)<sup>+</sup>.

**Compound 5:** Compound **4** (143 mg, 0.35 mmol) in dichloromethane (0.75 mL) and trifluoroacetic acid (0.25 mL) was stirred for 1.5 h then concentrated under reduced pressure. 1N Sodium hydroxide (5 mL) was added then extracted with dichloromethane (3 x 10 mL). The combined organic layers were dried (sodium sulfate), filtered and concentrated to give the compound **5** (107 mg, 100%) as a white solid. LRMS *m/z* 307 (M+H)<sup>+</sup>.

**Title Compound:** The title compound was prepared using methodology described in Example 390. LRMS *m/z* 440 (M+H)<sup>+</sup>.

EXAMPLE 938

4-[(2-Methoxy-benzoylamino)-methyl]-4-phenyl-piperidine-  
1-carboxylic acid dimethylamide

Synthesis

**Compound 1:** Compound 1 was prepared as described in Example 15.





**Compound 2:** A solution of compound 1 (0.63 g; 1.9 mmol) in tetrahydrofuran (25 mL) was treated with triethylamine (0.33 mL; 2.4 mmol) and 4-nitro phenyl chloroformate (0.47 g; 2.3 mmol) at room temperature. After 24 h the solvent was removed by rotary evaporation and the crude residue was purified directly by column chromatography on silica gel using 1:1 ethyl acetate:hexane as the eluent to give 0.53 g of compound 2 as a white foam. LRMS  $m/z$  491 ( $M+H$ )<sup>+</sup>.


**Title Compound:** Compound 2 (0.050 g; 0.10 mmol) was treated with 2 M dimethylamine in tetrahydrofuran (2 mL) and heated to 65°C in a sealed tube for 12 h. The solvent was removed by rotary evaporation and the crude residue was purified by

preparative reverse phase HPLC to give 0.018 g of the title compound as a white solid. LRMS  $m/z$  396 (M+H)<sup>+</sup>.

### EXAMPLES 939 TO 942


5 Examples 939 to 942 were prepared using methodology described in Example 938.

Example	Structure	Name	M+H
939		N-[1-(3-Hydroxy-pyrrolidine-1-carbonyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide	440
940		2-Methoxy-N-[1-(morpholine-4-carbonyl)-4-phenyl-piperidin-4-ylmethyl]-benzamide	440
941		2-Methoxy-N-[4-phenyl-1-(pyrrolidine-1-carbonyl)-piperidin-4-ylmethyl]-benzamide	424
942		4-[(2-Methoxy-benzoylamino)-methyl]-4-phenyl-piperidine-1-carboxylic acid isopropyl-methyl-amide	426

EXAMPLE 943

2-Methoxy-N-(4-phenyl-piperidin-4-ylmethyl)-thiobenzamide

5

Synthesis





10        **Compound 1:** Compound **1** was prepared using methodology described in Example **653**.

15        **Compound 2:** A solution of compound **1** (0.32 g; 0.76 mmol) and Lawesson's reagent (0.38 g; 0.95 mmol) in 10% pyridine in toluene (5 mL) was heated at reflux for 3 h. The reaction mixture was cooled to room temperature and  
 20        diluted with ethyl acetate (5 mL), water (2 mL) and saturated aqueous sodium bicarbonate (5 mL) and stirred for 0.5 h. Additional ethyl acetate was added and the organic layer was separated, washed with water and saturated aqueous sodium chloride, dried (sodium sulfate), filtered and concentrated. The crude residue was purified by column chromatography on silica gel using 3:1 hexane:ethyl acetate as the  
 20        eluent to give 0.24 g of compound **2** as a yellow solid. LRMS  $m/z$  442 (M+H)<sup>+</sup>.


25        **Title Compound:** Compound **2** (0.044 g; 0.10 mmol) was treated with 0.75 mL dichloromethane and 0.25 mL trifluoroacetic acid and the reaction mixture was stirred at room temperature for 0.5 h. The solvents were removed by rotary evaporation to give 0.040 g of the title compound as the trifluoroacetic acid salt as a  
 25        white solid that was used without additional purification. LRMS  $m/z$  341 (M+H)<sup>+</sup>.

EXAMPLES 944 TO 947

Examples 944 to 947 were prepared using methodologies described in Example 943 and Example 15, Example 16 or Example 25.


Example	Structure	Name	M+H
944		N-(1-Dimethylsulfamoyl-4-phenyl-piperidin-4-ylmethyl)-2-methoxy-thiobenzamide	450
945		2-Methoxy-N-(4-phenyl-1-sulfamoyl-piperidin-4-ylmethyl)-thiobenzamide	422
946			487
947			410

5

EXAMPLE 948

N-[1-(N-Ethylcarbamimidoyl)-4-phenyl-piperidin-4-ylmethyl]-2-methoxy-benzamide

10


Synthesis

5        **Compound 1:** Compound 1 was prepared using methodology described in Example 15.

**Compound 2:** Compound 2 was prepared as described in *J. Org. Chem.* **2002**, 67, 7553-7556.


**Compound 3:** A solution of compound 1 (0.63 g; 1.9 mmol) and compound 2  
 10 (0.45 g; 2.8 mmol) in tetrahydrofuran (15 mL) was heated at 50°C for 24 h. The solvent was removed by rotary evaporation and the residue was treated with ethyl acetate (50 mL) and water (50 mL). The organic layer was separated, washed with water and saturated aqueous sodium chloride. The organic layer was separated, dried (sodium sulfate), filtered and concentrated to give 0.7 g of compound 3 as a light  
 15 yellow foam that was used without additional purification. LRMS  $m/z$  419 (M+H)<sup>+</sup>.

**Title Compound:** Compound 3 (0.059 g; 0.14 mmol) was treated with 2 M ethylamine in tetrahydrofuran (2.5 mL) and heated to 60°C in a sealed tube for 48 h. The solvent was removed by rotary evaporation and the crude residue was purified by preparative reverse phase HPLC to give 0.022 mg of the title compound as the  
 20 trifluoroacetic acid salt as a white solid. LRMS  $m/z$  396 (M+H)<sup>+</sup>.

EXAMPLE 949

3-Amino-pyrazine-2-carboxylic acid (4-p-tolyl-tetrahydro-pyran-4-ylmethyl)-amide

5

Synthesis

10      **Compound 1:** Compound 1 is commercially available.







15      **Compound 2:** A solution of compound 1 (0.71 g; 5 mmol), 4-methyl benzyl cyanide (0.66 g; 5 mmol) and hexadecyl tributyl phosphonium bromide (0.13 g; 0.25 mmol) in 50% NaOH in water (8 mL) was heated at 100°C for 2 h. The reaction was cooled to room temperature, diluted with water (20 mL) and extracted with diethyl ether (3 x 10 mL). The combined organic layers were washed with water and saturated aqueous sodium chloride, dried (sodium sulfate), filtered and concentrated. The crude residue was purified by column chromatography on silica gel using 8.5:1.5 hexane: ethyl acetate as the eluent to give 0.776 g of compound 2 as a yellow oil. LRMS  $m/z$  202 (M+H)<sup>+</sup>.


20      **Compound 3:** Compound 3 was prepared using methodology described in Example 15. LRMS  $m/z$  206 (M+H)<sup>+</sup>.

**Title Compound:** The title compound was prepared using methodology described in Example **653**. LRMS  $m/z$  327 (M+H)<sup>+</sup>.


### EXAMPLES 950 TO 955

5 Examples 950 to 955 were prepared using methodology described in Example 949.

Example	Structure	Name	M+H
950		3-Amino-pyrazine-2-carboxylic acid (4-phenyl-tetrahydro-pyran-4-ylmethyl)-amide	313
951		3-Amino-pyrazine-2-carboxylic acid [4-(4-chloro-phenyl)-tetrahydro-pyran-4-ylmethyl]-amide	348
952		3-Amino-pyrazine-2-carboxylic acid [4-(3-fluoro-phenyl)-tetrahydro-pyran-4-ylmethyl]-amide	331
953		3-Amino-pyrazine-2-carboxylic acid [4-(4-methoxy-phenyl)-tetrahydro-pyran-4-ylmethyl]-amide	343
954		3-Amino-pyrazine-2-carboxylic acid [4-(2,4-difluoro-phenyl)-tetrahydro-pyran-4-ylmethyl]-amide	349
955		3-Amino-pyrazine-2-carboxylic acid [4-(4-fluoro-phenyl)-tetrahydro-pyran-4-ylmethyl]-amide	331

EXAMPLE 9562-Methoxy-*N*-(4-phenyl-azepan-4-ylmethyl)-benzamide

5

Synthesis**Compound 1:** Compound 1 is commercially available

10 **Compound 2:** To a solution of ketone 1 (500mg, 2.51 mmol) in glacial acetic acid (30mL) was added concentrated sulfuric acid (0.3mL) at room temperature. The solution was heated to 65°C and sodium azide (0.50g, 7.7mmol) was added in 3 equal portions over 5 min at 65°C. After a further 5 min at 65°C, the reaction mixture was

15 poured cautiously into a saturated NaHCO<sub>3</sub> solution (ca. 100mL), transferred to a separation funnel and the aqueous portion extracted with dichloromethane (3 x 50mL). The combined organic portions were washed with NaHCO<sub>3</sub> (20mL), dried over Na<sub>2</sub>SO<sub>4</sub>, decanted and concentrated yielding a pale yellow oil. Methanol (ca. 5mL) was added to the oil and the white precipitate was collected and dried under

- high vacuum (153mg). The methanol solution was purified by preparative HPLC. YMC ODS S5 30 x 50mm, 10 min gradient, 0 – 100% MeOH (90% in water, 0.1% TFA) UV detection 220nm, 50mL/min flow rate. The product retention time was 5.23 min. The product was neutralized with saturated NaHCO<sub>3</sub> and extracted into dichloromethane. A further 119 mg of product was isolated and combined with the product isolated by precipitation, (combined mass 272mg, yield 51%) . HPLC Rt 2.17min, Purity 98%, Phenomenex Luna S5 column 4.6 x 50 mm, 4min gradient 0 to 100% MeOH (90% in water, 0.1% PPA) UV detection at 220nm. LCMS Rt 1.12min, [M+1] 215.42 Phenomenex S5 column 4.6 x 30 mm, 2min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (DMSO) 1.98ppm, 1H, multiplet; 2.10ppm, 1H, multiplet; 2.30ppm, 1H, multiplet; 2.78ppm, 1H, multiplet; 3.21ppm, 1H, multiplet; 3.43ppm, 1H, multiplet; 7.31ppm, 1H, t, J=7.9Hz; 7.42ppm, 2H, dd, J=7.9Hz and J=7.9Hz; 7.55ppm, 2H, d, J=8.4Hz; 7.82ppm, 1H, s.
- Compound 3:** Benzoic anhydride (4.22 g, 18.7 mmol) and pyridine (3.1 mL, 37 mmol) were added to a suspension of compound **2** (2.00 g, 9.33 mmol) in toluene (20 mL). After heating at 100°C for 18 hours, the reaction mixture was concentrated and purified by ISCO hexane/EtOAc, 0%EtOAc – 10% EtOAc over 10 minutes, 10% EtOAc – 30% EtOAc over 10 minutes, 30% EtOAc for 20 minutes, 30% EtOAc – 100% EtOAc over 2 minutes, 100% EtOAc for 5 minutes. Compound **3** eluted at a retention time of 12.8min as a white solid (2.62 g, 88% yield). HPLC Rt 3.08min, Purity 86%, Phenomenex Luna S5 column 4.6 x 50 mm, 4min gradient 0 to 100% MeOH (90% in water, 0.1% PPA) UV detection at 220nm. LCMS Rt 1.57min, [M+1] 319.18 Phenomenex S5 column 4.6 x 30 mm, 2min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 2.23ppm, 2H, multiplet; 2.37ppm, 1H, multiplet; 2.47ppm, 1H, multiplet; 2.76ppm, 1H, multiplet; 3.34ppm, 1H, multiplet; 3.91ppm, 1H, dd, J=12.0 and 16.0Hz; 4.66ppm, 1H, dd, J=12.0 and 16.0Hz; 7.44ppm, 8H, multiplet; 7.57ppm, 2H, multiplet.
- Compound 4:** To a solution of compound **3** (500 mg, 1.57 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (8 mL) and THF (4 mL) was added a 1.0 M solution of lithium aluminum hydride in THF (9.4 mL, 9.4 mmol) at 0°C. The reaction mixture was allowed to warm up to


room temperature and stirred for 18 hours. The reaction was quenched with H<sub>2</sub>O (1.7 mL), 1 N NaOH (1.0 mL) and H<sub>2</sub>O (1.7 mL). After stirring at room temperature for 30 minutes, the reaction was filtered and the filtrate was concentrated. The resulting residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (75 mL). The organic layer was washed with brine (25 mL), dried over MgSO<sub>4</sub>, filtered, concentrated, and the resulting residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (2.0 mL). The solution was added to a solution of O-anisic acid (215 mg, 1.42 mmol) and EDCI (296 mg, 1.54 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL). After 2 hours, the reaction mixture was concentrated and purified by ISCO Hexane/EtOAc; 0% EtOAc – 30% EtOAc over 10 minutes, 30% EtOAc – 50% EtOAc over 10 minutes, 50% EtOAc for 10 minutes, 50% EtOAc – 100% EtOAc over 5 minutes, 100% EtOAc for 5 minutes. Compound **N+2** eluted at a retention time of 13.3min as a white solid compound (211 mg, 31%). HPLC Rt 2.40min, Purity 100%, Phenomenex Luna S5 column 4.6 x 50 mm, 4min gradient 0 to 100% MeOH (90% in water, 0.1% PPA) UV detection at 220nm. LCMS Rt 1.46min, [M+1] 429.22

Phenomenex S5 column 4.6 x 30 mm, 2min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.63ppm, 1H, multiplet; 1.75ppm, 1H, multiplet; 1.89ppm, 1H, multiplet; 1.99ppm, 1H, multiplet; 2.22ppm, 2H, multiplet; 2.61ppm, 4H, multiplet; 2.92ppm, 3H, s; 3.54ppm, 2H, s; 3.64ppm, 2H, d, J=4.0Hz; 6.75ppm, 1H, d, J=8.0Hz; 6.96ppm, 1H, t, J=6.0 Hz; 7.20ppm, 6H, multiplet; 7.30ppm, 5H, multiplet; 7.50ppm, 1H, multiplet; 8.11ppm, 1H, dd, J=4.0 and 8.0Hz.

**Title Compound:** At 0°C 1-chloroethyl chloroformate (159 µL, 1.48 mmol) was added to a solution of compound **4** (211 mg, 0.492 mmol) and TEA (341 µL, 2.45 mmol) in dichloroethane (5 mL). The reaction was allowed to warm up to room temperature and stirred for 3 hours. The reaction mixture was concentrated and dried on oil pump for 0.5 hours. The solution of the resulting residue in MeOH (5 mL) was heated under reflux for 5 hours and concentrated. Crude product was purified by preparative HPLC YMC ODS S5 30 x 100 mm Ballistic column 10-100% MeOH (90% in water, 0.1%TFA) gradient over 10min with flow rate 40mL/min and UV detection at 220nm. **Compound 5** eluted at a retention time of 6.6mins and was isolated as a yellow oil (101.3mg, yield 61%) NMR H (CDCl<sub>3</sub>) 1.60ppm, 1H,


multiplet; 1.91ppm, 3H, multiplet; 2.34ppm, 2H, multiplet; 3.22ppm, 2H, multiplet;  
 3.36ppm, 1H, multiplet; 3.52ppm, 1H, multiplet; 3.53ppm, 3H, s; 3.62ppm, 2H,  
 multiplet; 6.76ppm, 1H, d, J=1.8Hz; 6.80ppm, 1H, d, J=7.9Hz; 6.98ppm, 1H, t,  
 J=7.7 Hz; 7.06ppm, 1H, d, J=1.8Hz; 7.22ppm, 1H, multiplet; 7.34ppm, 3H,  
 5 multiplet; 7.56ppm, 1H, t, J=5.7Hz; 8.11ppm, 1H, dd, J=1.8 and 7.5Hz.

### EXAMPLE 957



10 *N*-(1-Benzyl-4-phenyl-azepan-4-ylmethyl)-2-methoxy-benzamide

### Synthesis:



15


**Compound 1:** Compound 1 was synthesized as described in **Example 956**.

**Compound 2:** NaH (95%) (71 mg, 2.8 mmol) was added to the suspension of  
 compound 1 (500 mg, 2.33 mmol) in DMF (10 mL). After 30 minutes benzylbromide  
 (333  $\mu$ L, 2.80 mmol) was added and the reaction was stirred for 18 hours. LiCl (10%,  
 20 50 mL) was added. The aqueous layer was extracted with EtOAc (3 x 25 mL). The  
 combined organic layers were dried over  $\text{MgSO}_4$ , filtered, concentrated and purified  
 by ISCO Hexane/EtOAc, 0% EtOAc to 10% EtOAc over 10 minutes, 10% EtOAc for


5 minutes, 10% EtOAc to 30% EtOAc over 10 minutes, 30% EtOAc for 10 minutes, 30% EtOAc to 100% EtOAc over 5 minutes. Compound N + 1' eluted at 23min as a clear oil (454.7 mg, 64% yield).

**Title Compound:** At room temperature a 1.0 M solution of Lithium aluminum hydride in THF (480  $\mu$ L, 0.480 mmol) was added to a solution of compound **2** (48.5 mg, 0.160 mmol) in THF (1 mL). After 3 hours the reaction mixture was quenched with H<sub>2</sub>O (60  $\mu$ L), 1 N NaOH (36  $\mu$ L) and H<sub>2</sub>O (60  $\mu$ L). The reaction was stirred at room temperature for 0.5 hours, dried over MgSO<sub>4</sub>, filtered and concentrated. The solution of the resulting residue in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) was added to the mixture of o-anisic acid (27 mg, 0.18 mmol) and EDCI (37 mg, 0.19 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL). After 1 hours the reaction was concentrated and purified by preparative HPLC yielding compound **N+2** as a yellow oil (17.5 mg, 26%).

#### EXAMPLE 958



2-Methoxy-N-(4-phenyl-1-sulfamoyl-azepan-4-ylmethyl)-benzamide

Synthesis

5            **Compound 1:** Compound 1 was synthesized as described in Example 956.

**Compound 2:** At 0°C N,N'-sulfuryl bis-2-methylimidazole mono-methyl  
triflate salt (317 mg, 0.390 mmol) was added to a solution of compound 1 (101 mg,  
0.300 mmol) in acetonitrile (10 mL) followed by the addition of TEA (100 µL). The  
reaction was allowed to warm up to room temperature and stirred for 18 hours. The  
10 mixture was concentrated and purified by ISCO hexane (0.1% TEA)/EtOAc, 0%  
EtOAc – 50% EtOAc over 15 minutes, 50% EtOAc for 15 minutes, 50% EtOAc –  
100% EtOAc over 5 minutes, 100% EtOAc for 5 minutes. Compound 2 eluted at a  
retention time of 26min as a yellow oil (78.8 mg, 54% yield) NMR H (CDCl<sub>3</sub>)  
1.58ppm, 1H, multiplet; 1.89ppm, 3H, multiplet; 2.35ppm, 2H, multiplet; 2.45ppm,  
15 3H, s; 3.20ppm, 2H, multiplet; 3.37ppm, 1H, multiplet; 3.51ppm, 1H, multiplet;  
3.53ppm, 3H, s; 3.62ppm, 2H, multiplet; 6.76ppm, 1H, d, J=1.8Hz; 6.80ppm, 2H,  
multiplet; 6.98ppm, 1H, t, J=7.5 Hz; 7.06ppm, 1H, d, J=1.8Hz; 7.23ppm, 2H,  
multiplet; 7.33ppm, 3H, multiplet; 7.56ppm, 1H, t, J=5.7Hz; 8.10ppm, 1H, dd,  
J=1.8 and 7.9Hz



20            **Title Compound:** A solution of compound 2 (34 mg, 0.070 mmol) in CH<sub>2</sub>Cl<sub>2</sub>  
(1 mL) was added methyl trifluoromethanesulfonate (8 µL, 0.08 mmol) at 0°C. After




1.5 hours, the reaction was concentrated to give crude product of compound **3** as a white foam. This product was used directly to the next step without further purification. A 2.0 M solution of NH<sub>3</sub> in MeOH (170  $\mu$ L, 0.35 mmol) was added to a solution of compound **4** and TEA (100  $\mu$ L) in acetonitrile (1 mL). The reaction mixture was heated at 80°C for 8 hours. The concentrated reaction mixture was purified by preparative HPLC YMC ODS S5 30 x 100 mm Ballistic column 10-100% MeOH (90% in water, 0.1%TFA) gradient over 12min with flow rate 40mL/min and UV detection at 220nm. **Compound 5** eluted at a retention time of 9.6mins and was isolated as a clear oil (18.27 mg, 63% yield from compound **3**). HPLC Rt 2.91 min, Purity 100%, Phenomenex Luna S5 column 4.6 x 50 mm, 4min gradient 0 to 100% MeOH (90% in water, 0.1% PPA) UV detection at 220nm. LCMS Rt 1.60min, [M+1] 418.17 YMC-ODS S5 column 4.6 x 33 mm, 2min gradient 0 to 100% MeOH (90% in water, 10 mM NH<sub>4</sub>OAc) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.65ppm, 1H, multiplet; 1.84ppm, 2H, multiplet; 1.99ppm, 2H, multiplet; 2.25ppm, 1H, multiplet; 2.37ppm, 2H, multiplet; 3.23ppm, 2H, multiplet; 3.31ppm, 1H, multiplet; 3.52ppm, 3H, s; 3.57ppm, 2H, s; 3.69ppm, 1H, multiplet; 4.40ppm, 2H, bs; 6.79ppm, 1H, d, J=8.4Hz; 6.97ppm, 1H, t, J=7.5Hz; 7.23ppm, 1H, multiplet; 7.34ppm, 5H, multiplet; 7.61ppm, 1H, multiplet; 8.09ppm, 1H, dd, J=1.7 and 7.9Hz.


20

EXAMPLES 959 TO 963


Compounds 959 to 963 were synthesized using methodology described in Example 958.

Example	Structure	Name	[M+1]
959		2-Methoxy-N-[1-(2-methoxy-ethylsulfamoyl)-4-phenyl-azepan-4-ylmethyl]-benzamide	476
960		N-[1-(2-Hydroxy-ethylsulfamoyl)-4-phenyl-azepan-4-ylmethyl]-2-methoxy-benzamide	462

Example	Structure	Name	[M+1]
961		2-Methoxy-N-{1-[(2-methoxy-ethyl)-methyl-sulfamoyl]-4-phenyl-azepan-4-ylmethyl}-benzamide	490
962		N-(1-Cyclopropylsulfamoyl-4-phenyl-azepan-4-ylmethyl)-2-methoxy-benzamide	458
963		2-Methoxy-N-(1-methylsulfamoyl-4-phenyl-azepan-4-ylmethyl)-benzamide	432

EXAMPLE 964

5 N-(1-Dimethylsulfamoyl-4-phenyl-azepan-4-ylmethyl)-2-methoxy-benzamide

Synthesis

10

**Compound 1:** Compound 1 was synthesized as described in Example 956.


**Title Compound:** Dimethylsulfomoyl chloride (3.44 mg, 0.0240 mmol) was added to a solution of compound 1 (6.76 mg, 0.0200 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 mL).

TEA (5  $\mu$ L) was added and the reaction was stirred at room temperature for 2 hours. The concentrated reaction mixture was purified by preparative HPLC YMC ODS S5 30 x 100 mm Ballistic column 20-100% MeOH (90% in water, 0.1%TFA) gradient over 10min with flow rate 40mL/min and UV detection at 220nm. **Compound 2**

- 5 eluted at a retention time of 9.0min and was isolated as a clear oil (5.7 mg, 64% yield). HPLC Rt 3.30min, Purity 100%, Phenomenex Luna S5 column 4.6 x 50 mm, 4min gradient 0 to 100% MeOH (90% in water, 0.1% PPA) UV detection at 220nm. LCMS Rt 1.68min, [M+1] 446.21 Phenomenex S5 column 4.6 x 30 mm, 2min gradient 0 to 100% MeOH (90% in water, 10 mM NH<sub>4</sub>OAc) UV detection at 220nm.
- 10 NMR H (CDCl<sub>3</sub>) 1.61ppm, 1H, multiplet; 1.89ppm, 3H, multiplet; 2.34ppm, 2H, multiplet; 2.61ppm, 6H, s; 3.18ppm, 2H, multiplet; 3.26ppm, 1H, multiplet; 3.54ppm, 3H, s; 3.57ppm, 3H, multiplet; 6.80ppm, 1H, d, J=8.0Hz; 6.98ppm, 1H, t, J=8.0Hz; 7.23ppm, 1H, multiplet; 7.33ppm, 5H, multiplet; 7.58ppm, 1H, multiplet; 8.11ppm, 1H, multiplet.

15


#### EXAMPLE 965



4-Aminomethyl-4-(3-fluoro-phenyl)-azepane-1-carboxylic acid tert-butyl ester

20

# Synthesis



5           **Compound 1:** Compound 1 was synthesized via the intermediate cyclohexanone according to the procedures described in Journal of Medicinal Chemistry, **1998**, 821. The intermediate cyclohexanone was converted to the lactam as described for **Example 956**.


10           **Compound 2:** To a suspension of the crude compound 1 in THF (20 mL) was added subsequently di-*tert*-butyl dicarbonate (2.33 mL, 10.1 mmol) and dimethylaminopyridine (1.24 g, 10.1 mmol). After 3 hours the reaction mixture was concentrated. The resulting residue was dissolved in Ethyl acetate (100 mL). The organic layer was washed with 1 N HCl (2 x 50mL) and dried over  $\text{MgSO}_4$ . Evaporation and purification by ISCO (hexane /EtOAc, 0% EtOAc – 10% EtOAc over 10 minutes, 10% EtOAc – 30% EtOAc over 10 minutes, 30% EtOAc for 15 minutes, 30% EtOAc – 100% EtOAc over 5 minutes, 100% EtOAc for 5 minutes) yielded compound 2 with a retention time of 16min as a white solid (2.69 g, 88% from ketone) NMR H ( $\text{CDCl}_3$ ) 2.03ppm, 1H, multiplet; 2.28ppm, 3H, multiplet; 2.76ppm, 1H, dd,  $J=7.2$  and 15.6 Hz; 3.21ppm, 1H, multiplet; 3.75ppm, 1H, dd,  $J=10.5$  and 16.3Hz; 4.44ppm, 1H, dd,  $J=6.5$  and 15.9Hz; 7.05ppm, 1H, multiplet; 7.16ppm, 1H, multiplet; 7.26ppm, 1H, multiplet; 7.38ppm, 1H, multiplet.

15

20


**Compound 3:** At  $-78^{\circ}\text{C}$  a 1.0 M solution of lithium triethylborohydride in THF (4.87 mL, 4.87 mmol) was added to a solution of compound **2** (1.35 g, 4.06 mmol) in THF (25 mL). After 30 minutes the reaction was quenched with saturated  $\text{NaHCO}_3$  (7.6 mL) and warmed to  $0^{\circ}\text{C}$ . At  $0^{\circ}\text{C}$   $\text{H}_2\text{O}_2$  (12 drops) was added and the  
5 reaction was stirred for 20 minutes. THF was removed and the aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (3 x 25 mL). The combined organic layers were dried over  $\text{MgSO}_4$ , filtered and concentrated. The resulting thick oil was dissolved in  $\text{CH}_2\text{Cl}_2$  (50 mL). At  $-78^{\circ}\text{C}$  triethylsilane (648  $\mu\text{L}$ , 4.06 mmol) was added followed by drop  
10 wise addition of boron trifluoride etherate (566  $\mu\text{L}$ , 4.47 mmol). After 30 minutes another portion of triethylsilane (648  $\mu\text{L}$ , 4.06 mmol) and boron trifluoride etherate (566  $\mu\text{L}$ , 4.47 mmol) were added and the reaction was continued to stir for 2 hours at  $-78^{\circ}\text{C}$ . The reaction was quenched with saturated  $\text{NaHCO}_3$  (15 mL). The aqueous  
layer was extracted with  $\text{CH}_2\text{Cl}_2$  (3 x 20 mL) and the combined organic layers were dried over  $\text{MgSO}_4$ , filtered and concentrated to give crude compound **3** (1.19 g) as an  
15 oil. This crude product was used to the next step directly without further purification.

**Title Compound:** To a solution of crude compound **3** (1.19 g, 3.73 mmol) in MeOH (35 mL) was added cobalt (II) chloride hexahydrate (1.41 g, 7.46 mmol). The resulted purple mixture was stirred at room temperature for 10 minutes. At  $0^{\circ}\text{C}$   $\text{NaBH}_4$  (1.41 g, 37.3 mmol) was added in three portions over 25 minutes. The  
20 reaction mixture was stirred at room temperature for 2 hours, concentrated to give a black residue. The black residue was dissolved in 30% ammonium hydroxide solution (100 mL), extracted with ethyl acetate (3 x 50 mL). The combined organic layers were dried over  $\text{MgSO}_4$ , filtered and concentrated to give a crude pink solid of compound **4** (1.46 g). This crude solid was used directly to the following acylation  
25 reaction.

EXAMPLE 966

N-[4-(3-Fluoro-phenyl)-azepan-4-ylmethyl]-2-methoxy-benzamide

5

Synthesis

10 **Compound 1:** The synthesis of **Compound 1** is described for **Example 965**.

**Compound 2:** To a solution of crude compound **1** (729 mg, 2.26 mmol), o-anisic acid (345 mg, 2.26 mmol) and TEA (314  $\mu$ L, 2.26 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was added PyBrOP (1.05 g, 2.26 mmol) at room temperature. After 3 hours, the reaction was concentrated and purified by flash chromatography using Hexane/EtOAc (2/1) to yield compound **2** as a white solid (536.2 mg, 58% from compound **N+13**).  
 15 HPLC Rt 3.79min, Purity 100%, Phenomenex Luna S5 column 4.6 x 50 mm, 4min gradient 0 to 100% MeOH (90% in water, 0.1% PPA) UV detection at 220nm.  
 LCMS Rt 2.04min,  $[\text{M}+1]$  357.43 Phenomenex S5 column 4.6 x 30 mm, 2min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR  
 20  $^1\text{H}$  ( $\text{CDCl}_3$ ) 1.38ppm, 9H, d,  $J=12.8\text{Hz}$ ; 1.64ppm, 1H, multiplet; 1.78ppm, 3H, multiplet; 2.32ppm, 2H, multiplet; 3.31ppm, 3H, multiplet; 3.65ppm, 6H, multiplet;





6.89ppm, 1H, d, J=8.4Hz; 6.99ppm, 1H, multiplet; 7.06ppm, 2H, multiplet;  
 7.16ppm, 1H, d, J=7.5Hz; 7.40ppm, 2H, multiplet; 7.62ppm, 1H, multiplet;  
 8.20ppm, 1H, d, J=7.5Hz.


- Title Compound:** To a solution of compound **2** (526 mg, 1.24 mmol) in  
 5 CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added a solution of TFA (1 mL) in CH<sub>2</sub>Cl<sub>2</sub> (4 mL) at room  
 temperature. After 1.5 hours the reaction was diluted with CH<sub>2</sub>Cl<sub>2</sub> (50 mL) and was  
 washed with saturated NaHCO<sub>3</sub> (2 x 20 mL), dried over MgSO<sub>4</sub>, filtered and  
 concentrated to give a crude product of compound **N+18** as a white solid (483 mg,  
 quantitative yield). The crude product was used directly to the next step without  
 10 further purification.

### EXAMPLES 967 TO 970


Compounds 967 to 970 were prepared using the methodology described for  
 Example 966 and Example 958.

15

Example	Structure	Name	[M+1]
967		N-[4-(3-Fluoro-phenyl)-1-(2-methyl-imidazole-1-sulfonyl)-azepan-4-ylmethyl]-2-methoxy-benzamide	501
968		N-[1-Cyclopropylsulfonyl-4-(3-fluoro-phenyl)-azepan-4-ylmethyl]-2-methoxy-benzamide	476
969		N-{4-(3-Fluoro-phenyl)-1-[(2-methoxy-ethyl)-methyl-sulfonyl]-azepan-4-ylmethyl}-2-methoxy-benzamide	508
970		N-[4-(3-Fluoro-phenyl)-1-sulfonyl-azepan-4-ylmethyl]-2-methoxy-benzamide	436

EXAMPLE 971

5

Synthesis

**Compound 1:** The synthesis of **Compound 1** is described for Example 965.




**Compound 2:** To a solution of crude compound **1** (729 mg, 2.26 mmol) and 3-aminopyrazine-2-carboxylic acid (377 mg, 2.26 mmol) in ethylene glycol dimethyl ether (10 mL) was added drop wise diethyl phosphoryl cyanide (411  $\mu$ L, 2.71 mmol) and TEA (630  $\mu$ L, 4.52 mmol) respectively at 0°C. The reaction was stirred at 0°C for 1 hour and at 40°C for 1 hour under N<sub>2</sub>. The mixture was diluted with EtOAc (100 mL) and washed with H<sub>2</sub>O (25 mL), saturated NaHCO<sub>3</sub> (25 mL) and H<sub>2</sub>O (25 mL) successively. The organic layer was dried over MgSO<sub>4</sub> and concentrated to afford the crude product, which was purified by ISCO Hexane/EtOAc; 0% EtOAc – 50% EtOAc over 10 minutes, 50% EtOAc for 25 minutes, 50% EtOAc – 100% EtOAc over 5 minutes, 100% EtOAc for 5 minutes. Compound **2** eluted at a retention time of 14min as a yellow solid (432.7 mg, 51% from compound **1**). HPLC Rt 3.67min, Purity 100%, Phenomenex Luna S5 column 4.6 x 50 mm, 4min gradient 0 to 100%


MeOH (90% in water, 0.1% PPA) UV detection at 220nm. LCMS Rt 1.90min, [M+1] 344.45 Phenomenex S5 column 4.6 x 30 mm, 2min gradient 0 to 100% MeOH (90% in water, 0.1% TFA) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.38ppm, 9H, d, J=12.8Hz; 1.62ppm, 1H, multiplet; 1.71ppm, 1H, mutiplet; 1.87ppm, 2H, multiplet; 2.34ppm, 2H, multiplet; 3.17ppm, 2H, multiplet; 3.38ppm, 1H, multiplet; 3.46ppm, 2H, multiplet; 3.67ppm, 1H, multiplet; 6.99ppm, 2H, multiplet; 7.11ppm, 1H, multiplet; 7.37ppm, 1H, multiplet; 7.71ppm, 2H, multiplet; 8.11ppm, 1H, s.

**Title Compound:** To a solution of compound 2 (457 mg, 1.03 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added a solution of TFA (1 mL) in CH<sub>2</sub>Cl<sub>2</sub> (4 mL) at room temperature. After 1.5 hours the reaction was diluted with CH<sub>2</sub>Cl<sub>2</sub> (50 mL) and was washed with saturated NaHCO<sub>3</sub> (2 x 20 mL), dried over MgSO<sub>4</sub>, filtered and concentrated to give a crude product of compound 3 as a yellow solid (291mg, 82% crude yield). The crude product was used directly to the next step without further purification.

#### EXAMPLES 972 TO 974

Compounds 972 to 974 were prepared using the methodology described for Example 971 and Example 961.

Example	Structure	Name	[M+1]
972		N-[1-Dimethylsulfamoyl-4-(3-fluorophenyl)-azepan-4-ylmethyl]-2-methoxybenzamide	464
973		3-Amino-pyrazine-2-carboxylic acid [1-dimethylsulfamoyl-4-(3-fluorophenyl)-azepan-4-ylmethyl]-amide	451
974		3-Amino-pyrazine-2-carboxylic acid [4-(3-fluorophenyl)-1-(2-methyl-imidazole-1-sulfonyl)-azepan-4-ylmethyl]-amide	488

EXAMPLE 975

5           **Compound 1:** **Compound 1** was prepared as described in **Example 671**.

**Compound 3:** N-(tert-Butoxycarbonyl)-N-[4-(dimethylazaniumylidene)-1,4-dihydropyridin-1-ylsulfonamide] was made according to Organic Letters, 2001, Vol. 3., No. 14, 2241-2243. **Compound 1** (19 mg, 0.058 mmol) was added to **Compound 2** (20 mg, 0.064 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) and was stirred at room temperature for 18 hours. The residue was concentrated and purified by preparative silica thin layer chromatography (25x25cm plate, 1mm thickness silica with UV indicator) using Hexane/EtOAc (1/2) as eluent to yield compound **3** as a clear oil (26 mg, 87% yield) NMR H (CDCl<sub>3</sub>) 1.44ppm, 9H, s; 1.64ppm, 1H, multiplet; 1.98ppm, 3H, multiplet; 2.37ppm, 2H, multiplet; 3.24ppm, 2H, multiplet; 3.49ppm, 3H, multiplet; 3.81ppm, 1H, multiplet; 6.99ppm, 2H, multiplet; 7.35ppm, 1H, multiplet; 7.67ppm, 1H, multiplet; 7.69ppm, 1H, d, J=4.0Hz; 8.00ppm, 1H, s; 8.12ppm, 1H, d, J=4.0Hz.

**Title Compound:** At room temperature, a solution of TFA (0.25 mL) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) was added to a solution of compound **3** (26 mg, 0.050 mmol). After 2 hours the reaction mixture was concentrated and purified by preparative HPLC YMC ODS S5 30 x 100 mm Ballistic column 20-100% MeOH (90% in water, 0.1%TFA) gradient over 10min with flow rate 40mL/min and UV detection at 220nm.


**Compound 4** eluted at a retention time of 7.0min and was isolated as a clear oil (20.2

mg, 95% yield). HPLC Rt 2.68min, Purity 100%, Phenomenex Luna S5 column 4.6 x 50 mm, 4min gradient 0 to 100% MeOH (90% in water, 0.1% PPA) UV detection at 220nm. LCMS Rt 1.39min, [M+1] 423.31 Phenomenex S5 column 4.6 x 30 mm, 2min gradient 0 to 100% MeOH (90% in water, 10 mM NH<sub>4</sub>OAc) UV detection at 220nm. NMR H (CDCl<sub>3</sub>) 1.66ppm, 1H, multiplet; 1.90ppm, 2H, multiplet; 2.00ppm, 1H, multiplet; 2.30ppm, 1H, multiplet; 2.41ppm, 1H, multiplet; 3.16ppm, 2H, multiplet; 3.41ppm, 1H, multiplet; 3.51ppm, 2H, multiplet; 3.66ppm, 1H, multiplet; 7.00ppm, 2H, multiplet; 7.10ppm, 1H, d, J=8.0Hz; 7.38ppm, 1H, multiplet; 7.60ppm, t, J=6.0Hz; 7.81ppm, 1H, d, J=4.0Hz; 7.93ppm, 1H, multiplet.

10

We claim:

1. A compound of formula I






enantiomers, diastereomers, salts and solvates thereof wherein

m and p are independently 0, 1, 2 or 3 provided that the sum of m and p is at least 2;

Q is NR<sup>1</sup>, O, S, S(O) or S(O)<sub>2</sub>;

- 10 R<sup>1</sup> is



-C(=NR<sup>8b</sup>)R<sup>8c</sup>, -SO<sub>2</sub>R<sup>8c</sup>, -OC(O)CCl<sub>3</sub>, -C(=S)R<sup>8c</sup>, optionally substituted aryl,

optionally substituted heteroaryl, optionally substituted heterocyclo,

perfluoroalkyl, cyano, hydroxy, optionally substituted alkoxy, optionally

- 20 substituted aryloxy, optionally substituted heteroaryloxy, optionally

substituted alkyl, optionally substituted cycloalkyl, optionally substituted

alkenyl, or optionally substituted alkynyl;


R<sup>2</sup> is heteroaryl, (heteroaryl)alkyl, aryl, (aryl)alkyl, heterocyclo, (heterocyclo)alkyl,

alkyl or cycloalkyl, any of which may be optionally independently substituted

- 25 with one or more groups T<sup>1</sup>, T<sup>2</sup> or T<sup>3</sup>;

J is a bond, C<sub>1-4</sub> alkylene optionally independently substituted with one or more groups T<sup>1a</sup>, T<sup>2a</sup> or T<sup>3a</sup>, or C<sub>1-4</sub> alkenylene optionally independently substituted with one or more groups T<sup>1a</sup>, T<sup>2a</sup> or T<sup>3a</sup>;

R<sup>3</sup> is




R<sup>4</sup> is H, alkyl, haloalkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, (aryl)alkyl or heteroaryl any of which may be optionally independently substituted with one or more groups T<sup>1b</sup>, T<sup>2b</sup> or T<sup>3b</sup>;

R<sup>5</sup> is

- (a) -NR<sup>6a</sup>R<sup>7a</sup>, cyano or
- (b) heteroaryl, (heteroaryl)alkyl, aryl, (aryl)alkyl, alkyl, cycloalkyl, (cycloalkyl)alkyl, heterocyclo, (heterocyclo)alkyl, or alkyl any of which may be optionally independently substituted with one or more groups T<sup>1c</sup>, T<sup>2c</sup> or T<sup>3c</sup>;

R<sup>6</sup>, R<sup>6a</sup>, R<sup>7</sup>, R<sup>7a</sup>, R<sup>8</sup>, R<sup>8a</sup>, R<sup>8a1</sup>, R<sup>8a2</sup>, and R<sup>8a3</sup> are independently H, alkyl, hydroxy, alkoxy, aryloxy, heterocycloxy, heteroaryloxy, (hydroxy)alkyl, (alkoxy)alkyl,

- (aryloxy)alkyl, (heterocyclooxy)alkyl, (heteroaryloxy)alkyl, (cyano)alkyl, (alkenyl)alkyl, (alkynyl)alkyl, cycloalkyl, (cycloalkyl)alkyl, aryl, (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, heterocyclo, (heterocyclo)alkyl,  $-C(O)R^9$ ,  $-CO_2R^9$ ,  $-C(O)-NR^9R^{10}$ , or  $-NR^9R^{10}$  any of which may be optionally
- 5 independently substituted with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$ ;  
 or  $R^6$  and  $R^7$ , or  $R^{6a}$  and  $R^{7a}$  together with the nitrogen atom to which they are attached may combine to form a 4 to 8 membered heterocyclo ring optionally independently substituted with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$ ;  
 or one of  $R^6$  or  $R^7$ , may combine with one of  $R^8$ ,  $R^{8a}$  or  $R^9$  to form a saturated or
- 10 unsaturated 5 to 8 membered ring optionally independently substituted with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$ .  
 or one of  $R^{6a}$  or  $R^{7a}$ , may combine with  $R^{8a1}$  to form a saturated or unsaturated 5 to 8 membered ring optionally independently substituted with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$
- 15  $R^{8b}$  is H, alkyl, aryl, cyano, nitro, acyl or  $-SO_2$ (alkyl) where the alkyl and aryl groups may be optionally independently substituted with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$ ;
- $R^{8c}$  is H, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, arylalkyl, heterocyclo, heteroaryl, alkoxy or aryloxy any of which may be optionally independently substituted
- 20 with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$ ;
- $R^{8d}$  is  $R^4$ ,  $COR^4$ ,  $CO_2R^4$ ,  $SO_2R^4$ ,  $CONR^6R^7$ , or  $SO_2NR^6R^7$ ;
- $R^9$  and  $R^{10}$  are independently H, alkyl, hydroxy, alkoxy, aryloxy, heterocyclooxy, heteroaryloxy, (hydroxy)alkyl, (alkoxy)alkyl, (aryloxy)alkyl, (heterocyclooxy)alkyl, (heteroaryloxy)alkyl, cycloalkyl, (cycloalkyl)alkyl, aryl,
- 25 (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, heterocyclo, or (heterocyclo)alkyl any of which may be optionally independently substituted with one or more groups  $T^{1f}$ ,  $T^{2f}$  or  $T^{3f}$
- or  $R^9$  and  $R^{10}$  together with the nitrogen atom to which they are attached may combine to form a saturated or unsaturated ring which may be optionally independently
- 30 substituted with one or more groups  $T^{1f}$ ,  $T^{2f}$  or  $T^{3f}$ ;
- W is  $=NR^{8a1}$ ,  $=N-CO_2R^{8a1}$ ,  $=N-COR^{8a1}$ ,  $=N-CN$ ,  $=N-SO_2R^{8a1}$ , or



$X^1$  is O, S,  $\text{NR}^{8a2}$  or  $\text{CH}_2$ ;

Z,  $Z^1$  and  $Z^2$  are independently =O, =S,  $\text{NR}^{8a3}$  or =N-CN;

$\text{R}^X$  is one or more optional substituents, attached to any available ring carbon atom,

5 independently selected from  $\text{T}^{1g}$ ,  $\text{T}^{2g}$  or  $\text{T}^{3g}$ ;

$\text{T}^{1-1g}$ ,  $\text{T}^{2-2g}$ , and  $\text{T}^{3-3g}$  are each independently

(1) hydrogen or  $\text{T}^6$ , where  $\text{T}^6$  is

(i) alkyl, (hydroxy)alkyl, (alkoxy)alkyl, alkenyl, alkynyl, cycloalkyl,  
 (cycloalkyl)alkyl, cycloalkenyl, (cycloalkenyl)alkyl, aryl,  
 10 (aryl)alkyl, heterocyclo, (heterocylco)alkyl, heteroaryl, or  
 (heteroaryl)alkyl;

(ii) a group (i) which is itself substituted by one or more of the same or  
 different groups (i); or

(iii) a group (i) or (ii) which is independently substituted by one or  
 15 more (preferably 1 to 3) of the following groups (2) to (13) of the  
 definition of  $\text{T}^{1-1g}$ ,  $\text{T}^{2-2g}$  and  $\text{T}^{3-3g}$ ,

(2)  $-\text{OH}$  or  $-\text{OT}^6$ ,

(3)  $-\text{SH}$  or  $-\text{ST}^6$ ,

(4)  $-\text{C}(\text{O})_t\text{H}$ ,  $-\text{C}(\text{O})_t\text{T}^6$ , or  $-\text{O}-\text{C}(\text{O})\text{T}^6$ , where  $t$  is 1 or 2;

20 (5)  $-\text{SO}_3\text{H}$ ,  $-\text{S}(\text{O})_t\text{T}^6$ , or  $\text{S}(\text{O})_t\text{N}(\text{T}^9)\text{T}^6$ ,

(6) halo,

(7) cyano,

(8) nitro,

(9)  $-\text{T}^4-\text{NT}^7\text{T}^8$ ,

25 (10)  $-\text{T}^4-\text{N}(\text{T}^9)-\text{T}^5-\text{NT}^7\text{T}^8$ ,

(11)  $-\text{T}^4-\text{N}(\text{T}^{10})-\text{T}^5-\text{T}^6$ ,

(12)  $-\text{T}^4-\text{N}(\text{T}^{10})-\text{T}^5-\text{H}$ ,

(13) oxo,

$\text{T}^4$  and  $\text{T}^5$  are each independently

30 (1) a single bond,

(2)  $-\text{T}^{11}-\text{S}(\text{O})_t-\text{T}^{12}-$ ,

- (3)  $-T^{11}-C(O)-T^{12}-$ ,
- (4)  $-T^{11}-C(S)-T^{12}-$ ,
- (5)  $-T^{11}-O-T^{12}-$ ,
- (6)  $-T^{11}-S-T^{12}-$ ,
- 5 (7)  $-T^{11}-O-C(O)-T^{12}-$ ,
- (8)  $-T^{11}-C(O)-O-T^{12}-$ ,
- (9)  $-T^{11}-C(=NT^{9a})-T^{12}-$ , or
- (10)  $-T^{11}-C(O)-C(O)-T^{12}-$

$T^7$ ,  $T^8$ ,  $T^9$ ,  $T^{9a}$  and  $T^{10}$


- 10 (1) are each independently hydrogen or a group provided in the definition of  $T^6$ , or
- (2)  $T^7$  and  $T^8$  may together be alkylene or alkenylene, completing a 3- to 8-membered saturated or unsaturated ring together with the atoms to which they are attached, which ring is unsubstituted or substituted with one or
- 15 more groups listed in the description of  $T^{1-1g}$ ,  $T^{2-2g}$  and  $T^{3-3g}$ , or
- (3)  $T^7$  or  $T^8$ , together with  $T^9$ , may be alkylene or alkenylene completing a 3- to 8-membered saturated or unsaturated ring together with the nitrogen atoms to which they are attached, which ring is unsubstituted or substituted with one or more groups listed in the description of  $T^{1-1g}$ ,  $T^{2-2g}$
- 20 and  $T^{3-3g}$ , or
- (4)  $T^7$  and  $T^8$  or  $T^9$  and  $T^{10}$  together with the nitrogen atom to which they are attached may combine to form a group  $-N=CT^{13}T^{14}$  where  $T^{13}$  and  $T^{14}$  are each independently H or a group provided in the definition of  $T^6$ ; and

$T^{11}$  and  $T^{12}$  are each independently

- 25 (1) a single bond,
- (2) alkylene,
- (3) alkenylene, or
- (4) alkynylene;

provided said compound is other than

- 30 (i) a compound of formula I




where


$R^{1i}$  is H, alkyl, aralkyl,  $-C(O)alkyl$ ,  $-C(O)aryl$ ,  $-C(O)aralkyl$ ,  
 $-C(O)alkylene-CO_2alkyl$ ,  $-CO_2alkyl$ ,  $-CO_2alkenyl$ ,  $-CO_2aralkyl$ , or  
 $-SO_2alkyl$ ; and  
 $R^{8i}$  is H, or alkyl;

(ii) a compound of formula iia, iib or iic

(a)



(b)




where

q is 0, 1 or 2; and

$R^{1iib}$  is H or  $-C(O)_2alkyl$ ;

(c)




where

$R^{1iic}$  is H or alkyl; and

$R^{8iic}$  is H or alkyl;

5

(iii) a compound of formula iii




where

$R^{1iii}$  is -alkylene-C(O)-Ar where Ar is phenyl, alkylphenyl, xylyl, halophenyl, methoxyphenyl or thienyl; and

$R^{2iii}$  is thienyl, phenyl, halophenyl, methoxyphenyl, alkylphenyl, xylyl, or trifluoromethylphenyl;


10

(iv) a compound of formula iv



15

where



$T^{1h}$  is alkyl, aryl, heteroaryl, (aryl)alkyl, (heteroaryl)alkyl,

(cycloalkyl)alkyl, -alkylene-M-alkyl, -alkylene-M-alkylene-aryl,

-alkylene-M-alkylene-heteroaryl, -alkylene-M-alkylene-cycloalkyl

20

where alkyl and alkylene groups may be optionally substituted with 1 to 5 halo,  $-S(O)_qH$ ,  $-S(O)_qalkyl$ , or 1 to 3 hydroxy, alkoxy, carboxyl or  $-C(O)Oalkyl$ , and where the aryl and heteroaryl groups may be optionally substituted with phenyl, phenoxy, (aryl)alkoxy, (heteroaryl)alkoxy, halophenyl, 1 to 3 halo or alkyl, 1 to 3 methoxy, alkoxy, or cycloalkoxy, 1 to 3 trifluoromethyl or trifluoromethoxy, 1 to 2 methylenedioxy,  $S(O)_qR^{5b}$ , nitro,  $-NR^{5b}R^{5b}$ ,  $-NR^{5b}C(O)R^{5b}$ ,  $-C(O)OR^{5b}$ ,  $-C(O)NR^{5b}R^{5b}$ ,  $-SO_2NR^{5b}R^{5b}$ ,  $-NR^{5b}SO_2aryl$ ,  $-NR^{5b}SO_2heteroaryl$ , or  $-NR^{5b}SO_2R^{5b}$ ;

M is O,  $SO_q$ ,  $NR^{5b}C(O)$ ,  $C(O)NR^{5b}$ ,  $OC(O)$ ,  $C(O)O$ ,  $-CR^{5b}=CR^{5b}-$ , or  $-C\equiv C-$ ;

$T^{2h}$  is H, alkyl, or cycloalkyl;

or  $T^{1h}$  and  $T^{2h}$  combine to form a 3 to 8-membered ring optionally including heteroatoms;

$T^{3h}$  is  $-LC(O)ANR^{6d}R^{7d}$ ;

L is  $NR^{5c}$ , O or  $CH_2$ ,

$R^{5c}$  is H, alkyl or  $R^{5c}$  combines with  $T^{2h}$  to form a 3 to 8-membered ring;

$R^{6d}$  and  $R^{7d}$  are independently H, alkyl, or substituted alkyl where the

substituents are selected from phenyl, phenoxy, 2-furyl,

$-C(O)Oalkyl$ ,  $-S(O)_qalkyl$ , 1 to 5 halo, 1 to 3 hydroxy, 1 to 3

$-OC(O)alkyl$ , or 1 to 3 alkoxy, or  $R^{6d}$  and  $R^{7d}$  may combine to form an unsaturated ring optionally containing additional heteroatoms;

A is a bond or  $-Q^1-\left(CH_2\right)_r-\overset{\overset{R^{15}}{|}}{\underset{\underset{R^{15a}}{|}}{C}}-\left(CH_2\right)_s-$  where r and s are independently 0-3;

$Q^1$  is a bond,  $NR^{5b}$  or O;

$R^{15}$  and  $R^{15a}$  are independently H, alkyl, trifluoromethyl, phenyl or optionally substituted alkyl where the substituents are selected

from imidazolyl, phenyl, indolyl, p-hydroxyphenyl,  $OR^{5b}$ ,

$S(O)_qR^{5b}$ ,  $C(O)OR^{5b}$ , cycloalkyl,  $NR^{5b}R^{5b}$ ,  $C(O)NR^{5b}R^{5b}$ , or  $R^{15}$

and R<sup>15a</sup> can independently be joined to one or both of R<sup>6d</sup> and R<sup>7d</sup> to form alkylene bridges between the terminal nitrogen and the alkyl portion of the R<sup>15</sup> and R<sup>15a</sup> group, or R<sup>15</sup> and R<sup>15a</sup> may combine to form a 3 to 7-membered ring;

5 R<sup>2iv</sup> is alkyl, cycloalkyl, aryl or heteroaryl, each optionally substituted with one to three halo, methyl, methoxy or trifluoromethyl;

$-J^{iv}-R^{3iv}$  is  $-OR^{5b}$ ,  $-C(O)OR^{5c}$ ,  $-NR^{6b}R^{7b}$ ,  $-\overset{\overset{O}{\parallel}}{C}NR^{6c}R^{7c}$  or  $-J^{iva}-R^{3iva}$ ,

where

10 J<sup>iva</sup> is a bond or alkylene optionally substituted with one or more halo or hydroxy;

R<sup>3iva</sup> is a 5 or 6-membered cycloalkyl or heterocyclo ring to which is fused an optionally substituted 5 or 6-membered aryl or heteroaryl ring where the optional substituents are selected from halo, methyl, methoxy or trifluoromethyl;

15 R<sup>5b</sup> is H, alkyl or cycloalkyl;

R<sup>5c</sup> is H, alkyl, haloalkyl, cycloalkyl, aryl, heteroaryl, (aryl)alkyl, or (heteroaryl)alkyl where the aryl and heteroaryl groups are optionally substituted with 1 to 3 halo, methyl, methoxy or trifluoromethyl;

20 R<sup>6b</sup> is  $-C(O)-R^{12a}$  or  $-SO_2-R^{12a}$ ;

R<sup>7b</sup> is aryl or heteroaryl, each optionally substituted with 1 to 3 halo, methyl, methoxy or trifluoromethyl;


25 R<sup>6c</sup> is H, alkyl, phenyl, thiazolyl, imidazolyl, furyl or thienyl each optionally substituted with 1 to 3 halo, methyl, methoxy, trifluoromethyl or trifluoromethoxy;

R<sup>7c</sup> is H or alkyl optionally substituted with phenyl, phenoxy,  $-C(O)Oalkyl$ ,  $-SO_qalkyl$ , 1 to 5 halo, 1 to 3 hydroxy, 1 to 3 alkoxy, or 1 to 3  $-OC(O)alkyl$

30 or R<sup>6c</sup> and R<sup>7c</sup> may combine to form a cycloalkyl or heterocyclo ring optionally substituted with alkyl or cycloalkyl;

$R^{12a}$  is 4-morpholinyl, 4-(1-methylpiperazinyl), cycloalkyl or alkyl, each optionally substituted with 1 to 3 fluorine, hydroxy, methoxy, trifluoromethoxy, trifluoromethyl or cycloalkyl;

5 (v) a compound of formula v



where

$R^{1v}$  is H, (alkoxy)alkyl, aryl, (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, heterocyclo, (heterocyclo)alkyl, cycloalkyl, or (cycloalkyl)alkyl

10 wherein the alkyl moieties may be optionally substituted with 1 to 7 fluoroines, and wherein the aryl, heteroaryl and heterocyclo moieties may be optionally independently substituted with 1 to 3 halo, alkyl, fluoralkyl, phenyl, benzyl, hydroxy, acetyl, amino, cyano, nitro, alkoxy, (alkyl)amino and (dialkyl)amino

15  $T^1$  is hydroxy, (hydroxy)alkyl, (alkoxy)alkyl,  $\text{NHSO}_2R^{16}$ ,  $\text{C(OH)R}^{16}R^{17}$ , halo, heteroaryl, or  $\text{C(O)NHR}^{16}$ ;

$T^2$  is H, halo or alkyl

$R^{16}$  and  $R^{17}$  are independently H, alkyl, alkoxy and (alkoxy)alkyl

20 wherein the alkyl moieties are optionally substituted with 1 to 7 fluorine atoms;

$n$  is 0 or 1;


$E$  and  $F$  are independently O, N, S or CH provided  $E$  and  $F$  cannot both be either O or S; and

25  $T^{1c}$  is H, aryl, halo, heteroaryl, heterocyclo,  $-\text{SO}_2R^{18}$ ,  $-\text{C(O)R}^{18}$ ,  $-\text{C(O)NR}^{18}R^{19}$ ,  $-\text{COOR}^{18}$ , or  $-\text{C(OH)R}^{18}R^{19}$ ;

$R^{18}$  and  $R^{19}$  are independently H, (alkoxy)alkyl, aryl, (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, heterocyclo, (heterocyclo)alkyl,

5 cycloalkyl, or (cycloalkyl)alkyl wherein the alkyl moieties may be optionally substituted with 1 to 7 fluoroines, and wherein the aryl, heteroaryl and heterocyclo moieties may be optionally independently substituted with 1 to 3 halo, alkyl, fluoralkyl, phenyl, benzyl, hydroxy, acetyl, amino, cyano, nitro, alkoxy, (alkyl)amino and (dialkyl)amino;

(vi) a compound of formula vi




10

where

$R^{1vi}$  is alkyl optionally substituted with hydroxy or  $-OC(O)alkyl$ ; and  $R^{20}$  is H or  $-C(O)alkyl$ ;

(vii) a compound of formula vii



15


where

$R^{1vii}$  is (aryl)alkyl, (aryl)alkenyl, (alkoxy)alkyl, (aryloxy)alkyl, (arylalkoxy)alkyl, (cycloalkyloxy)alkyl, (heterocyclo)alkyl, or  $-X-CH_2-Y$ -phenyl, where X is  $-CH_2-$  or  $-C(O)-$ , and Y is  $-C(O)-$ ,  $-C(=NOH)-$  or  $-CH_2-$  optionally substituted with hydroxy;

20  $R^{21}$  is  $-C(O)alkyl$ ,  $-C(O)aryl$ ,  $-C(O)Oalkyl$ ,  $-O-C(O)alkyl$ , or  $-C(O)NR^{6e}R^{7e}$ , where  $R^{6e}$  and  $R^{7e}$  are independently alkyl or combine to form an alkylene chain; and


$T^{1vii}$  is H, hydroxy or alkoxy;

(viii) a compound of formula viii



where

5  $R^{lviii}$  is  $SO_2R^{23}$  or a group




$R^{22}$  is H, alkyl, phenyl, benzyl or biphenyl, wherein alkyl is optionally substituted with biphenyl, carboxy, alkoxy,  $-C(O)alkyl$ , or  $-C(O)N(H)alkyl$ , and wherein benzyl is optionally substituted with hydroxy, alkoxy or halo;

$R^{23}$  is alkyl or phenyl, wherein phenyl is optionally substituted with 1 or 2 halo, methoxy, halomethoxy,  $-N(H)C(O)methyl$ ,  $CF_3$ , alkyl or CN;


$X^{viii}$  is a bond,  $CH_2$  or  $CHMe$ ;

15  $R^{24}$  is H or  $-N(H)R^{25}$ ; and

$R^{25}$  is



20 (ix) a compound of formula ixa or ixb



where

$R^{1x}$  is H or alkyl;

$R^{2ixa}$  is heteroaryl or (heteroaryl)alkyl;

5  $R^{2ixb}$  is alkyl optionally substituted with  $C(O)_tH$  or  $C(O)_t$ alkyl where t is 1 or 2;


$R^{26a}$  is alkyl optionally substituted with  $C(O)_tH$  or  $C(O)_t$ alkyl;

$R^{26b}$  is optionally substituted heteroaryl,  $-NHR^{27}$ , or  $-OR^{27}$ ; and

$R^{27}$  is optionally substituted heteroaryl;

10

(x) a compound of formula x



where

$R^{1x}$  is  $-C(O)-(CHR^{28})-NHR^{29}$ ;

15  $R^{2x}$  is phenyl, (phenyl)alkyl, tetrahydropyranyl, piperidyl, alkyl, cycloalkyl or (cycloalkyl)alkyl;

$R^{26c}$  is  $-C(O)R^{30}$ ,  $-C(O)_2$ alkyl,  $-CH_2-R^{31}$ , or  $-NR^{32}R^{33}$ ;

$R^{28}$  is H, alkyl, cycloalkyl, (cycloalkyl)alkyl, aryl, (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, aryloxy, (aryloxy)alkyl, aralkoxy or

20 (aralkoxy)alkyl, any of which may be optionally substituted;


$R^{29}$  is H or  $-C(O)-(CR^{34}R^{34})_s$ -het\* where s is 0-2;  $R^{34}$  is H, alkyl, cycloalkyl, (cycloalkyl)alkyl, heteroaryl or (heteroaryl)alkyl; and het\* is an optionally substituted N-containing heterocyclo ring fused to an optionally substituted aryl, heteroaryl, heteroclo or cycloalkyl ring;

25


$R^{30}$  is alkyl, heterocyclo,  $-NR^{35}R^{36}$ , where  $R^{35}$  and  $R^{36}$  are independently H, alkyl, cycloalkyl or (cycloalkyl)alkyl;

R<sup>31</sup> is -S-alkyl, -S(O)-alkyl, -S(O)<sub>2</sub>-alkyl, -O-alkyl, -NR<sup>32</sup>R<sup>33</sup>, or triazolyl;  
R<sup>32</sup> and R<sup>33</sup> are independently H, alkyl, -S(O)<sub>2</sub>alkyl, or -C(O)alkyl;



5 (xi) a compound of formula xi





where

 $R^{1 \times i}$  is

10




15

 $R^{2xi}$  is

20

 $R^{3xi}$  is



R<sup>34</sup> is phenyl or naphthyl optionally substituted with cyano or –  
 5 C(O)NH<sub>2</sub>;

R<sup>35</sup> is H or alkyl;

R<sup>36</sup> is phenyl optionally substituted with alkoxy;

R<sup>37</sup> is H, alkyl, or –C(O)NH<sub>2</sub>;


10 T<sup>1xi</sup> is H, alkyl, alkoxy, haloalkoxy, cyano, nitro, –CH<sub>2</sub>NH<sub>2</sub>,  
 –CH<sub>2</sub>N(H)–(C(O)OCH<sub>2</sub>–phenyl), or –NH–SO<sub>2</sub>alkyl;

T<sup>1axi</sup> is H, alkyl, halo, alkoxy, haloalkoxy, or cyano;

T<sup>1bxi</sup> is H, alkyl, halo alkoxy or hydroxy;

T<sup>1cxi</sup> is H or alkyl;

15 (xii) a compound of formula xii





where

R<sup>38</sup> is H or methyl;

R<sup>39</sup> is

20





$R^{40}$  is H or Cl; and

$v$  is 0 or 1.

5


2. A compound of claim 1 wherein

$Q$  is  $NR^1$  or O;

$R^1$  is H,  $-\overset{\overset{O}{\parallel}}{C}-R^4$ ,  $-\overset{\overset{O}{\parallel}}{C}-O-R^4$ ,  $-\overset{\overset{O}{\parallel}}{S}-NR^6R^7$ ,  $-\text{SO}_2R^{8c}$ ,  $-\overset{\overset{O}{\parallel}}{C}-NR^6R^7$ ,  $-\overset{\overset{O}{\parallel}}{C}-NR^6R^7$ ,  
 $-\text{C}(=\text{S})R^{8c}$ ,  $-\text{C}(=\text{NR}^{8b})R^{8c}$  or heteroaryl;

10  $R^2$  is aryl, (aryl)alkyl or heteroaryl any of which may be optionally independently substituted with one or more groups  $T^1$ ,  $T^2$  or  $T^3$ ;

$J$  is a bond or methylene; and



15 3. A compound of claim 2 wherein

$Q$  is  $NR^1$ ;

$R^4$  is alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, (aryl)alkyl, heteroaryl or (heteroaryl)alkyl any of which may be optionally independently substituted with one or more  $T^{1b}$ ,  $T^{2b}$   $T^{3b}$ ;

20  $R^5$  is

(a)  $-\text{NR}^{6a}R^{7a}$  or

(b) aryl, (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, heterocyclo or (heterocyclo)alkyl any of which may be optionally independently substituted with one or more  $T^{1c}$ ,  $T^{2c}$   $T^{3c}$ ;

25  $R^6$ ,  $R^{6a}$ ,  $R^7$  and  $R^{7a}$  are independently H, alkyl, alkenyl, alkynyl, aryl, (aryl)alkyl, (alkoxy)alkyl, cycloalkyl, (cycloalkyl)alkyl, (hydroxy)alkyl, heteroaryl, (heteroaryl)alkyl, heterocyclo, (heterocyclo)alkyl, (aryloxy)alkyl,  $-\text{C}(\text{O})R^9$ ,

$-\text{CO}_2\text{R}^9$ , or  $-\text{C}(\text{O})-\text{NR}^9\text{R}^{10}$  any of which may be optionally independently substituted with one or more  $\text{T}^{1d}$ ,  $\text{T}^{2d}$   $\text{T}^{3d}$ ;

or  $\text{R}^6$  and  $\text{R}^7$ , or  $\text{R}^{6a}$  and  $\text{R}^{7a}$  together with the nitrogen atom to which they are attached combine to form an optionally substituted 4 to 8 membered

5 heterocyclo ring optionally independently substituted with one or more groups  $\text{T}^{1d}$ ,  $\text{T}^{2d}$  or  $\text{T}^{3d}$ ;

$\text{R}^{8a}$  is H, alkyl, or (aryl)alkyl where the alkyl and aryl groups may be optionally independently substituted with one or more groups  $\text{T}^{1d}$ ,  $\text{T}^{2d}$  or  $\text{T}^{3d}$ ;

$\text{R}^{8c}$  is

10 (a) alkyl, aryl, heteroaryl any of which may be optionally independently substituted with one or more  $\text{T}^{1d}$ ,  $\text{T}^{2d}$   $\text{T}^{3d}$ ; or

(b)  $-\text{NR}^9\text{R}^{10}$ ;

W is  $=\text{N}-\text{CN}$ ; and


$\text{Z}^1$  is  $=\text{O}$  or  $=\text{N}-\text{CN}$ ;

15


4. A compound of claim 3 wherein  $\text{T}^1$ ,  $\text{T}^{1b}$ ,  $\text{T}^{1c}$ ,  $\text{T}^{1d}$ ,  $\text{T}^2$ ,  $\text{T}^{2b}$ ,  $\text{T}^{2c}$ ,  $\text{T}^{2d}$ ,  $\text{T}^3$ ,  $\text{T}^{3b}$ ,  $\text{T}^{3c}$  and  $\text{T}^{3d}$  are independently halo, cyano, alkyl, aryl, (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, haloalkyl,  $-\text{OH}$ ,  $-\text{OT}^6$ ,  $-\text{C}(\text{O})\text{T}^6$ ,  $-\text{SO}_2\text{T}^6$ ,  $-\text{T}^4\text{NT}^7\text{T}^8$ , or  $-\text{T}^4\text{N}(\text{T}^{10})\text{T}^5-\text{T}^6$ .

20



5. A compound of claim 1 having the structure



where  $\text{R}^{1*}$  is




25




- 5  $\text{---C(=S)R}^{8c}$ , optionally substituted heterocyclo, cyano, hydroxy, optionally substituted alkoxy, optionally substituted aryloxy, or optionally substituted heteroaryloxy.

6. The compound of claim 5 where at least one of  $\text{T}^{1c}$ ,  $\text{T}^{2c}$  or  $\text{T}^{3c}$  is alkoxy.

- 10 7. A compound of claim 1 having the structure




where  $\text{R}^{3*}$  is



- 15  $\begin{array}{c} \text{Z}^{1*} \\ \parallel \\ \text{---N---H} \end{array}$   $\text{NR}^{6a}\text{R}^{7a}$ ,  $\begin{array}{c} \text{O} \\ \parallel \\ \text{---N---S---} \end{array}$   $\text{NR}^{6a}\text{R}^{7a}$ ; and

$\text{Z}^{1*}$  is  $=\text{O}$  or  $=\text{N-CN}$ .


8. A compound of claim 1 having the structure



where  $R^{1**}$  is  $-C(O)R^{4*}$ ,  $-SO_2R^{8c*}$ ;


- 5  $R^{4*}$  is (aryl)alkyl, (aryloxy)alkyl, or (aryl)cycloalkyl any of which may be optionally independently substituted with one or more groups  $T^{1b}$ ,  $T^{2b}$  or  $T^{3b}$ ; and  $R^{8c*}$  is aryl optionally independently substituted with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$ .

9. A compound of claim 1 having the structure



10

10. A compound of claim 1 having the structure




- 15 where  $R^{3**}$  is heteroaryl or (heteroaryl)alkyl, either of which may be optionally substituted with one or more groups  $T^{1c}$ ,  $T^{2c}$  or  $T^{3c}$ .

11. A pharmaceutical composition comprising at least one compound of claim 1 together with a suitable vehicle or carrier therefor.

12. A pharmaceutical composition of claim 11 further comprising at least one additional therapeutic agent selected from anti-arrhythmic agents, calcium channel blockers, anti-platelet agents, anti-hypertensive agents, anti-thrombotic/anti-thrombolytic agents, anti-coagulants, HMG-CoA reductase inhibitors, anti-diabetic agents, thyroid mimetics, mineralocorticoid receptor antagonists, or cardiac glycosides.
13. The pharmaceutical composition of claim 12 wherein
- (a) the additional anti-arrhythmic agent is selected from sotalol, dofetilide, diltiazem and verapamil;
  - (b) the anti-platelet agent is selected from clopidogrel, ifetroban and aspirin;
  - (c) the anti-hypertensive agent is selected from beta adrenergic blockers, ACE inhibitors, A II antagonists, ET antagonists, Dual ET/A II antagonists, and vasopepsidase inhibitors;
  - (d) the anti-thrombotic/anti-thrombolytic agent is selected from tPA, recombinant tPA, TNK, nPA, factor VIIa inhibitors, factor Xa inhibitors and thrombin inhibitors;
  - (e) the anti-coagulant is selected from warfarin and heparins;
  - (f) the HMG-CoA reductase inhibitor is selected from pravastatin, lovastatin, atorvastatin, simvastatin, NK-104 and ZD-4522;
  - (g) the anti-diabetic agent is selected from biguanides and biguanide/glyburide combinations;
  - (h) the mineralocorticoid receptor antagonist is selected from spironolactone and eplerinone; and
  - (i) the cardiac glycoside is selected from digitalis and ouabain.
14. The pharmaceutical composition of claim 13 wherein
- (a) the ACE inhibitors are selected from captopril, zofenopril, fosinopril, enalapril, ceranopril, cilazopril, delapril, pentopril, quinapril, ramipril, and lisinopril; and




- (b) the vasopepsidase inhibitors are selected from omapatrilat and gemopatrilat.

15. A method of treating  $I_{Kur}$ -associated disorders comprising the step of  
 5 administering to a patient in need thereof an effective amount of at least one compound of the following formula I



enantiomers, diastereomers, salts and solvates thereof wherein

- 10 m and p are independently 0, 1, 2 or 3 provided that the sum of m and p is at least 2;  
 Q is  $NR^1$ , O, S, S(O) or S(O)<sub>2</sub>;  
 $R^1$  is




- 20  $-C(=NR^{8b})R^{8c}$ ,  $-SO_2R^{8c}$ ,  $-OC(O)CCl_3$ ,  $-C(=S)R^{8c}$ , optionally substituted aryl,  
 optionally substituted heteroaryl, optionally substituted heterocyclo,  
 perfluoroalkyl, cyano, hydroxy, optionally substituted alkoxy, optionally  
 substituted aryloxy, optionally substituted heteroaryloxy, optionally  
 substituted alkyl, optionally substituted cycloalkyl, optionally substituted  
 alkenyl, or optionally substituted alkynyl;

$R^2$  is heteroaryl, (heteroaryl)alkyl, aryl, (aryl)alkyl, heterocyclo, (heterocyclo)alkyl, alkyl or cycloalkyl, any of which may be optionally independently substituted with one or more groups  $T^1$ ,  $T^2$  or  $T^3$ ;

J is a bond,  $C_{1-4}$  alkylene optionally independently substituted with one or more groups  $T^{1a}$ ,  $T^{2a}$  or  $T^{3a}$ , or  $C_{1-4}$  alkenylene optionally independently substituted with one or more groups  $T^{1a}$ ,  $T^{2a}$  or  $T^{3a}$ ;

$R^3$  is



15

$R^4$  is H, alkyl, haloalkyl, alkenyl, alkynyl, cycloalkyl, heterocyclo, aryl, (aryl)alkyl or heteroaryl any of which may be optionally independently substituted with one or more groups  $T^{1b}$ ,  $T^{2b}$  or  $T^{3b}$ ;


$R^5$  is

- 20 (a)  $-\text{NR}^{6a}\text{R}^{7a}$ , cyano or  
 (b) heteroaryl, (heteroaryl)alkyl, aryl, (aryl)alkyl, alkyl, cycloalkyl, (cycloalkyl)alkyl, heterocyclo, (heterocyclo)alkyl, or alkyl any of which

- may be optionally independently substituted with one or more groups  $T^{1c}$ ,  $T^{2c}$  or  $T^{3c}$ ;
- $R^6$ ,  $R^{6a}$ ,  $R^7$ ,  $R^{7a}$ ,  $R^8$ ,  $R^{8a}$ ,  $R^{8a1}$ ,  $R^{8a2}$ , and  $R^{8a3}$  are independently H, alkyl, hydroxy, alkoxy, aryloxy, heterocyclooxy, heteroaryloxy, (hydroxy)alkyl, (alkoxy)alkyl, (aryloxy)alkyl, (heterocyclooxy)alkyl, (heteroaryloxy)alkyl, (cyano)alkyl, (alkenyl)alkyl, (alkynyl)alkyl, cycloalkyl, (cycloalkyl)alkyl, aryl, (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, heterocyclo, (heterocyclo)alkyl,  $-C(O)R^9$ ,  $-CO_2R^9$ ,  $-C(O)-NR^9R^{10}$ , or  $-NR^9R^{10}$  any of which may be optionally independently substituted with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$ ;
- or  $R^6$  and  $R^7$ , or  $R^{6a}$  and  $R^{7a}$  together with the nitrogen atom to which they are attached may combine to form a 4 to 8 membered heterocyclo ring optionally independently substituted with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$ ;
- or one of  $R^6$  or  $R^7$ , may combine with one of  $R^8$ ,  $R^{8a}$  or  $R^9$  to form a saturated or unsaturated 5 to 8 membered ring optionally independently substituted with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$ ;
- or one of  $R^{6a}$  or  $R^{7a}$ , may combine with  $R^{8a1}$  to form a saturated or unsaturated 5 to 8 membered ring optionally independently substituted with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$ ;
- $R^{8b}$  is H, alkyl, aryl, cyano, nitro, acyl or  $-SO_2$ (alkyl) where the alkyl and aryl groups may be optionally independently substituted with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$ ;
- $R^{8c}$  is H, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, arylalkyl, heterocyclo, heteroaryl, alkoxy or aryloxy any of which may be optionally independently substituted with one or more groups  $T^{1d}$ ,  $T^{2d}$  or  $T^{3d}$ ;
- $R^{8d}$  is  $R^4$ ,  $COR^4$ ,  $CO_2R^4$ ,  $SO_2R^4$ ,  $CONR^6R^7$ , or  $SO_2NR^6R^7$ ;
- $R^9$  and  $R^{10}$  are independently H, alkyl, hydroxy, alkoxy, aryloxy, heterocyclooxy, heteroaryloxy, (hydroxy)alkyl, (alkoxy)alkyl, (aryloxy)alkyl, (heterocyclooxy)alkyl, (heteroaryloxy)alkyl, cycloalkyl, (cycloalkyl)alkyl, aryl, (aryl)alkyl, heteroaryl, (heteroaryl)alkyl, heterocyclo, or (heterocyclo)alkyl any of which may be optionally independently substituted with one or more groups  $T^{1f}$ ,  $T^{2f}$  or  $T^{3f}$

or R<sup>9</sup> and R<sup>10</sup> together with the nitrogen atom to which they are attached may combine to form a saturated or unsaturated ring which may be optionally independently substituted with one or more groups T<sup>1f</sup>, T<sup>2f</sup> or T<sup>3f</sup>;

5 W is =NR<sup>8a1</sup>, =N-CO<sub>2</sub>R<sup>8a1</sup>, =N-COR<sup>8a1</sup>, =N-CN, =N-SO<sub>2</sub>R<sup>8a1</sup>, or



X<sup>1</sup> is O, S, NR<sup>8a2</sup> or CH<sub>2</sub>;

Z, Z<sup>1</sup> and Z<sup>2</sup> are independently =O, =S, =NR<sup>8a3</sup> or =N-CN;

R<sup>X</sup> is one or more optional substituents, attached to any available ring carbon atom, independently selected from T<sup>1g</sup>, T<sup>2g</sup> or T<sup>3g</sup>;

10 T<sup>1-1g</sup>, T<sup>2-2g</sup>, and T<sup>3-3g</sup> are each independently

(1) hydrogen or T<sup>6</sup>, where T<sup>6</sup> is

(i) alkyl, (hydroxy)alkyl, (alkoxy)alkyl, alkenyl, alkynyl, cycloalkyl, (cycloalkyl)alkyl, cycloalkenyl, (cycloalkenyl)alkyl, aryl, (aryl)alkyl, heterocyclo, (heterocylco)alkyl, heteroaryl, or (heteroaryl)alkyl;

(ii) a group (i) which is itself substituted by one or more of the same or different groups (i); or

(iii) a group (i) or (ii) which is independently substituted by one or more (preferably 1 to 3) of the following groups (2) to (13) of the definition of T<sup>1-1g</sup>, T<sup>2-2g</sup> and T<sup>3-3g</sup>,

20

(2) -OH or -OT<sup>6</sup>,

(3) -SH or -ST<sup>6</sup>,

(4) -C(O)<sub>t</sub>H, -C(O)<sub>t</sub>T<sup>6</sup>, or -O-C(O)T<sup>6</sup>, where t is 1 or 2;

(5) -SO<sub>3</sub>H, -S(O)<sub>t</sub>T<sup>6</sup>, or S(O)<sub>t</sub>N(T<sup>9</sup>)T<sup>6</sup>,

25

(6) halo,

(7) cyano,

(8) nitro,

(9) -T<sup>4</sup>-NT<sup>7</sup>T<sup>8</sup>,

(10) -T<sup>4</sup>-N(T<sup>9</sup>)-T<sup>5</sup>-NT<sup>7</sup>T<sup>8</sup>,

30

(11) -T<sup>4</sup>-N(T<sup>10</sup>)-T<sup>5</sup>-T<sup>6</sup>,

(12) -T<sup>4</sup>-N(T<sup>10</sup>)-T<sup>5</sup>-H,

(13) oxo,

T<sup>4</sup> and T<sup>5</sup> are each independently

- (1) a single bond,
- (2) -T<sup>11</sup>-S(O)<sub>r</sub>-T<sup>12</sup>-,
- 5 (3) -T<sup>11</sup>-C(O)-T<sup>12</sup>-,
- (4) -T<sup>11</sup>-C(S)-T<sup>12</sup>-,
- (5) -T<sup>11</sup>-O-T<sup>12</sup>-,
- (6) -T<sup>11</sup>-S-T<sup>12</sup>-,
- (7) -T<sup>11</sup>-O-C(O)-T<sup>12</sup>-,
- 10 (8) -T<sup>11</sup>-C(O)-O-T<sup>12</sup>-,
- (9) -T<sup>11</sup>-C(=NT<sup>9a</sup>)-T<sup>12</sup>-, or
- (10) -T<sup>11</sup>-C(O)-C(O)-T<sup>12</sup>-

T<sup>7</sup>, T<sup>8</sup>, T<sup>9</sup>, T<sup>9a</sup> and T<sup>10</sup>


- (1) are each independently hydrogen or a group provided in the definition of  
15 T<sup>6</sup>, or
- (2) T<sup>7</sup> and T<sup>8</sup> may together be alkylene or alkenylene, completing a 3- to 8-  
membered saturated or unsaturated ring together with the atoms to which  
they are attached, which ring is unsubstituted or substituted with one or  
more groups listed in the description of T<sup>1-1g</sup>, T<sup>2-2g</sup> and T<sup>3-3g</sup>, or
- 20 (3) T<sup>7</sup> or T<sup>8</sup>, together with T<sup>9</sup>, may be alkylene or alkenylene completing a  
3- to 8-membered saturated or unsaturated ring together with the  
nitrogen atoms to which they are attached, which ring is unsubstituted or  
substituted with one or more groups listed in the description of T<sup>1-1g</sup>, T<sup>2-  
2g</sup> and T<sup>3-3g</sup>, or
- 25 (4) T<sup>7</sup> and T<sup>8</sup> or T<sup>9</sup> and T<sup>10</sup> together with the nitrogen atom to which they are  
attached may combine to form a group -N=CT<sup>13</sup>T<sup>14</sup> where T<sup>13</sup> and T<sup>14</sup>  
are each independently H or a group provided in the definition of T<sup>6</sup>; and

T<sup>11</sup> and T<sup>12</sup> are each independently

- (1) a single bond,
- 30 (2) alkylene,
- (3) alkenylene, or
- (4) alkynylene;

provided said compound is other than

(i) a compound of formula i



where

- 5             $R^{1i}$  is H, alkyl, aralkyl, -C(O)alkyl, -C(O)aryl, -C(O)aralkyl, -C(O)alkylene-CO<sub>2</sub>alkyl, -CO<sub>2</sub>alkyl, -CO<sub>2</sub>alkenyl, -CO<sub>2</sub>aralkyl, or -SO<sub>2</sub>alkyl; and
- $R^{8i}$  is H, or alkyl.

- 10            16. The method of claim 15 wherein the  $I_{Kur}$ -associated condition is arrhythmia.
- 15            17. The method of claim 16 wherein the arrhythmia is a supraventricular arrhythmia.
18. The method of claim 17 wherein the supraventricular arrhythmia is atrial fibrillation.
- 20            19. The method of claim 17 wherein the supraventricular arrhythmia is atrial flutter.
20. The method of claim 15 wherein the  $I_{Kur}$ -associated condition is a gastrointestinal disorder.
- 25            21. The method of claim 20 wherein the gastrointestinal disorder is reflux esophagitis.

22. The method of claim 20 wherein the gastrointestinal disorder is a motility disorder.
23. The method of claim 20 wherein the  $I_{Kur}$ -associated condition is an inflammatory or immunological disease.
24. The method of claim 23 wherein the inflammatory disease is chronic obstructive pulmonary disease.
25. A method of treating diabetes comprising administering to a patient in need thereof an effective amount of at least one compound of claim 1.
26. A method of treating cognitive disorders comprising administering to a patient in need thereof an effective amount of at least one compound of claim 1.
27. A method of treating migraine comprising administering to a patient in need thereof an effective amount of at least one compound of claim 1.
28. A method of treating epilepsy comprising administering to a patient in need thereof an effective amount of at least one compound of claim 1.